



## Evaluation of Inhalation Effect of Insecticides on Haemorheological Profiles of Rabbits

Emily Funmilayo Eyitayo <sup>a</sup>, Olusegun Taiwo Oke <sup>b</sup>  
and Emmanuel Ifeanyi Obeagu <sup>c\*</sup>

<sup>a</sup> Oyo State Management Board, Oyo State, Nigeria.

<sup>b</sup> Department of Medical Laboratory Science, Achievers University, Owo, Ondo State, Nigeria.

<sup>c</sup> Department of Medical Laboratory Science, Kampala International University, Western Campus, Uganda.

### Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

### Article Information

#### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/90286>

Original Research Article

Received 25 May 2022  
Accepted 01 August 2022  
Published 04 August 2022

### ABSTRACT

**Background:** High malaria burden has led to the increase use of insecticides in the tropics and subtropics. This study thus aimed at assessing the effect of insecticides inhalation on haemorheological parameters using experimental animal model.

**Materials and Methods:** Sixteen adult male rabbits divided into four groups A, B, C and D. Group A, B and C were exposed to 20ml of pyrethroid insecticides containing 0.05% pralletrin and 0.15% cyfluthrin by inhalation for 10mins, 20mins, and 30mins respectively. Group D serve as the control and was not exposed. Baseline study was done on all the animals before grouping for exposure. The pyrethroid insecticides was soaked in cotton wool (2.5g) in a container that was able to prevent the animal from ingesting it, which was placed inside the room A, B and C for 10mins, 20mins and 30mins respectively. The rabbits were exposed for three weeks and sample were collected at the end of each week. Exposure was discontinued after day 21 and samples were collected again on day 28 and 35 respectively which is the fourth and fifth week. All the animals were monitored twice daily for clinical signs like jerky movement, skin scratching, licking of legs and other body parts.

**Results:** Data analysis revealed that there was significant effect of inhalation of insecticides on some haemorheological parameters of rabbits at 10min, 20mins and 30mins of exposure. There was decreased in Plasma viscosity. But there was no significant effect on whole blood viscosity and plasma fibrinogen concentration.

\*Corresponding author: E-mail: [emmanuelobeagu@yahoo.com](mailto:emmanuelobeagu@yahoo.com);

**Conclusion:** The results from this study have shown that aerosol of these pyrethroid insecticides (pralletrin and cyfluthrin) has effect on haemorrhological parameters. It is thus recommended that one should avoid exposure to the aerosol of these insecticides during domestic, veterinary, agricultural or industrial use.

*Keywords: Inhalation; insecticides; haemorheological profiles; rabbits.*

## 1. INTRODUCTION

Excessive use of various chemicals including insecticides has become a public health concern. Use of insecticides and other organophosphates is one of the major ways through which manufacturing workers and farmers are exposed to toxicants and this has impact on the ecosystem and public health [1]. Pyrethroid insecticides are used extensively in agriculture, commercial facilities and in residential homes to control insect pests [2]. An insecticide is a natural or man-made preparation that is used to kill or control insects such as mosquitoes, cockroaches, bees, and wasp. The most common active ingredient in insecticide are synergist, carbamate, whose common name is propoxur, pyrethrin (or synthetic pyrethroids), D-trans-allevrin, permethrin, pralletrin, tetrametrin, deltametrin, cyfluthrin imiprothrin chlopyritos, Diaznon, Malathion, Silical gel, Boric acid, Arsenicals, paradichlorobenzene, Naphthalene, N,N-diethyl meta-toluamide (Deet), Dimethylphthalate [3]. Pyrethrins, pyrethroids and carbamate are effective insecticides that are often used in household sprays, insect repellent, pet shampoo, and lice treatment. They are often combined commercially with other chemicals called synergists, which enhance their insecticidal activities. Synergists are chemicals that activate some insecticides making them more poisonous to insects and thereby enhancing the effectiveness of the active ingredients. MGK 264 and piperonyl-butoxide are two commonly used synergist [4]. Synthetic pyrethroid insecticides are now used as substitutes for pest control [5] thus accounting for over 30% of insecticide used globally [6]. Cypermethrin, a pyrethroid has been documented to cause clinical sign such as increased urination, licking of legs, jerky movements ataxia, incoordination, staggering gait dizziness, altered blood chemistry, hepatotoxicity [7,8] and neutotoxicity [9].

The study was done to evaluate inhalation effect of insecticides on haemorheological profiles of rabbits.

## 2. MATERIALS AND METHODS

Sixteen (16) adult male and female rabbits weighing 1.5 - 2kg were obtained from Animal Care and Use Research Ethics Committee (ACUREC) University of Ibadan Oyo State, which served as subjects for this experiment. They were housed in a clean, quiet, well ventilated and temperature controlled room ( $21\pm 4^{\circ}$  C) in experimental animal house of University of Ibadan, Oyo State.

The rabbits were fed continuously with pelletized guinea feed containing 16.5% protein and water, they were provided with clean drinking water throughout the duration of the experiment and were allowed to acclimatized to their new environment for two (2) weeks in separate rooms and their weight was taken before and after each week.

### 2.1 Procedure for Insecticides Exposure

#### 2.1.1 Study design

The rabbits were grouped into four (4) A B C and D before acclimatization for two weeks in a separate room.

Group A consists of four animals shaved at the spine for group identification and were exposed to 20ml of pyrethroid insecticides containing 0.05% imiprothrin, 0.05% pralletrin and 0.15% cyfluthrin by inhalation for 10mins.

Group B consists of four animals shaved at the left leg for group identification and were exposed for 20mins.

Group C consists of four animals shaved at the head for group identification and were exposed for 30mins daily respectively for three weeks [10] (Akhigbe et al. 2011).

Group D was used as control and was not shaved and exposed.

The rabbits in group A, B, C i.e the test group in poorly ventilated room for exposure, were

exposed to 20mls of pyrethroid insecticides which was soaked in cotton wool (2.5g) and place inside the rooms (A, B, C) respectively, using a container that prevented the rabbits from ingesting them. The rabbits were exposed daily by inhalation for 10mins (group A) 20mins (group B) and 30mins (group C) for three weeks.

Exposure was discontinued after day 21 to day 35 [11] (Kingsley et al. 2016).

The baseline parameters (Whole blood viscosity, plasma viscosity, packed cell volume) were carried out on the sixteen rabbits before exposure. The insecticides was purchased directly from the company at 13/14 Abimbola Street Isolo Industrial Estate Isolo Nigeria (Johnson Wax Nigeria).

All animals received humane care in compliance with the guidelines of the University of Ibadan Animal Care and Use Research Ethics Committee (ACUREC).

The post exposure samples were collected on day 28 and 35 respectively after three weeks of exposure i.e. last sample was collected on the fourth and fifth weeks after the last exposure.

## 2.2 Collection of Blood Samples

### 2.2.1 Procedure

The hair at the ear vein of the rabbits was wiped with xylene and about 9ml of venous blood sample was collected from the ear vein of the rabbits in which 4.5ml of blood was dispensed into a 5ml bottle containing 0.5ml of sodium citrate and was mixed gently to avoid clotting, while the remaining 4.5ml of blood was dispensed into bottle containing EDTA and mixed to avoid clotting.

The EDTA blood was used for Whole blood and plasma Viscosity. Samples were labeled properly to avoid error. The analysis of blood samples was carried out at the Laboratory Unit, Jericho Specialist Hospital Jericho, and Ibadan.

### 2.2.2 Packed cell volume (PCV) [12]

Micro haematocrit method.

**Procedure:** The anticoagulated blood was mixed carefully and the plain capillary tubes was filled with blood up to  $\frac{3}{4}$  length of the capillary tube in which one end was sealed with plasticine, then

spinned at 12000 rpm for 5 minutes. Therefore it was read with the aid of microhaematocrit reader [13].

### 2.2.3 Total white blood cell count [12]

**Procedure:** 1 in 20 dilution of Turk's solution to blood was made by adding 0.02ml of blood to 0.38ml of Turk's fluid into a clean tube. The dilution was then charged into an improved Neubaur counting chamber with the use of pipette, the chamber was left undisturbed for 2 minutes to allow time for white cells to settle, and these cells were counted using x10 objective lens of microscope by counting the four outer square of the chamber.

### 2.2.4 Platelets count [12]

**Procedure:** 1 in 20 dilution of blood to ammonium oxalate was made by adding 0.02ml of well mixed anticoagulated blood to 0.38ml of ammonium oxalate into clean tube. This solution was then charged into improved Neubaur counting chamber and was counted under X 40 objective lens of the Microscope by counting the inner five (5) squares.

## 2.3 Plasma Fibrinogen Concentration (PFC)

### 2.3.1 Procedure

1 ml of plasma was dispensed into the test tube and 1ml of prewarmed 0.025M calcium chloride was added. The content was thoroughly mixed, and an applicator stick was inserted, so that fibrin clot can be wound around it, an7d this was incubated for 30 minutes at 37°C in a water bath. When all the fibrin has wound round the applicator stick, the adhered fibrin was removed from the the applicator stick after washing 3 – 4 times with distilled water and then blot dried, it was carefully removed with whatman filter paper no1. The adherent fibrin was carefully removed from the stick into a clean petridish for 3 – 4 days at room temperature to dry. The dried fibrin was weighed, and the plasma fibrinogen concentration was calculated using the formular.

## 2.4 Plasma Viscosity and Whole Blood Viscosity

### 2.4.1 Procedure

A syringe (1ml) was clamped in a retort stand vertically. The blood /plasma was drawn up and

**Table 1. Comparison of the haemorrhological parameters of those exposed in Group A (10 mins) with those not exposed in Group D (Controls) using paired sample t-test (n=8)**

Haemorrhological parameters	Group A (n=4)	Controls (n=4)	t-test	p-value
Plasma viscosity	3.36 ± 0.31	3.65 ± 0.15	-2.998	0.012*
Whole Blood viscosity	228.50 ± 2.31	229.58 ± 0.79	-1.384	0.194
Plasma Fibrinogen Concentration	1.92 ± 0.48	1.82 ± 0.24	0.583	0.571
Packed cell volume (PCV)	35.0 ± 1.51	30.5 ± 1.73	4.648	0.001*
White blood cell	13387.5±5846.47	5225.0±206.15	2.724	0.021*
Platelet	520.75 ± 79.16	239.75 ± 39.68	6.583	0.000*

N=8, \*p<0.05 (i.e. Significant).

**Table 2. Comparison of the haemorrhological parameters of those exposed in Group B (20 mins) with the controls (Group D) using paired sample t-test (n=8)**

Haemorrhological parameters	Group B (n=4)	Controls (n=4)	t-test	p-value
Plasma viscosity	3.30 ± 0.33	3.65 ± 0.15	-3.461	0.005*
Whole Blood viscosity	229.08 ± 1.56	229.58 ± 0.79	-0.897	0.389
Plasma Fibrinogen Concentration	1.71 ± 0.24	1.82 ± 0.24	-1.114	0.289
White blood cell	7616.66±1667.24	4825.0±525.19	3.229	0.006*
Platelet	386.41 ± 84.55	239.75 ± 39.68	3.292	0.005*
Packed cell volume (PCV)	32.58 ± 2.42	36.17 ± 1.11	-4.634	0.001*

N=8, \*p<0.05 (i.e. Significant).

care is taken to exclude all air bubbles into the vertical syringe until the plunger pass the 1.0ml mark. The plunger was then completely withdrawn and immediately the lower meniscus falls to the 1.0 mark, a stop watch was started. The time required for 1ml of blood /plasma to drain down the syringe was taken. This was repeated 3 times for each sample and average value was taken. The relative whole blood viscosity /plasma viscosity was then calculated from the formular.

Table 1 shows the comparison of the haemorrhological parameters of those exposed in Group A (10mins) with the control (Group D). There is a significant effect of inhalation of insecticides on plasma viscosity, packed cell volume, total white blood cell counts and platelets when compared with control, while (the whole blood viscosity and plasma fibrinogen concentration were not statistically significant.

Table 2 shows comparison of the haemorrhological parameters of those exposed in Group B (20mins) with the control (Group D). There is significant effect of inhalation of insecticides on plasma viscosity (*P-value of 0.005*), packed cell volume, total white cell counts and platelet counts when compared with the control, But there is no significant effect on whole blood viscosity and plasma fibrinogen/ concentration after 20mins of exposure when compare with the control.

### 3. RESULTS AND DISCUSSION

This research was designed to look at the effect inhaling insecticide on short, midium and long term on haemorheological parameters using rabbits as model. The short term exposure was 10 minutes for seven days. Statistical significant difference decreased was observed in the packed cell volume and significant increase were observed in and Platelet count when compared with the control after inhalation. The relative increase seen in the total white blood cell count though not significant. corroborate the study done by Yousef et al. [13] in which changes in some haematological and biochemical indices of rabbits induced by cypermetin were evaluated, they reported that the increased in WBC may be indicative of activation of defense and immune system of the body but contrary to the results from work done by Kamal et al. [8] in which effect of cypermetrin on clinico haematological parameters in rabbit were evaluated, they reported that. decrease in WBC may be due to viral infection that temporarily disrupt the work of bonemarrow., However, the exposed animal total white blood cell counts did not return from the effect of the exposure after 7days of withdrawal from exposure, this was in accordance with the work done by Adhikari et al. [14] in which effect of cypermetrin and carbonfuran on certain haematological parameters and prediction of recovery in a fresh water teleost, was evaluated and stated that increase in WBCs count

occurred as a pathological response since these WBCs play a great role during infestation by stimulating the haemopoietic tissues and the immune system by producing antibodies and chemical substances working as defense against infection.

The packed cell volume of the treated rabbits was significantly increased when compared to that of the control group, which is contrary to the work done by Guyton and Hall, [15] stated that decrease in PCV of the treated rabbits could have resulted from a likely bonemarrow aplasia reported to have been caused by aerosols of some chemicals such as insecticides.

The midterm exposure (20mins) show significant increased in packed cell volume when compared with the control, which differ from the work done by Sembulingam and Sembulingam, [16] reported that decrease in PCV was time dependent that longer treatment could result to the animals developing aplastic anaemia. but inline with the work done by Iteire who reported that the increase in PCV may be caused by dehydration [17]. The white blood cell counts, show significant increase when compared with the control. Increased in platelets count was also observed which is the same with the work done by Iteire where he reported that the increase in platelets may be due to infection or inflammatory disease [17].

All investigated haemorrhological factors at short term exposure (10 mins), midterm exposure (20 mins) and long term (30 mins) exposure were statistically significant except whole blood viscosity and plasma fibrinogen concentration. which were statistically non significant when compared with control. Increased in plasma viscosity, packed cell volume, white blood cell counts and platelet counts were observed in all the three phases which were in accordance with the work done previously where it was recorded that the increased in plasma viscosity and total white blood cell counts may be caused by infection or inflammatory disease [18]. Plasma fibrinogen concentration and whole blood viscosity were statistically non significant.

#### 4. CONCLUSION

The results from this study have shown that aerosol of these pyrethroid insecticides (prallethrin and cyfluthrin) has effect on haemorrhological parameters. It is thus recommended that one

should avoid exposure to the aerosol of these insecticides during domestic, veterinary, agricultural or industrial use.

#### ETHICAL APPROVAL

Animal Ethic committee approval has been taken to carry out this study.

Animal Ethic committee approval has been collected and preserved by the author(s).

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Solomon KR, Giddings JM, Maund, SJ. Probabilistic risk assessment of cotton pyrethroid: 1. Distributions analysis of laboratory aquatic toxicity data. *Environmental toxicology and chemistry*. 2001; 20(3):652-659.
2. Crago J, Schlenk D. The effect of bifenthrin on the dopaminergic pathway in juvenile rainbow trout (*Oncorhynchus mykiss*). *Aquatic Toxicology*. 2015;162:66-72.
3. Stuart M, Bennett. Classification and Composition of Insecticide. 2003; 1-3.
4. Olawale O. Sub-chronic Effect of Rambo Insect Power Contaminated Diet on Superoxide Dismutase (SOD) Activity in Wistar Albino Rats. *Biokemistri*. 2005; 17:157 – 163.
5. Sangha GK, Khera KS, Singh B. Toxicology effect of cypermethrin on female Albino rats *Toxicol int*. 2011;18:5-8.
6. Shukla Y, Yadav A, Arora A. Carcinogenic and Carcinogenic potential of cypermethrin on mouse skin. *Cancer Lett*. 2000;182: 33-41.
7. Yavasoglu A, Sayim F, Uyanikgil Y, Turgut M, Karabay-Yavasoglu NU. The pyrethroid cypermethrin-induced biochemical and histological alterations in rat liver. *Journal of Health Science*. 2006;52(6):774 – 780.
8. Kamal SM, Khan F, Rizvi M, Siddique S. Effect of cypermethrin on clinical haematological parameters in rabbits. *Pak, Vet. J*. 2007;27:171-175.
9. Sayim FN, Yavasoglu UK, Uyanikgil U, Aktug H, Yavasoglu A, Turgut M. Neurotoxic effects of cypermethrin in Wistar rats: a haematological biochemical and histopathological study. *J. Health. Sci*. 2005;51:300 – 307.

10. Akhigb RE, Ige SF, Adegunlola GJ, Adewunmi MO, Azees MO. Malaria, Haemoglobin genotypes and ABO blood groups, in Ogbomoso, Nigeria. *J. Trop. Med.* 2011;6:73-76.
11. Kingsley CK, Solomon NI, Odudu A. Haematology, Biochemical and Antioxidant changes in wistar rats exposed to dichlorvos based insecticide formulation used in Southeast Nigeria. 2016.
12. Dacie JV, Lewis MS. *Practical Haematology* 7<sup>th</sup> Edn, ELBS Churchill Livingstone, England; 2012.
13. Yousef MI, El-Deerdash FM, Kamel KI, Al-Salhen KS. Changes in some haematological and biochemical indices of rabbits induced by isoflavones and cypermethrin. *Toxicology.* 2003;189: 223 – 234.
14. Adhikari S, Sarkar B, Chatterjee A, Mahapatra CT, Ayyappan S. Effect of cypermethrin and carbofuran on certain hematological parameters and prediction of recovery in a freshwater teleost, *Labeo rohita* (Hamilton). *Ecotoxicol. Environ. Saf.* 2013;58:220-226.
15. Guyton AC, Hall JE. *Guyton and Hall Text book of Medical physiology.* 12<sup>th</sup> ed. Elsevier New Delhi India. 2011; P420.
16. Sembulingam K, Sembulingam P. *Essentials of medical physiology* 4<sup>th</sup> ed. JAYPEE Brothers medical publishers limited New Delhi India. 2006:89.
17. Iteire AK. Haematological evaluation of Albino rats orally administered pyrethroids formulation. *African Journal of Cellular Pathology.* 2016;1:6-10.
18. Dong WY, Kim JK, Junk JS. Increased plasma viscosity in plasma cell dyscrasia and whole blood viscosity in polycythemia vera. *Clinical Haemorrhological and Microcirculation.* 2018;1:59- 67.

© 2022 Eyitayo et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/90286>