

Histopathological Patterns of Nephrotic Syndrome in a Tertiary Care Center in Ile-Ife, Southwestern Nigeria: A 16-Year Retrospective Study

**Adeyemi Abiola Adefidipe ^{a*}, Akinwunmi Komolafe ^a,
Bamikefa Titilope ^b, Kayode Adelusola ^a,
Adegoke Aremu ^a and Fatiu Arogundade ^c**

^a *Department of Morbid Anatomy and Forensic Medicine, Obafemi Awolowo University, Teaching Hospital's Complex Ile Ife, Nigeria.*

^b *Department of Medicine, Osun State University, Teaching Hospital Osogbo, Nigeria.*

^c *Nephrology Unit, Department of Medicine, Obafemi Awolowo University, Teaching Hospital's Complex Ile Ife, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors.

Author AAA performed the research design, did data analysis and wrote first draft of the manuscript. Author KA did the definition of intellectual content and edited the manuscript. Author AK did statistical analysis and edited the manuscript. Author FA did the definition of intellectual content and reviewed the manuscript. Author BT did the sample collection and reviewed the manuscript. Author AA did the Laboratory analysis of sample. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95601>

Original Research Article

Received 20/10/2022

Accepted 29/12/2022

Published 02/01/2023

ABSTRACT

Background: Nephrotic syndrome is one of the major ways in which kidney disease presents clinically the world over. Where biopsies are carried out, the microscopic appearances observed in the affected patients vary according to age, gender, race and geographical location. In most cases,

*Corresponding author: E-mail: Adeyemiabiola87@gmail.com;

the vital information obtained from the histological assessment of these patients is pivotal to instituting effective management of their diseases.

In this study, we set out to describe the spectrum of microscopic findings in renal biopsies obtained from nephrotic syndrome patients in our hospital between January 2002 to December 2017.

Methodology: Archival formalin-fixed-paraffin-embedded renal biopsy tissue blocks were subjected to routine Hematoxylin & Eosin(H&E), Jones Methelamine Silver (JMS), Masson's Trichrome (MT) and Periodic Acid Schiff stains (PAS) as well as immunoperoxidase stains for IgG, IgM, IgA and C3. The slides were viewed using binocular light microscope.

Results: A total of 73 cases were reviewed with a male to female ratio of 1.7:1 and mean age of 20.2 years SD of 11.6). Overall, focal segmental glomerulosclerosis (FSGS) (48.3%) was the most frequent histological pattern observed. FSGS and minimal change disease were the commonest histological patterns seen in children while FSGS and membranous glomerulopathy were the most common patterns in adult population. There were four cases of IgA nephropathy on immunoperoxidase stain. The previous diagnosis was changed in 13 cases, with use of immunostaining which was not carried out ab initio.

Conclusion: This study shows that FSGS is the most common histopathological finding among patients with the nephrotic syndrome in our hospital irrespective of age or gender. In addition, we found that the use of immunoperoxidase in assessment of renal biopsy is important in making a more accurate and clinically relevant diagnosis.

Keywords: Nephrotic syndrome; renal; biopsies; patterns; histopathological.

1. INTRODUCTION

The nephrotic syndrome is the most common manifestation of glomerular diseases. It is a complex clinical condition that is characterized by massive proteinuria, hypoalbuminemia, hypercholesterolemia and generalized oedema [1,2]. Establishing the histopathological basis of this syndrome in the individual case is most desirable for proper management and follow-up subsequently

For a full evaluation of renal disease, clinical data and serological tests, need to be combined with a complete assessment of renal biopsy using light microscopy, immunofluorescence and electron microscopy. While this standard approach is the routine practice in the developed countries of the world, it however still remains a challenge in many of the developing countries (Nigeria inclusive). Up until recently we have depended solely on the use of Hematoxylin & Eosin, JMS, PAS, and MT and these come with all their diagnostic limitations [3]. This study therefore is aimed at evaluating renal biopsy specimens of patients who presented with nephrotic syndrome in our centre over a period of 16 years using light microscopy with the aid of Hematoxylin & Eosin, special histochemical stains and immunoperoxidase stains, since use of immunofluorescence and electron microscopy for paraffin embedded sections are not routinely available in the Nigeria yet. This will enable us to improve on the diagnostic characterization and documentation of the histological basis of nephrotic syndrome in our centre.

2. METHODOLOGY

2.1 Study Design

This was a 16-year retrospective histological descriptive cross-sectional study of renal biopsies obtained from patients who presented with nephrotic syndrome in the Renal Unit of the Obafemi Awolowo University Teaching Hospitals Complex (OAUTHC), Ile-Ife, Nigeria, between January 2002 and December 2017).

2.2 Case Selection

About one hundred and thirty cases of renal biopsies of nephrotic syndrome patients received at the Morbid Anatomy Department O.A.U.T.H.C, Ile-Ife from January 2002 to December 2017 were recruited for the study. The clinical and demographic information of the individual patient was collected from the accession register as well as data from the original consultation request cards of the patients. The slides of the cases were retrieved for review of previous diagnosis. Fresh 3micrometer sections were cut from archival tissue blocks and stained with H and E, Periodic Acid Schiff, Jones silver methenamine, and the Masson's trichrome stains.

Immunoperoxidase stains using IgG, IgM, IgA, and C3 antibodies after thermal antigen retrieval were also applied. Appropriate positive and negative tissue controls were used for each of the stains applied.

All the slides were reviewed using a standard compound binocular light microscope and histological findings were noted.

2.3 Exclusion Criteria

Fifty-seven cases whose tissue blocks have been used up as well as those with inadequate tissue samples and incomplete clinical data were excluded from the study.

2.4 Data Analysis

The results were presented using simple descriptive statistical analysis, frequency tables and charts. Statistical analysis was carried out using Statistical Package for Social Sciences software (SPSS Inc. Chicago, IL, USA) version 20.

3. RESULTS

There were 73 patients with the nephrotic syndrome between 2002 and 2017 who had biopsy carried out on them. Of these 73 cases, 46 were males and 27 were females (male to female ratio 1.7 to 1). The mean age was 20.8 ± 12.4 years, the ages ranging from 5 to 56 years.

Most of the patients were children below the age of 16 years and FSGS was the most predominant pattern.

Excepting IgA Nephropathy and post-infectious GN, which occurred more commonly among females, all the histological abnormalities were commoner among males. Fig. 1 depicts the range of histopathological abnormalities that we found.

3.1 Histopathological Abnormalities

3.1.1 Morphological patterns

FSGS was the most common histological abnormality across all age groups in this study

accounting for 42.5% of the cases (Thirty-one cases). Minimal change disease in children and membranous glomerulopathy in adults were the next most common. Membranous glomerulopathy was responsible for sixteen cases (21.9%). There were thirteen cases of minimal change disease (17.8%), six cases of membranoproliferative glomerulonephritis (MPGN) (8.2%), four cases of IgA (5.5%) Nephropathy, two cases of Diabetic nephropathy (2.7%) and one case of Postinfectious glomerulonephritis (1.4%). Table (1).

3.1.2 Immunoperoxidase staining patterns

The immunoperoxidase staining was positive in sixty cases and negative in thirteen cases. These immunoperoxidase-negative were the cases of minimal change disease.


3.1.3 Review of initial diagnosis

The initial diagnosis was changed in eleven cases after special stains and immunoperoxidase were applied. Initially, eight of these eleven cases had been diagnosed as minimal change disease, one as FSGS, one as Membranoproliferative glomerulonephritis and one as Mesangioproliferative glomerulonephritis (MSPGN). Six of the initial minimal change disease group were subsequently re-designated as membranous glomerulopathy and two as IgA nephropathy. The cases that had been diagnosed as MSPGN and MPGN were re-designated as IgA nephropathy, because of the striking IgA staining on immunoperoxidase. The initial case of FSGS had to be changed to Membranous glomerulonephritis Table (1).

Table 1. Showing the various frequencies of morphological patterns found in renal biopsies of the Nephrotic syndrome cases seen at the O.A.U.T.H.C using routine and immunoperoxidase stains

Histological patterns	Frequency (n/%) routine stains	Frequency(n/%) immuno stains
FSGS	32(43.8%)	31(42.5%)
MGN	9(12.3%)	16(21.9%)
MCD	21(28.8%)	13(17.8%)
MPGN	7(9.6%)	6(8.2%)
IGAN	0(0%)	4(5.5%)
DN	2(2.7%)	2(2.7%)
MSGN	1(1.4%)	0(0%)
PIGN	1(1.4%)	1(1.4%)
Total	73(100%)	73(100%)

FSGS, Focal segmental glomerulosclerosis; MGN, Membranous glomerulopathy; MCD, Minimal change disease; IGAN, IgA Nephropathy; DN, Diabetic Nephropathy; MSGN, Mesangioproliferative glomerulonephritis; MPGN, Membranoproliferative glomerulonephritis, PIGN, Post-infectious glomerulonephritis



DN, Diabetic Nephropathy; MGN, Membranous glomerulopathy; FSGS, Focal segmental glomerulosclerosis; MCD, Minimal change disease; IGAN, IgA Nephropathy; MPGN, Membranoproliferative glomerulonephritis; PIGN, Postinfectious glomerulonephritis

Fig. 1. Gender distribution of the various morphological patterns in nephritic syndrome patients in O.A.U.T.H.C.




Fig. 2. Subtypes of Thirty-one Cases of Focal segmental glomerulosclerosis in nephrotic syndrome patients at the O.A.U.T.H.C




Fig. 3. Photomicrographs of glomeruli showing thickened glomerular basement Membrane in a 23years old. (PAS stain x 400)




Fig. 4. Granular IgG staining along the glomerular basement membrane in membranous glomerulopathy. (IgG Immunoperoxidase stain x 100)




Fig. 5. Photomicrograph shows area of segmental (Black arrow) sclerosis in a glomerulus in a case of FSGS in a 12-year old boy. (Masson's Trichrome x 100)

4. DISCUSSION

Nephrotic syndrome is the most common indication for renal biopsy worldwide [4]. It is a major indication for renal biopsy in our center within the study period and accounted for 58.5% of the cases of renal diseases where biopsies were carried out. This is similar to studies done by Obiagu et al [5], Patricia et al. [6] and Okpechi et al. [7] in which nephrotic syndrome was the most common indication for renal biopsies. Our findings is however at variance with what Ibrahim et al in Egypt reported in which isolated sub-nephrotic proteinuria constituted the major indication for renal biopsy [8]. This difference may be attributable to the variation in each hospital's renal biopsy protocol. We observed that nephrotic syndrome was commoner in males than in females across all the age groups and this is in agreement with what has been previously reported [9-11]. Umuiezudike et al. [12] in a prospective study of 30 cases in a Lagos clinic however recorded equal prevalence in both genders and this may be because of their smaller sample size.

Most of the patients in our study were below 16 years in whom focal segmental glomerulosclerosis and minimal change disease constituted the major histological patterns. Focal segmental glomerulosclerosis was seen in 40.5% of 37 children in this study. Several local studies in Nigeria recorded similar findings, [13-15]. Doe

et al. [16] in Ghana also reported Focal segmental glomerulosclerosis and minimal change disease as the main histological patterns in their cohort of children with nephrotic syndrome, especially in those associated with steroid resistance. In contrast to our study however, they reported no case of membranous or membranoproliferative glomerulonephritis, and this may be because of their small sample size. They only analyzed thirteen cases. In contrast to our work however, previous work on nephrotic syndrome in adults by Oviasu et al in South-south Nigeria documented minimal change and proliferative glomerulonephritis each accounting for 33.3% as the leading causes of nephrotic syndrome [17]. Obiorah et al. [18] in Port Harcourt also, in a study of nineteen childhood and adult cases of nephrotic syndrome, reported membranoproliferative glomerulonephritis as the most common basis of nephrotic syndrome. It is to be noted however, that many of these Nigerian studies examined smaller cohort than we did and that they only employed routine and special stains. That may well explain the differences in our findings.

Minimal change disease is the second most commonly associated morphological finding in children and third commonest histological pattern in adults with nephrotic syndrome in our series. Even though minimal change disease is the most common cause of nephrotic syndrome in children, most of them do not get to be biopsied as they would have responded to steroid therapy

and hence the lower incidence reported compared to FSGS in this our study. This is in agreement with obtains in other studies both locally and globally. Asinobi et al. [13] in Ibadan reported minimal change disease as the third commonest histological pattern associated with nephrotic syndrome in their patients.

Chijoke et al. [9] in a clinicopathological study of nephrotic syndrome in adults in North–Central Nigeria reported minimal change disease as the least commonly associated histological pattern (5.3%).

This is however contrary to the finding by Rico et al. [19]. who in a Colombian paediatric population with nephrotic syndrome, reported minimal change disease as the most common histological pattern observed in children with nephrotic syndrome followed by FSGS and membranous glomerulopathy. The difference from our finding could be due to regional variation in renal biopsy indications. Although Bornilla-Fellix et al. [20] had noted a change in the trend of morphological patterns in the renal biopsy of children with nephrotic syndrome in the American population, they still reported minimal change disease as the most common cause of nephrotic syndrome, the essential difference being an increase in the reported incidence of FSGS particularly in the African-American paediatric population.

IgA nephropathy (IgAN) is recognized to typically present with haematuria, but could be a basis of nephrotic syndrome in the rare cases. We found four cases of IgA nephropathy on immunoperoxidase stain. These cases are characterized by endocapillary proliferation and segmental glomerulosclerosis in some of the glomeruli. Leal et al. [21] in a study of 17 cases of IgAN presenting with nephrotic syndrome suggested that these are cases of IgAN with coexistent minimal change disease.

5. CONCLUSION

Our study showed that FSGS was the most common histopathological pattern seen amongst nephrotic syndrome patients in our hospital cutting across age and gender and this was followed by membranous glomerulopathy in adults and minimal change disease in children. It also underscores the importance of the use of special stains and immunoperoxidase stains in the assessment of renal biopsies. This study, while highlighting the need for improving the infrastructural repertoire for the tissue diagnosis

of the nephrotic syndrome and other renal diseases in resource-limited settings such as ours, provides important preliminary data to guide local nephrologists and provide the basis for further studies on the aetiopathogenesis of the disease in Tropical Africa.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Gesualdo L, DiPalma AM, Morrone LF, Strippoli GF, Schena FP. Italian Immunopathology Group, Italian Society of Nephrology. The Italian experience of the national registry of renal biopsies. *Kidney Int.* 2004;66:890-4.
2. Charles EA. Glomerular diseases. In: Kumar.V, Fausta. N, Aster.J AK, editor. *Robbins and Cotran Pathological Basis of Disease.* 8th ed. Philadelphia: Saunders Elsevier. 2010;978–9.
3. Onwubuya IM, Adelusola KA, Sabageh D, Ezike KN, Olafoe OO. Biopsy- proven renal disease in Ile Ife, Nigeria. A histopathological review. *Indian Soc Nephrol.* 2016;26(1):17–21.
4. Hull RP, Goldsmith DJ. Nephrotic syndrome in adults. *BMJ [Internet].* 2008; 336(7654):1185–9.
5. Obiagwu PN, Atanda AT, Aliyu A. Nephrotic syndrome among children in Kano: A clinicopathological study. *Niger J Med Pract.* 2014;17(3):370–4.
6. Patricia M et al. Paulista registry of glomerulonephritis: 5-year data report. *Nephrol Dial Transplant.* 2006;21(11):3098–105.
7. Okpechi IG, Ameh OI, Bello AK, Ronco P, Swanepoel CR, Kengne AP. Epidemiology of histologically proven Glomerulonephritis in Africa: A systematic review and meta-analysis. *Plos One.* 2016;11(3):1–15.
8. Ibrahim S, Fayed A, Fadda S. BD. A five-year analysis of the incidence of glomerulonephritis at Cairo University Hospital-Egypt. *Saudi J Kidney Dis Transpl.* 2012;23:866–70.
9. Chijioke A, Adeniyi AB. Clinicopathological study of adult nephrotic syndrome in

- Ilorin, Nigeria. Niger Med Pract. 2003;43(2):28–32.
10. Kazi JI, Mubarak M, Ahmed E, Akhter F, Naqvi SAA, Rizvi SAH. Spectrum of glomerulonephritides in adults with nephrotic syndrome in Pakistan. Clin Exp Nephrol. 2009;13(1):38–43.
 11. Caneta PA, Radhakrishnan J. The evidence-based approach to Adult onset Idiopathic Nephrotic syndrome. Vol3. Front peditric; 2015.
 12. Umezudike TI et al. Renal histology pattern in prospective study of nephrology clinics in Lagos Nigeria. Clin Nephrol. 2016;23:354–66.
 13. Asinobi AO, Ademola AD, Okolo CA, Yaria JO. Trends in the histopathology of childhood nephrotic syndrome in Ibadan Nigeria: preponderance of idiopathic focal segmental glomerulosclerosis. BMC Nephrol [Internet]. 2015;1–9. Available:<http://dx.doi.org/10.1186/s12882-015-0208-0>
 14. Olowu WA, Adelusola KA, Adefehinti O. Childhood Idiopathic Steroid Resistant Nephrotic Syndrome. Saudi J Kidney Dis Transplant. 2010;21(5):979–90
 15. Obiagwu PN, Atanda AT, Aliyu A. Nephrotic syndrome among children in Kano: A clinicopathological study. Niger J Med Pract. 2014;17(3):370–4.
 16. Doe JY, Funk M, Mengel M, Doehring E, Ehrich E, Jochen HH. Nephrotic syndrome in African children: Lack of evidence for ‘tropical nephrotic syndrome’? Nephrol Dial Transpl. 2006;21(December 2005): 672–6.
 17. Oviasu E, Ojogwu L. Another look at the nephrotic syndrome in adult Nigerians: Pathological and immunological findings. West Afr J Med. 1992;11:18–24.
 18. Obiorah CC, Okoro PE, Clinicopathological report of children and young adults with nephrotic syndrome undergoing renal biopsy at workshops in Port Harcourt, Nigeria. African J Nephrology. 2019; 21(1):3-7.
 19. Rico MP et al. Characterization and Etiopathogenic Approach of Pediatric Renal Biopsy Patients in a Colombian Medical Center 200-7-2017. International Journal of Nephrology. 2018; 205-206.
 20. Bonilla-Felix M et al. Changing patterns in the histopathology of idiopathic nephrotic syndrome in children. Kidney Int [Internet]. 1999;55(5):1885–90.
 21. Leal CH, Andrew SB, Micheal BS, Jai R, D’Agati VD, Glen SM. IgA Nephropathy with minimal change disease. CJASN. 2014;9(6):1033-1039.

© 2023 Adefidipe et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/95601>