



Predisposing Factors Determining the Rearing Performance of Muga Silkworm (*Antheraea assamensis* Helfer): A Review

Abhishek Singh ^{a++*}, Vikram Kumar ^{b#}, Lopamudra Guha ^{c#},
H. Hridya ^{c#}, K. Indirakumar ^{d#} and M. Majumdar ^{e#}

^a MESSO, Central Silk Board, Nongpoh, Meghalaya, India.

^b MESSO, Central Silk Board, Rompara, Meghalaya, India.

^c MESSO, Central Silk Board, Guwahati, Assam, India.

^d MESSO, Central Silk Board, Tura, Meghalaya, India.

^e MESSO, Central Silk Board, Mendipathar, Meghalaya, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i242697

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95886>

Review Article

Received: 20/10/2022

Accepted: 30/12/2022

Published: 30/12/2022

ABSTRACT

Muga silkworm (*Antheraea assamensis* Helfer) is endemic to North-Eastern India, and colour designates it as golden silk. From time immemorial, many ethnic and tribal groups have produced muga silk. Muga silk has a special status due to its natural golden-yellow colour. Semi-domesticated muga silkworm (*Antheraea assamensis* Helfer) is multivoltine in nature and commercial rearing is conducted during the spring (Jethua) and autumn (Kotia) seasons. The muga

⁺⁺ Scientist – B;

[#] Scientist – C;

*Corresponding author: E-mail: abhisheksinghcsa@gmail.com;

silkworm is a mono race with little genetic variation among populations, rearing is completely in an open environment and is subject to various diseases, pests and predators. Due to unpredictable weather conditions, natural calamities along with disease outbreak and natural enemies, the muga silk production has been affected dramatically. To improve the productivity of this silkworm it is important to have a better knowledge of the factors that affect the most to this silkworm. So, the present study is mainly discussed about factors responsible for the rearing of muga silkworm.

Keywords: Muga silkworm; seed; factor affecting; rearing seasons.

1. INTRODUCTION

“Muga silkworm *Antheraea assamensis* Helfer is a Lepidopteran insect of *Saturniidae* family and, is geographically isolated only to NE region of India. The geographical endemicity of this silkworm is indicative of its special requirements for geo-climatic conditions that prevail in this region i.e. high humid temperate climate and forest vegetation of primary and secondary host plants” [1]. “Thus, this species is phylogenetically less adaptive reaching its ecological isolation that is indicative of being on the verge of extinction. Although Muga silkworm has been reared for production of silk but outdoor practice on host plant under natural conditions poses a big challenge. Being exposed to the natural environment muga culture faced lots of problems right from brushing of worms to spinning of cocoons. Outdoor silkworm larvae are invariably exposed to nature’s vagaries such as seasonal climate change, rainfall, strong wind, and soaring temperature, besides pests, predators and pathogens inflicting heavy loss particularly in early three instars reported that in an average in all seasons more than 50% larval loss was due to abiotic factors and 80% of the total loss of muga silkworm occurred in second/third instars only” [2]. “Rearing being outdoor, the success of crops is always bestowed on the mercy of environmental conditions as muga rearers practice traditional cultural methods. Out of the 5-6 broods of rearing in a year, the commercial crops fall in favorable period while all seed crops rearing coincide with the period of extremities of temperature, humidity, rainfall etc. The Aghenua crop which is pre-seed crop for Chatua seed crop falls during December-January. Low temperature prevailing during this season emanated to the lengthening of larval duration and outbreak of fungal diseases mainly muscardine leading to heavy larval mortality. The resultant in turn is low rate of seed multiplication affecting the availability of seed to Chatua Seed and subsequent multiplication for Jethua commercial crops. Similarly, seed supply for Kotia commercial crop has also become inadequate as

the seed multiplication rate of its preceding pre-seed and seed crops i.e. Aherua and Bhadia are very low. These two crops fall during unfavorable climatic conditions is characterized by high temperature, high humidity and high rainfall” [3]. According to [4] “relative contribution of several factors responsible for a successful crop harvest were estimated as: Host plant (38.2%), climate (37.0%), rearing technique (9.3%), silkworm race (4.2%), silkworm egg (3.1%) and other factors (8.2%). Since, the quality of host plants played a major role in successful cocoon production, several workers tried indoor rearing of muga silkworm to avoid early instar loss, unfavorable seasons of extremities of temperature, humidity and rainfall. Several works had been carried out on different factors affecting muga silkworm rearing which are discussed in the present study”.

2. REARING SEASON

Muga is a multivoltine silkworm which completes its life cycle 5 to 6 times a year. Duration of the lifecycle of the muga silkworm may vary from 50 days in summer to 150 days (maximum) in winter [5]. Each life cycle takes about two months on an average making it possible to organize 5 or 6 crops in a year. In Assam, muga farmers have named the crops as Kotia (October to November) in the autumn season, Jarua (November to January) in the winter season, Chotua (February to April) in the early spring season, Jethua (April to May) in the spring season, Aherua (June – July) in early summer and Bhodia (August – September) in late summer season. The Katia is the main commercial crop and the Jethua is the second commercial crop whereas Jarua and the Aherua are the two pre-seed crops and the Bhodia and the Chotua are the seed crops of muga silkworm farming practice [6]. Multivoltine nature is one of the major problem for which the maintenance of different seed broods of muga silkworm is difficult, time consuming, laborious and hazardous making the unavailability of good quality seed cocoons in specific seasons. Dutta et al. [7] reported that “muga silkworm has the

potential to lay a good number of eggs (250-280) but realized fecundity (120-150) was comparatively poor even during the favorable seasons of *Jethua* and *Katia* compared to eri (440-470) and mulberry (450-550)".

3. INFLUENCE OF CLIMATE AND SEASON

"The performance of muga silkworm rearing has been impacted in recent years by climate change and pollution i.e., prolonged summer and short winter. Rescheduling the crop cycle in coordination to changing climate is a prerequisite. The ideal conditions for rearing Muga silkworms are 20–31 °C temperatures and 65–95% relative humidity" [8]. "Change of climatic factors, especially temperature and relative humidity affected almost every aspect of the life cycle of silkworms including their development and survival" [6]. "Even, the change of season can influence the consumption and utilization of food in muga worms" [9]. "Muga silkworm is semi-domesticated and rearing is conducted in outdoor condition, they may not be able to adjust to the new changing environment" [10] and thus "the differential seasonal conditions greatly influence the growth and development of muga silkworm" [11]. "Because of unpredicted climatic conditions as well as other biotic factors, the seed crops are low productive (14-40% crop loss) and sometimes highly uncertain leading to uneconomic crops" [12]. However, these seed crops are mandatory to maintain the linkage for production of commercial seed for *Kotia* and *Jethua* crops. There is no clear cut demarcation of seed and commercial zone for muga silk worm rearing. As a result, the main constraint remains the availability of high-quality muga silkworm seed, which is critical to the muga industry's productivity, sustainability, and profitability. "During the last few decades, atmospheric pollution and the resultant variability in temperature and relative humidity due to global warming along with abnormal rainfall patterns, drought, and floods caused the continuous failure of the crop or low crop yield despite all efforts and utilization of resources. Besides an abnormal increase in temperature, the other reasons enlisted for the heavy loss of muga silkworm were air pollution caused by rampant use of pesticides in neighboring tea gardens, pollution from the brick kilns and burning of natural gases emitting from oil wells and a seismic survey by ONGC for oil exploration" [13]. "It was also reported that large-scale of muga culture was destroyed in Upper Assam by pesticides sprayed in nearby tea gardens" [14].

4. EFFECT OF FOOD PLANT

"The muga silkworm, *Antheraea assamensis* Helfer feeds on a wide range of host plants among which Som, *Persea bombycina* and Soalu, *Litsea monopetala* are the primary food plants of this silkworm" [15]. "*Persea bombycina* plantation is evergreen in nature, easy to maintain and propagate, and overall, its rearing performance is better than that of other food plants. Although *L. monopetala* is a primary food plant, muga silkworm farmers rarely raise nursery or plantation of this plant because it defoliation and has poor leaf quality during winter" [16]. "The silkworms were mostly attracted to semi-mature leaves of *P. bombycina* rather than other primary and secondary food plants. Som leaves are said to be suitable in rearing muga silk for higher yield. Chawki rearing can be done in Soalu plants and then switched to Som plants for late age rearing if there is a paucity of Som leaves during commercial rearing" [17]. "However, the pupal weight and oviposition rate was higher when the silkworms fed on *L. monopetala* leaves compared to *P. bombycina*, *L. salicifolia*, and *Cinnamomum glaucescens*" [18]. "Larval duration did not vary when Muga silkworms fed on *P. bombycina* and *L. monopetala* leaves in different seasons" [19]. "There were no significant differences in shell weight, shell ratio, filament denier and length, or silk recovery, and there was no significant variation in fecundity, hatching percentage, effective rate of rearing (ERR), and cocoon:dfi (disease-free layings) ratio between the two food plants", as reported by Kakati [19]. "The primary food plant Soalu (*Litsea monopetala*) is mostly utilized for production of seed cocoon purpose. The plant is distributed in plains as well as hills, and the propagation occurs by falling seeds. Proteins and amino acids are important for silkworm larvae as they are utilized for synthesis of silk protein" [20].

5. EFFECT OF NUTRITIONAL STATUS OF THE LEAF

"Feeding is one of the major factors in silkworm rearing. Healthy host plant leaves that are rich in essential nutrients are crucial for optimal feeding. Muga silkworms obtain their all essential nutrients from the leaves which they take as food. The nutritional status of the leaves can be improved by enriching them with vitamins and other nutrients" [21]. "The feeding of nutritionally enriched leaves showed better growth and development of silkworm larvae, as well as

directly influencing the quality and quantity of silk production” [22]. “The host leaves mainly constitute proteins, carbohydrates, vitamins, sterols, stimulants, and minerals. Such nutritional requirements in food consumption have a direct impact on all genetic traits such as larval and cocoon weight, the quantity of silk production, pupation and reproductive traits” [23]. Yadav & Goswami [24] reported “16.2% crude protein in the tender leaves of *P. bombycina*, and 15.5% in medium and mature leaves, whereas leaves of *L. monopetala* contained 20.7% (tender), 18.2% (medium), and 15.5% (mature) crude protein”.

6. INCIDENCE OF DISEASE

“Disease and pest infestation pose serious threats to seed crop rearing. Muga silkworm is subjected to viral, bacterial, fungal, and protozoan diseases that resulted in heavy crop losses up to 40% for individual diseases” [25]. “It is estimated that 20–30% of the loss is due to pebrine (microsporidia) disease, which sometimes kills an entire silkworm culture” [26]. “Flacherie is another common disease of muga silkworm. It is caused by a virus followed by a secondary infection with bacteria. The incidence of this disease is highest in summer and this is thought to be due to sudden fluctuations in temperature coupled with poor food quality” [27]. “Grasserie disease is caused by a virus and can cause heavy crop losses. It infects larvae mainly in summer and rarely in winter. Muscardine is a less prevalent disease caused by a fungus (*Fusarium* spp.). It infects larvae mainly in winter” [8]. Patnaik, [28] also stated that fungal infection is rare in muga crops.

7. INCIDENCE OF PEST

Muga Silkworm is attacked by several parasitoids and predators (ants, wasps, birds etc) [29]. “Among these *Exorista sorbillans* is one of the serious larval end parasitoids of muga silkworm (*Antheraea assamensis*). It causes extensive damage to the sericulture industry. During Jarua (December-January) and Chotua (February-March) crop seasons and reported 20-90% loss in Jarua and Chotua season (December-March) and 50-70% cocoon rejection during February-March” [30]. “The mature maggots come out of the larvae/pupae and undergo pupation in the rearing field or grainage hall. The uzi infested muga silkworm dies during the larval or pupal stage. This parasitoid was also reported on 95 species of insects belonging to 20 families of Lepidoptera and one family of Hymenoptera

worldwide in the absence of silkworm” [31]. Thangavelu and Sahu, [32] reported that “the maggots of the Uzi fly exhibited considerable variation in their body size. The maggots developed within *B. mori* larvae were generally smaller than those developed within *A. assamensis* larvae were larger. They also suggested that the much larger muga silkworm might have provided a better niche for the uzi fly maggot than the smaller silkworm”. “The recent survey results indicate that the maximum infestation of uzi fly was recorded in 5th instar larvae (43.0%) and harvesting of cocoons (35.0%) during Chotua crop (March-April) followed by Jarua crop,(19.0%) infestation at larval stage & 27.50% at harvesting stage of cocoons reported in upper Assam” [33]. “The muga cocoons are also infested by yellow fly. The fly completely fed on the head region of the pupae and emerged from the same by making a hole that is about 1cm in diameter” [30]. Instances of a pupal parasitoid Ichneumon wasp (*Xanthopimpla pedator*) on muga silkworm have been reported for the first time from various muga growing locales in West Garo Hills, Meghalaya, during various muga summer crops [34].

8. QUALITY OF MUGA SILKWORM SEED

The foundation of the sericulture industry is the seed. Silkworm seed of high quality means that the laying is disease-free, that there are more viable eggs, that the hatching is uniform, and that the crop is stable. Healthy seed production is indeed the primary requisite for the conducive growth of the entire silk industry [35]. A timely supply of superior quality silkworm seed can alone sustain sericulture as a commercial crop in competition with other cash crops. Not only is the augmentation of silkworm food plants under suitable climatic conditions important for the success of the sericulture industry, but so is the timely supply of superior quality silkworm seed to farmers. The quality of silkworm seed may be defined as to the one where the laying are entirely free from diseases, has maximum nos., of viable eggs, gives uniform hatching and assures a stable crop [6].

9. PRESERVATION OF SEED COCOON

Species of *Antheraea* undergo diapause in the pupal stage, except *A. yamamai*, which undergoes diapause in both the egg and pupal stages [36]. For rearing commercial crops (either *Kotia* or *Jethua*) four seed broods are to be

maintained in a year. Because seed broods are reared during unfavorable seasons, inducing pupal diapause to delay moth emergence by storing seed cocoons at low temperatures may aid in avoiding the unfavorable seasons. Domesticated muga silkworm did not either over winter or diapause at high altitude [37]. Haniffa and Thatheyus, [38] reported that increase in the duration of preservation led to the aging of male moths which was also attributed to less pairing efficiency and less productivity. Khatri, [39] also reported that the rearing and grainage performance of muga silkworm during four crops, namely Aherua, Jarua, Bhodia, and Kotia, were found to be better in Doon valleys than in Assam from 1998 to 2000. Preservation of seed cocoons at 5 °C beyond 30 days led to detrimental effects on economic characteristics in muga silkworms [40]. Tolerance of this temperature beyond 30 days was fully detrimental, but its tolerance beyond 120 days showed almost complete pupal mortality [41]. Sahu et al. [42] “preserved cocoon from *Jethua* crop (April–May) at 10± 1.5 °C and RH of 80-85% for 25, 30, and 35 days and observed that reproductive physiology was affected when cocoons preserved beyond 30 days. They also observed that the hatching percentage in the treated lot was 16-46%, compared to 5% in the control”. “Whereas, long-term seed cocoon preservation for up to 42 days was achieved by Rajkhowa et al. [43] following the double-step preservation method without affecting the grainage parameters”. “Several other studies also revealed that refrigeration of muga seed cocoons at 5-12°C for 10-120 days gave satisfactory results in terms of moth emergence, pairing, fecundity, and hatchability” [44,45].

10. CONCLUSION AND FUTURE PERSPECTIVES

The golden silk of North East India is the precious silk for India. Many difficulties are being faced by the farmers and its related peoples. We have discussed briefly almost all the threats related to silkworm rearing. Muga silk is the heritage of Assam as well as for Assamese people. There is a need to study scientifically all the effecting causes of muga silkworms and the silk production. Development of proper indoor rearing technique to grow the larvae and turn them into completely domesticated needs vast and tedious scientific study. Muga silkworms are very much vulnerable to climatic and any other change in the environment. Keen investigations and observations should be made to develop some

new technologies which will help the muga silkworms to continue to exist but not to be in near extinction state. At last we should welcome a silk revolution in Assam to save this priceless heritage which is one of our social identities and the symbol of our rich history.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Singh A, Kumar V, Indirakumar K, Guha L, Majumdar M. Distribution, Ontogeny and Ethology of Muga Silkworm *Antheraea assamensis*, Advances in Agricultural Entomology. 2022;(20):83-95.
2. Sengupta AK, Siddique AA, Barah A, Negi BK. Improved technologies for Muga silkworm rearing, a development perspective. Indian Silk. 1993;31: 21-24.
3. Neog K, Dutta P, Changmai A, Goswami D, Choudhury B. Comparative Study on the Rearing Performance of Muga Silkworm under Indoor and Outdoor Rearing Conditions. Int J Agricultural Sci Food Technology. 2015;1(1):20-24.
4. Choudhury CC, Dubey OP, Choudhury A, Sengupta K. Effect of refrigeration and preservation of muga seed cocoons *Antheraea assamensis* Ww. Sericologia.1987;27(3):547-552.
5. Khanikor DP, Dutta SK. Rearing schedule of Katia crop and Jethua crop cycle of muga silkworm (*Antheraea assama* Ww). Muga silkworm biochemistry, biotechnology, molecular biology (Ed. Unni, B.G) RRL, Jorhat. 2006;51-59.
6. Saikia M, Ghosh K, Richard, Peigler S. Factors affecting on quality Muga silkworm (*Antheraea assamensis* Helfer) seed crop production: A review, Journal of Entomology and Zoology Studies. 2016;4(6):806-810.
7. Dutta P, Dutta P, Rai AK, Neog K. Improvement of rearing performance in muga silkworm *Antheraea assamensis* with hormone treatment. Mun. Ent. Zool. 2013;8(2):767-771.
8. Tikader A, Vijayan K, Saratchandra B. Muga silkworm, *Antheraea assamensis* (Lepidoptera: Saturniidae) – An overview of distribution, biology and breeding. Eur. J Entomol. 2013; 110(2):293-300.

9. Das P, Unni BG, Bhattacharya PR, Deka PC. Seasonal changes in food consumption and utilization pattern of semi-domesticated muga silkworm, *Antheraea assama* Westwood (Lepidoptera: Saturniidae). J Entomol. Res. 2002;26(4):277-284.
10. Singh RN, Maheshwari M. Conservation and utilization of sericigenous insects in North East Region of India. Sericologia. 2003;43(1):1-15.
11. Chiang HC. Insects and their environment. Fundamentals of applied Entomology. Macmillan Publishing Company, New York. 1985;128-161.
12. Chakravorty R, Das R, Neog K, Das K, Sahu M. A diagnostic manual for diseases and pest of Muga silkworm and their host plants. CMER&TI, CSB, Lahdoigarh, Jorhat, Assam. 2007;1-47.
13. Bhattacharyya PR. Studies on the effect of oil field population upon nutritional quality of "Som" (*Machilus bombycina*) and the biology and certain biochemical aspects of muga silkworm "*Antheraea assama* Ww". Unpublished Ph. D Thesis, Gauhati University; 1994.
14. Bhattacharyya S. Tea planters' friend turns muga farmers' foe – Pesticides sprayed in nearby gardens are destroying large tracts of silkworms in Upper Assam. The Telegraph; 2012.
15. Kumar V, Singh A, Indirakumar K, Majumdar M, Guha L. Effect of different host plants on rearing and grainage activity on Muga silkworm (*Antheraea assamensis*). International Journal of Agriculture Sciences. 2022;14(8):11559-11562.
16. Yadav GS, Goswami BC. Golden muga silk in Mizoram, Indian Silk. 1987;26: 14-15.
17. Borpuzari SP, Das MR, Rahman A, Comparative study on the Rearing performance of muga silkworm (*Antheraea assamensis* Helfer) on two primary host plants Som (*Persea bombycina* Kost) and Soalu (*Litsea monopetala* Roxb) in different rearing seasons. Eco. Env. & Cons. 2022;28(1):568-573.
18. Barah, A. and Sengupta, A. K. Correlation and regression studies between pupal weight and fecundity of muga silkworm *Antheraea assama* Ww. (Lepidoptera: saturniidae) on four different food plants. Acta Physiologica Hungarica. 1991;78: 261–264.
19. Kakati BT. Altitudinal effect and host plant preference on growth and production of *Antheraea assama* Westwood, PhD Dissertation, Nagaland University, Lumami, India; 2012.
20. Bose PC, Bindroo BB. Response of mulberry (*Morus alba* L.) to varying combinations of nitrogen, phosphorus and potassium under rainfed condition of subtropical region growth, productivity and economics. Indian Journal of Sericulture. 2001;40:116-118.
21. Kanafi RR, Ebadi R, Mirhosseini SZ, Seidavi SR, Zolfaghari M, Etebari K. A review on nutritive effect of mulberry leaves enrichment with vitamins on economic traits and biological parameters of silkworm *Bombyx mori* L. ISJ. 2007;4:86-9
22. Neto JB, Stefan V, Mendonca BB, Bloise W, Ana Valeria B, Castro MD. The essential role of zinc in growth. Nutrition Research. 1995;15:335-358.
23. Khurram S. Combined effect of mineral nutrients on silkworm growth and silk yield of *Bombyx mori* L. M.Sc. Thesis, Department Agriculture Entomology University Agriculture Faisalabad, Pakistan; 1998.
24. Yadav GS and Goswami BC. Effect of ringing and auxins on the vegetative propagation of foliated cuttings of som (*Machilus bombycina* King) (Laurales: Lauraceae), *Sericologia*. 1992;32:437-446.
25. Veeranna G. Integrated silkworm disease management: China vs. India. Indian Silk. 1999;7:27-29.
26. Chakrabarti S, Manna B. Studies on cross infection of microsporidian spores of mulberry, eri and tasar silkworm to muga silkworm (*Antheraea assamensis* Helfer) and a new approach for management to pebrine disease in muga silkworm in Assam and west Bengal, India. Zool. Res. Human Welfare. 2008;47:447-460.
27. Aruga H, Fukuda H, Yoshitake N. Observations on a polyhedrosis virus within the nucleus of the silk gland cells of the silkworm, *Bombyx mori* L. J Sericult. Sci. Jpn. 1963;32:213-218.
28. Patnaik RK. Sericulture Manual. Biotech. Books Publ. Tri Nagar Delhi-110035, India. 2008; 139.
29. Singh PK, Das PK, Negi BK. Studies on Seasonal Uzi Fly. Annual Report, Regional Muga Research Station, Boko, Assam. 1993;7.

30. Baruah JP and Kalita C. Integrated pest management of uzi fly (*Exorista sorbillans*) in Muga silkworm *Antheraea assamensis* Helfer (Lepidoptera: saturniidae): A review. Journal of Entomology and Zoology Studies 2020;8(4):341-343.
31. Narayanaswamy KC, Devaiah MC. Silkworm Uzi fly. Zen publishers, No. 1489, 40th Cross, 4th 'T' Block, Jayanagar, Bangalore, Karnataka. 1998;232.
32. Thangavelu K, Sahu AK. Some studies on the bionomics of *Exorista sorbillans* Nied from North Eastern India. Sericologia. 1986;26:77-82.
33. Eswara Reddy SG. Integrated management of Uzi fly, *Exorista bombycis* (Louis) (Diptera: Tachinidae) in Muga Silkworm, *Antheraea assamensis* Helfer (Lepidoptera: Saturniidae) under outdoor rearing conditions of Assam (India). Munis Entomology & Zoology. 2011;6(2):1012-1013.
34. Majumdar M, Guha L, Kumar V, Das B, Borpuzari P. Ichneumon wasp *Xanthopimpla pedator*-a pupal parasitoid of Muga silkworm *Antheraea assamensis*. Int. J. Adv. Res. Biol. Sci. 2021;8(7):152-156.
35. Majumdar M, Kumar V, Borpuzari P, Singh A, Cost effective management of summer grainage (May-August) of Muga silkworm (*Antheraea assamensis* Helfer) for better performance in Garo Hills, Meghalaya. Int. J. Adv. Res. Biol. Sci. 2020;7(10):11-23.
36. Khanikor DP, Dutta SK. Low temperature preservation of seed cocoons for delaying moth emergence in muga silkworm (*Antheraea assama* Westwood). Int. J Wild Silkworm Silk. 2000;5:216-220.
37. Prasad GK, Sinha SS. Induction of diapause in muga silkworm. Annual Report, Central Muga Eri Research & Training Institute, Jorhat, Assam; 1982.
38. Haniffa MA, Thatheyus AJ. Effect of age related mating schedule on reproduction in the silkworm, *Bombyx mori* L. Sericologia. 1992;32(2):217-225.
39. Khatri RK. Effect of seasonal variation on muga silkworm (*A. assama* Ww) rearing and grainage under different conditions of Doon Valley. Unpublished Ph.D. Thesis, HNB Garhwal University, Sri Nagar, Garhwal; 2003.
40. Sengupta K, Singh K. Some studies on the effect of refrigeration of muga *A. assamensis* cocoons on moth emergence, egg laying and hatching of eggs. First International Seminar on non-mulberry silks, CTR&TI, Ranchi. 1974;174.
41. Chowdhury SN, Muga Silk Industry. Directorate of Sericulture, Govt. of Assam, Guwahati. 1982;1-177.
42. Sahu AK, Singha BB, Kumar N. Preservation of seed cocoons during summer. Ann. Rep. RMRS, Boko, Assam; 1999.
43. Rajkhowa G, Kumar R, Rajan RK. Studies on the long-term preservation method of Muga Cocoon (*Antheraea assamensis* Helfer) at low temperature. Munis Entomology & Zoology. 2011;6(2):815-818.
44. Bora A, Samson MV, Borah B, Bhuyan N. Studies on preservation of muga cocoons of higher altitude and low temperature. Ann. Report RMRS, CSB, Boko, Assam. 1988-89;32-33.
45. Bora A, Singh PK, Borah B, Sengupta AK. Studies on preservation of muga cocoons at low temperature. Ann. Report RMRS, CSB, Boko, Assam. 1992; 10-11.

© 2022 Singh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/95886>