



## **Estimation of Optimum Plot Size and Shape from a Uniformity Trial for Field Experiment with Sunflower (*Helianthus annuus*) Crop in Soil of Hisar**

**Nishu Lohmor<sup>1</sup>, Mujahid Khan<sup>1\*</sup>, Kiran Kapoor<sup>1</sup> and Sudha Bishnoi<sup>1</sup>**

<sup>1</sup>*Department of Mathematics, Statistics and Physics, Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana, India.*

### **Authors' contributions**

*This work was carried out in collaboration between all authors. Author NL designed and carried out the study, performed the statistical analysis and wrote the protocol. Author MK wrote the first draft of the manuscript and managed the literature searches. Author KK guided the other authors during whole research period. Authors NL, MK and SB managed the analyses of the study. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJPSS/2017/31613

#### Editor(s):

- (1) Peter A. Roussos, Agricultural University of Athens, Lab. Pomology, Greece.  
(2) Fatemeh Nejatzadeh, Department of Horticulture, Faculty of Agriculture, Khoj Branch, Islamic Azad University, Iran.

#### Reviewers:

- (1) Wafula Nelson Wekha, Kenyatta University, Kenya.  
(2) Mustafa Turkmen, Giresun University, Turkey.

Complete Peer review History: <http://www.sciencedomain.org/review-history/18907>

**Original Research Article**

**Received 16<sup>th</sup> January 2017**  
**Accepted 28<sup>th</sup> February 2017**  
**Published 4<sup>th</sup> May 2017**

### **ABSTRACT**

A study of uniformity trial for assessing the nature and magnitude of soil variability and to determine the optimum size and shape of plots was conducted on 66A507 Pioneer hybrid of Sunflower crop at Research Farm of CCS Haryana Agricultural University, Hisar, Haryana during the February 2014 to June 2014, on a field of size 35 m × 40 m which after eliminating border effects reduced to 32 m × 36 m. The total area (1152 m<sup>2</sup>) divided into 1152 basic units, each have size 1 m × 1 m and yield data of all the basic units was recorded separately for further investigations. The coefficient of variation of yield of individual harvested units was observed to be as high as 13.92 per cent indicating high degree of soil heterogeneity. The coefficient of variation decreased with increase in plot size in both the directions *i.e.* when plots were elongated in N-S direction or elongated in E-W direction and the decrease was near about same for both the directions but was more when plots were elongated in N-S direction (96.48 per cent decrease). The long-narrow plots elongated in N-S direction were found to be more useful than the compact and square plots. It was observed that the smallest plot has the maximum efficiency and the optimum plot size was estimated to be 2 m<sup>2</sup>.

\*Corresponding author: E-mail: [mkhanstat@gmail.com](mailto:mkhanstat@gmail.com);

**Keywords:** Coefficient of variation; direction; optimum plot size and shape; sunflower; uniformity trial.

## 1. INTRODUCTION

The experimental material consists of certain variations which may be inherent like soil variability hence the agricultural field experiments are subject to high degree of error variation. This variability causes variations in the yield from plot to plot in the entire area even when the crops are grown in the similar sized plots and given same treatments, under exactly identical conditions. This sort of variation in the field experiments is measured by the coefficient of variation (CV). Coefficient of variation is directly proportional to the variation in soil fertility and hence high coefficient of variation indicates high variation in the soil fertility and low coefficient of variation indicates low variation in the soil fertility.

In practice, soil fertility has different magnitude for different sizes and shapes of plots. Thus for efficient planning of experiments, problem will be to find out the best possible sizes and shapes of the plots for experimentation, so that the error variation has minimum effect on treatment comparisons. The selection of suitable sizes and shapes of plots depends both on statistical consideration as well as practical feasibility. From statistical consideration, the estimate of treatment on a given experimental area should be obtained with maximum accuracy, and from a practical point of view, the plots should be sufficiently large so that the various field operations can be done correctly and probably reduce the experimental error.

For finding the suitable size and shape of plots arrangements that will be most accurate for estimating the treatment means for the given amount of experimental area, it is necessary to have an idea of the magnitude of the experimental error associated with different sizes and shapes of plots. This can be studied by conducting the uniformity trials on the crop in a given area. Numerous reports [1-6] suggest that optimum plot size for different crops of the region differ. Realizing the importance of sunflower (*Helianthus annuus*), being the third most important oilseeds crop in India after groundnut and mustard, the present study was undertaken to estimate the magnitude of the experimental error associated with the varying sizes and shapes of plots.

## 2. MATERIALS AND METHODS

The experiment of uniformity trial on sunflower hybrid 66A507 Pioneer was carried out at Research Farm, Department of Genetics and Plant Breeding, CCS Haryana Agricultural University, Hisar during February 2014 to June 2014. The uniformity trial was conducted over a field of area 35 m × 40 m. At the time of harvest, the experimental field was divided into rows (E-W direction) and columns (N-S direction). But to eliminate the border effects, the border area from all sides was left as non-experimental area, thereby making out net area of 32 m × 36 m at the centre of the field. Harvesting of the crop was done separately for each basic unit (1 m × 1 m) and the produce of each unit were recorded separately in grams for further investigation.

The adjacent basic units were combined to form plots of different shapes and sizes, and yield was recorded. These plots were formed by taking 1, 2, 4, 8 and 16 units along the rows (E-W direction) and also 1, 2, 3, 4, 6, 9, 12 and 18 units along the columns (N-S direction), thus having different shapes and sizes. Coefficient of variation for each size and shape of plot was calculated and the coefficient of variation so obtained was utilized to determine optimum size and shape of plots.

Relationship between CV and size and shape of plots was computed using [7], which states that

$$V_x = V_1 / X^b \quad (1)$$

Where;

$V_x$  is the variance of yield per unit area among plots of size X units,  
 $V_1$  is the variance among plots of size unity,  
 b is the linear regression coefficient and  
 X is the number of basic units per plot.

The relative efficiencies (R.E.) of different plot sizes were calculated using method suggested by [8]. Taking the efficiency of smallest plot as unity, the relative efficiencies of various plot sizes has been calculated.

$$R.E. = (CV_1 / CV_2)^2 \times (X_1 / X_2)^2 \quad (2)$$

Where;

$CV_1$  and  $CV_2$  are the coefficients of variation corresponding for plot sizes  $X_1$  and  $X_2$  respectively.

The optimum plot size was calculated using Maximum curvature method and Smith's variance law method. The maximum curvature method [9] has frequently been used to determine plot size for various field crops. The formula given by [9]

$$X_{opt}^{2(1+b)} = V_1^2 b^2 \{ [3(1+b)/(2+b)] - 1 \} \quad (3)$$

Smith [7] worked out optimum plot size for different values of costs under assumption of linear cost structure.

$$X_{opt} = \frac{bC_1}{(1-b)C_2} \quad (4)$$

Where;

$X_{opt}$  is the optimum plot size which provides the maximum information per unit of cost,  $C_1$  is that part of total cost which is proportional to no. of plots per treatment and  $C_2$  is that part of total cost which is proportional to the total area per treatment.

### 3. RESULTS AND DISCUSSION

The coefficient of variation of yields of harvested units for various sizes and shapes of plots is given in Table 1.

A high degree of variability i.e. 13.92 per cent was observed which indicates high degree of soil heterogeneity (Table 1). This variation further reduced with increase in plot size in either direction but the decrease was more when plots were elongated in N-S direction (96.48 per cent) than those elongated in E-W direction (95.83 per cent), indicating thereby that the plots become

more homogenous when elongated along N-S direction. This is in agreement with earlier reports of [10-16,1].

The coefficient of variation for various plot shapes for a given plot size have been calculated and are presented in Table 2.

The long-narrow plots elongated in N-S direction had less coefficient of variation than compact and square plots for a given particular plot size. Thus, best plot shape was 1:X, where '1' is the number of units in E-W direction and 'X' is the number of units in N-S direction. The same results were obtained by [13-15,17,18,6].

After having known the best shape, a functional relationship between plot size and coefficient of variation was examined by fitting the equation (1), which comes out to be

$$V_x = 24.78X^{-0.299} \quad (R^2=0.679)$$

The equation was in conformity with Smith's law, where the soil variability index (b) was 0.299, indicating the positive correlation between the adjacent basic units.

The relative efficiencies were computed using equation (2) and are presented in Table 3. Relative efficiency of smallest plot was highest but efficiency decreases as the plot size increases. Hence, smallest plot was most efficient but convenience of practical operation is to be given due attention.

The optimum plot size was worked out by maximum curvature method using equation (3) and was found to be 2 units i.e. 2 m<sup>2</sup>. The optimum plot sizes were also calculated by Smith's method using equation (4) and results are presented in Table 4. It was observed that the optimum plot size increases with the increase in cost ratio for a given plot arrangement.

**Table 1. Coefficient of variation for various plot sizes**

		No. of units in E-W direction							
		1	2	3	4	6	9	12	18
No. of units in N-S direction	1	13.92	8.45	7.71	7.08	4.36	1.62	1.75	0.58
	2	10.93	7.51	4.30	3.34	3.26	1.97	0.91	-
	4	8.20	6.14	6.26	5.18	3.77	1.61	-	-
	8	4.21	5.36	4.30	2.21	1.77	-	-	-
	16	0.49	-	-	-	-	-	-	-

**Table 2. Coefficient of variation for various plot sizes and plot shapes**

Plot size (in units)	Plot shape	CV (%)	Minimum CV (%)
1	1:1	13.92	13.92
2	1:2	8.45	8.45
	2:1	10.93	
3	1:3	7.71	7.71
4	1:4	7.08	7.08
	2:2	7.51	
	4:1	8.20	
6	1:6	4.36	4.30
	2:3	4.30	
8	2:4	3.34	3.34
	4:2	6.14	
	8:1	4.21	
12	1:12	1.75	1.75
	2:6	3.26	
	4:3	6.26	
16	4:4	5.18	0.28
	8:2	5.36	
	16:1	0.28	
18	1:18	0.05	0.05
	2:9	1.97	

**Table 3. Relative efficiencies of various plot sizes**

Plot size (in units)	Plot shape	C.V.	Relative efficiency
1	1:1	13.92	1
2	1:2	8.45	0.678
3	1:3	7.71	0.362
4	1:4	7.08	0.241
6	2:3	4.30	0.292
8	2:4	3.34	0.272
12	1:12	1.75	0.440
16	16:1	0.28	0.076
18	1:18	0.05	0.059

**Table 4. Optimum plot size under cost consideration**

Value of b = 0.299	
C <sub>1</sub> /C <sub>2</sub>	Optimum size of plot (m <sup>2</sup> )
0.5	0.214
1.0	0.428
2.0	0.855
3.0	1.283
4.0	1.710
5.0	2.138
6.0	2.565
7.0	2.993
8.0	3.420
9.0	3.848
10.0	4.275

## 4. CONCLUSIONS

The coefficient of variation decreases as the plot size increases in case of both the directions i.e. when plots were elongated in N-S direction or elongated in E-W direction. However, it was observed that the plots elongated in the N-S direction were more beneficial in controlling the soil heterogeneity than those elongated in the E-W direction. Relative efficiency of smallest plot was found maximum but efficiency decreases as the plot size increases. The optimum plot size worked out by maximum curvature method was found to be 2 units i.e. 2 m<sup>2</sup>. The optimum plot size increases with the increase in cost ratio for a given plot arrangement.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Shafi S, Mir SA, Nazir N, Rashid A. Optimum plot size for tomato by using S-PLUS and R-software's in the soils of Kashmir. *An Asian Jour of Soil Sci.* 2009;4(2):311-314.
- Storck L, Filho AC, Lopes SJ, Toebe M, de Silveira TR. Experimental plan for single, double and triple hybrid corn. *Maydica.* 2010;55:27-32.
- Patil SL, Reddy MN, Rao PB. Experimental plot size and shape based on data from a uniformity trial in dryland Bengalgram (*Cicer arietinum* L.) during winter season in the vertisols of semi-arid tropics of South India. *Indian Journal of Dryland Agriculture Research and Development.* 2010;25(1):102-105.
- Prajapati BH, Chaudhary GK, Chaudhary MK, Loria JM. Optimum size and shape of plot for field experiments on Mustard under North Gujarat condition. *Journal of Indian Society of Agricultural Statistics.* 2011;65(1):39-58.
- Masood MA, Raza I. Estimation of optimum field plot size and shape in paddy yield trial. *American-Eurasian Journal of Scientific Research.* 2012;7(6):264-269.
- Khan M, Hasija RC, Tanwar N. Optimum size and shape of plots based on data from a uniformity trial on Indian mustard in Haryana. *Mausam.* 2017;68(1):67-74.
- Smith HF. An empirical law describing heterogeneity in the yields of agricultural

- crops. Journal of Agricultural Science. 1938;28:1-23.
8. Agarwal KN, Deshpande MR. Size and shape of plots and blocks in field experiments with dibbled paddy. Indian Jour of Agril Sci. 1967;37(6):445-455.
  9. Agarwal KN. Uniformity trial on apple. Jour of Horti Sci. 1973;30:525-528.
  10. Kaushik LS, Daulta BS, Arora RK. A study of size and shape of plots and blocks in field experiment with mandarin orange. Haryana Jour of Horti Sci. 1974;3:124-133.
  11. Kaushik LS, Singh RP, Yadav TP. A preliminary study of size and shape of plots in field experiment with groundnut. Oilseeds Jour. 1976;613:20-25.
  12. Kaushik LS, Singh RP, Yadav TP. A uniformity trial with mustard. Ind Jour Agric Sci. 1977;47(10):515-518.
  13. Hasija RC, Kaushik LS, Mehta SL, Paroda RS. Uniformity trial in guar (*Cyamopsis tetragonoloba* Taub.). Forage Research. 1985;11(1):21-25.
  14. Kaushik LS, Kapoor K, Hasija RC, Tyagi CS. A uniformity trial on Moong. Indian Jour Agric Res. 1989;28(3):138-142.
  15. Kumar M, Hasija RC. A study in size and shape of plots with wheat (*Triticum aestivum* L.). Annals of Agric Bio Res. 2002;7(1):89-93.
  16. Kumar A, Kapoor K, Gupta SC, Hasija RC. Uniformity trial on sesame (*Sesamum indicum*). Environment and Ecology. 2007;255(2):295-298.
  17. Naliyadara CM, Upadhyay SM, Ramani CV. Optimum Size and Shape of Plots for Field Experiments in Tomato. Jour Ind Soc Agril Statist. 2005;59(1):34.
  18. Chaudhary GK, Prajapati BH, Patel JK, Prajapati RI, Loria JM. Optimum Size and Shape of Plot for Field Experiments on Wheat under North Gujarat Condition. Jour Ind Soc Agri Stat. 2011;65(1):39-58.

© 2017 Lohmor et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<http://sciencedomain.org/review-history/18907>