



***In-vitro* Screening of EMS and Sodium Azide Induced Mutant Population at Seedling Stage for Drought Tolerance in Wheat (*Triticum aestivum* L.)**

Ritu Rani ^{a*}, Manoj Kumar Yadav ^a, Akash Tomar ^b,
Vaishali ^a, Lokesh Kumar Gangwar ^c
and Rakesh Singh Sengar ^a

^a Department of Agriculture Biotechnology, College of Agriculture, S.V.B. Patel University of Agriculture and Technology, Meerut, Uttar Pradesh, India.

^b Department of Recombination Techniques, College of Biotechnology, S.V.B. Patel University of Agriculture and Technology, Meerut, Uttar Pradesh, India.

^c Department of Genetics and Plant Breeding, College of Agriculture, S.V.B. Patel University of Agriculture and Technology, Meerut, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i103104

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106924>

Original Research Article

Received: 14/07/2023

Accepted: 21/09/2023

Published: 28/09/2023

ABSTRACT

Drought caused by climate change result in water shortages, decreased global wheat production and unevenly distributed heavy rains. About 50% of the global wheat production is affected by water deficit conditions. The effects of different concentrations of EMS and SA on HD-3226 and HI-1620 wheat genotype for drought tolerance capability were investigated at seedling stage under

*Corresponding author: E-mail: rituchaudhary246@gmail.com;

water and osmotic stress conditions. In the present study, 15% concentration of polyethylene glycol (PEG-6000, to mimic drought stress) was used for in-vitro screening of EMS and SA induced mutant lines of both wheat genotypes. Current study revealed that SA decreased germination of seeds, seedling length, seed vigour index and survival rate percentage while it increased morphological mutation frequency percentage as comparison to EMS treatment in both wheat genotypes under water and PEG stress. Moreover, osmotic stress significantly reduced seed germination, shoot length, root length, seed vigour index, survival rate percentage compared to control-wild type.

Keywords: Water stress; seed vigour index; mutagen; wheat; drought stress; morphological characters.

ABBREVIATIONS

EMS : Ethyl Methane Sulphonate

SA : Sodium Azide

PEG : Polyethylene Glycol

1. INTRODUCTION

Wheat (*Triticum aestivum* L.) is a member of Triticeae tribe and Poaceae family. Since 10,000 years ago, tetraploid and hexaploid varieties of wheat have been domesticated [1] and Hexaploid form is modern day bread wheat and fulfils dietary needs of global population. By 2050, global population is expected to reach 10 billion, which would require double of current global food production [2]. Global climate change affects a variety of factors associated with water stress and extreme drought land area is likely to increase from 1 to 35% by year 2100 [3]. Bread wheat, being a rabi season crop is exposed to a number of environmental stresses namely, drought, cold, salinity and heat stress among which drought stress is major contributor to crop losses. Moisture or drought stress occurs at all stages irrespective of growth stages which largely depend on local environmental conditions [4]. Drought conditions particularly affect yield and quality of wheat (*Triticum aestivum* L.), one of the most important and widespread grain crops, indispensable in human nutrition and animal feed production [5]. Development of stress tolerant varieties is always a main objective of many breeding programs, but success has been limited by adequate screening techniques, and lack of genotypes that show clear differences in response to various environmental stresses [6]. Screening and identification of wheat genotypes that can tolerate water stress is important to boost wheat production which can be achieved by exploring genetic potential from available germplasm of wheat. Seed germination is first stage of growth that is sensitive to water deficit. Therefore, seed germination, vigour and coleoptile length are

rudiments for success of stand establishment of crop plants [7]. Thus there is a need to improve genetic tolerance of crops at seedling stages. Improvement in grain yield of wheat has traditionally relied on direct selection for this trait [8]. Genetic modification is best method for improving wheat production [9]. Induction of mutations by various mutagens is one way to generating large number of variability and has been successfully employed [10]. Mutant screening involves evaluation of a large number of mutant plants to identify rare mutant individuals that meet desired trait. The screening for drought tolerant lines of wheat under field condition is time consuming and labour intensive. Therefore, there is an urgent need to develop a simple and effective early screening method [11,12]. For drought stress induction, one of most popular approaches is to use high molecular weight osmotic substances, such as polyethylene glycol (PEG). These agents have no detrimental or toxic effects on plant [13]. They inhibit plant's growth by reducing water potential in a way similar to soil drying [14]. It affects how quickly plants' roots and shoots grow and generates osmotic stress in them [15]. The non-ionic, almost impermeable chains of PEG molecules (PEG 6000) maintain homogeneity of water potential throughout experiment without harming people's health [6]. The research on identifying drought resistant wheat genotypes have demonstrated that various features respond differently with varying amounts of PEG-6000 [16,17]. In several studies, it has been observed that PEG based screening may be a option to screen drought tolerant wheat lines at seedling stage. Therefore, present experiment was designed to screening M₁ seeds using PEG solution for drought tolerance in wheat combining with EMS and SA.

2. MATERIALS AND METHODS

Mature dry seeds of latest released varieties-HD-3226 (susceptible) and HI- 1620 (tolerance)

were used for the present investigation conducted during 2020-2021 in SVPUA&T, Meerut, UP, India. Seeds (M_0 seeds) of genotypes HD-3226 and HI-1620 were sterilized with 70% ethanol for 3-5 minutes at room temperature and then pre-soaked for 12 hours. Experiment was conducted during Rabi session 2020-2021 (M_1 generation). After overnight pre-soaking, one hundred fifty seeds (M_0) of each genotype were treated with four different concentrations of Ethyl Methane sulphonate (EMS)-0.25%, 0.5%, 0.75%, and 1% (v/v) and three different concentrations of sodium azide (SA)-0.02%, 0.04%, and 0.08% (w/v) for 2 hours were used. After mutagen treatment, treated seeds (M_1) seeds were transfer into a small cotton bag and these bags tie on stopcock to remove excess chemical mutagens under running tap water for 2 hours. Then, twenty-five air dried M_1 and non-treated seeds as control (Wild Type) were placed on moistened double whatman filter paper in each Petri plates with three replications in water and 15% PEG-6000 solution (w/v) for drought tolerance screening [18]. Five ml of 15% PEG solution was added to each Petri plates under osmotic stress conditions and distilled water was added to each Petri dish under normal conditions every 2 days to compensate for losses through evaporation up to 15 days [19]. When seedlings were at stage of

first true leaf initiation (after 15 days), successful germination in both water and 15% PEG solution, the germination data was collected for calculation of germination percentage, root and shoot length of the germinated seedlings were measured in centimetre before transferring in the small plastic pots and survival data along with morphologically mutant plants, was collected for survival percentage calculation at maturity of M_1 and M_2 generation, respectively. After 20 days, all plants were transferred in the field in a randomly block design (RBD). All treated (M_1) and control seeds (Wild Type) were sown in the field in three replications by maintaining row to row and plant to plant distance. The plant to plant distances was kept around 15 cm and spacing between adjacent rows was kept 20cm. The gap between two rows of different genotype was kept 50cm for maintaining proper distance from each genotype to another genotype. After transplanting of all plants, fields were irrigated at regular interval of 20-25 days. The crop was maintained in the field using conventional agronomic practices to keep crop in good condition. All plants (mutant as well as control- Wild Type) sown in field, were tagged properly with their genotype name along with number and other details. For drought screening at seedling stage data were calculated using following formulas:

1. Germination %: Germination % is an estimate of viability of a population of seeds. The germination % of seeds was measured by counting total germinated seeds from the total seeds were sown for the experiment after 15 days of sowing.

$$\text{Germination percentage (GP)} = \frac{\text{Seeds germinated}}{\text{Total seeds}} \times 100$$

2. Seedling Length (SLs): The length of seedlings was measured in centimetre from starting point of root tip to the top node of shoot using a ruler against a millimetre paper.

3. Seed Vigour Index [SI (%)]: The vigour index of each seed (control-WT and treated seed) was calculated using the formula which proposed by Abdul-Baki and Anderson [20] as germination percentage multiplied by seedling length.

$$\text{Seed Vigour Index (SI)} = \text{Germination percentage} \times \text{Seedling Length}$$

Where,
Seedling Length = Root length + Shoot length in cm,

4. Survival Rate % [SR]: The surviving plants in control and different treatments were counted at the time of maturity and calculated by following formula:

$$\text{Survival rate \%} = \frac{\text{Number of living seedlings}}{\text{Number of Total seedlings}} \times 100$$

5. Morphological Mutation Frequency [MF%]: Morphological mutations were screened throughout the growth period of plants and Gustafsson method [21] used for mutation frequency calculation.

$$\text{Morphological Mutation Frequency} = \frac{\text{Number of mutated plants}}{\text{Number of Total plants}} \times 100$$

Statistical analysis: The recorded data were subjected to analysis of variance based on Random Block Design (RBD) with two factors was performed by OPSTAT software.

3. RESULTS AND DISCUSSION

The in vitro screening of M₁ populations was carried out by germination of seeds on 15% PEG-6000 and analysing the mutagenic effects of EMS and Sodium azide were studied on seed germination, seedling length, seed vigour index, plant survival in M₁ generation while morphological mutation frequency in M₂ generation of both wheat genotype.

Germination %: In HD-3226 genotype, seed germination in control-wild type was 97.00% in water and 96.67% in 15% PEG (Table1). In case of EMS, it range from 94.67% (EMS 0.75%) to 98.33% (EMS 0.5%) and 88.67% (EMS 0.5%) to 94.00% (EMS 0.75%) in water and 15% PEG treatments, respectively. While SA decreased the germination of seeds as comparison to EMS, it range from 84.67% (SA 0.04%) to 89.33% (SA 0.02%) and 88.00% (SA 0.04%) to 90.33% (SA 0.08%) in water and 15% PEG treatments, respectively. The average mean value across all treatments in water was 97.08% (EMS) and 86.33% (SA) while it was 90.41% (EMS) and 89.22% (SA) in PEG. In HI-1620 genotype, seed germination in control-wild type was 97.33% in water and 98.67% in 15% PEG. In case of EMS, it range from 97.33% (EMS 0.75%) to 98.67% (EMS 0.25%, EMS 0.5% and EMS 1%) and 88.00% (EMS 1%) to 93.33% (EMS 0.5%) in water and 15% PEG, respectively. SA decreased the germination of seeds as comparison to EMS in water and it range from 88.00% (SA 0.04% and SA 0.08%) to 92.00% (SA 0.02%) while SA increased the germination of seeds as comparison to EMS in PEG and it range from 90.67% (SA 0.02% and SA 0.08%) to 96.00% (SA 0.04%). The average mean value across all treatments in water was 98.33% (EMS) and 89.33% (SA) while it was 90.66% (EMS) and 92.44% (SA) in PEG. The germination percentage decreased in PEG as comparison to water in EMS treatments while it increased in PEG as comparison to water in SA treatments according to average mean value of EMS and

SA in both wheat genotypes (Table1). In mutagenic treatments reduction in seed germination is due to delay or inhibition in physiological and biological processes necessary for seed germination which includes inhibition of mitotic process [22], enzyme activity [23] and hormonal imbalance [24]. Similar findings have also been reported, like reduction in germination rate with the increase PEG were noted in different crops [17,10,9]. Rajoriya *et al.*, also reported that seed germination decreased with increase in concentration of mutagens [25].

Seedling Length (cm): In HD-3226 genotype, seedling length in control-wild type was 33.90cm in water and 15.97cm in 15% PEG (Table 2). In case of EMS, it ranges from 22.81cm (EMS 0.5%) to 29.95cm (EMS 0.25%) and 16.62cm (EMS 1%) to 22.68% (EMS 0.75%) in water and 15% PEG treatments, respectively. While SA decreased the seedling length as comparison to EMS, it range from 15.73cm (SA 0.04%) to 22.23cm (SA 0.08%) and 11.91cm (SA 0.08%) to 17.74cm (SA 0.04%) in water and 15% PEG treatments, respectively. The average mean value across all treatments in water was 26.84cm (EMS) and 18.87cm (SA) while it was 20.38cm (EMS) and 15.19cm (SA) in PEG. In HI-1620 genotype, seedling length in control-wild type was 19.73cm in water and 19.12cm in 15% PEG. In case of EMS, it ranges from 23.78cm (EMS 1%) to 28.99cm (EMS 0.5%) and 15.96cm (EMS 1%) to 19.17cm (EMS 0.75%) in water and 15% PEG, respectively. SA also decreased seedling length as comparison to EMS, it range from 18.50cm (SA 0.08%) to 19.07cm (SA 0.02%) and 15.27cm (SA 0.04%) to 15.73cm (SA 0.02%) in water and 15% PEG treatments, respectively. The average mean value across all treatments in water was 26.22cm (EMS) and 18.87cm (SA) while it was 17.67cm (EMS) and 15.53cm (SA) in PEG. Seedling length decreased in PEG as comparison to water in both EMS and SA treatments according to average mean value of EMS and SA in both wheat genotypes. Roots are important in up taking water and nutrients, perceiving and transducing water deficit to shoots which will trigger different morpho-physiological responses and it was an important trait for selection of drought resistant genotypes. According to Fraser *et al.*, [26], reduction in root

and shoot lengths may be due to an impediment of cell division and elongation leading to a kind of tuberization which allow the stressed conditions to become favourable. Similarly, Rajoriya *et al.*, [24] reported that seedling length decreased with

increase in concentration of mutagens. Similar results also reported by Vuković *et al.*, [6] and Memon *et al.*, [27], seedling length of wheat cultivars significantly decreased under osmotic stress.

Table 1. Germination percentage of seeds of HD-3226 and HI-1620 wheat genotype after Ethyl Methane Sulphonate (EMS) and Sodium azide (SA) treatments in water and 15% PEG for *in-vitro* screening at seedling stage

Treatment	HD-3226			HI-1620		
	Water	PEG	Mean	Water	PEG	Mean
Control	97.00	96.67	96.83	97.33	98.67	98.00
EMS 0.25%	97.33	89.33	93.33	98.67	90.67	94.67
EMS 0.5%	98.33	88.67	93.50	98.67	93.33	96.00
EMS 0.75%	94.67	94.00	94.33	97.33	90.67	94.00
EMS 1%	98.00	89.67	93.83	98.67	88.00	93.33
SA 0.02%	89.33	89.33	89.33	92.00	90.67	91.33
SA 0.04%	84.67	88.00	86.33	88.00	96.00	92.00
SA 0.08%	85.00	90.33	87.67	88.00	90.67	89.33
EMS Mean	97.08	90.41	93.74	98.33	90.66	94.49
SA Mean	86.33	89.22	87.77	89.33	92.44	90.88
Total Mean	93.04	90.75		94.83	92.33	
	Factor (A)	Factor (B)	Factor (AxB)	Factor (A)	Factor (B)	Factor (AxB)
C.D.	3.87	1.94	5.48	3.52	1.76	4.98
SE(d)	1.89	0.94	2.67	1.72	0.86	2.43
SE(m)	1.33	0.67	1.89	1.21	0.61	1.72

Table 2. Seedling length (cm) of HD-3226 and HI-1620 wheat genotype after Ethyl Methane Sulphonate (EMS) and Sodium azide (SA) treatments in water and 15% PEG for *in-vitro* screening at seedling stage

Treatment	HD 3226			HI 1620		
	Water	PEG	Mean	Water	PEG	Mean
Control	33.90	15.97	24.93	19.73	19.12	19.47
EMS 0.25%	29.95	21.49	25.69	28.35	17.67	23.01
EMS 0.5%	22.81	20.76	21.78	28.99	17.91	23.47
EMS 0.75%	28.87	22.68	25.77	23.79	19.17	21.48
EMS 1%	25.74	16.62	21.18	23.78	15.96	19.87
SA 0.02%	18.65	15.93	17.29	19.07	15.73	17.40
SA 0.04%	15.73	17.74	16.74	19.05	15.27	17.16
SA 0.08%	22.23	11.91	17.076	18.50	15.60	17.05
EMS Mean	26.84	20.38	23.61	26.22	17.67	21.94
SA Mean	18.87	15.19	17.03	18.87	15.53	17.20
Total Mean	24.73	17.88		22.66	17.05	
	Factor (A)	Factor (B)	Factor (AxB)	Factor (A)	Factor (B)	Factor (AxB)
C.D.	1.19	0.59	1.68	1.93	0.97	2.75
SE(d)	0.59	0.29	0.82	0.94	0.47	1.33
SE(m)	0.41	0.20	0.59	0.66	0.33	0.94

Seed Vigour Index [SI (%)]: In HD-3226 genotype, seed vigour index in control-wild type was 32.86% in water and 15.44% in 15% PEG (Table 3). In case of EMS, it range from 22.43% (EMS 0.5%) to 29.13% (EMS 0.25%) and 14.92% (EMS 1%) to 21.32% (EMS 0.75%) in water and 15% PEG treatments, respectively. While SA decreased the seed vigour index as comparison to EMS, it range from 13.31% (SA 0.04%) to 18.89% (SA 0.08%) and 10.74% (SA 0.08%) to 15.63% (SA 0.04%) in water and 15% PEG treatments, respectively. The average mean value across all treatments in water was 26.03% (EMS) and 16.29% (SA) while it was 18.45% (EMS) and 13.54% (SA) in PEG. In HI-1620 genotype, seed vigour index in control-wild type was 19.20% in water and 18.87% in 15% PEG. In case of EMS, it range from 23.16% (EMS 0.75%) to 28.61% (EMS 0.5%) and 14.02% (EMS 1%) to 17.39% (EMS 0.75%) in water and 15% PEG, respectively. SA decreased the seed vigour index as comparison to EMS, it range from 16.23% (SA 0.08%) to 17.50% (SA 0.02%) and 14.15% (SA 0.08%) to 14.67% (SA 0.04%) in water and 15% PEG treatments, respectively. The average mean value across all treatments in water was 25.80% (EMS) and 16.84% (SA) while it was 16.04% (EMS) and 14.37% (SA) in PEG. The seed vigour index decreased in PEG as comparison to water in both EMS and SA treatments according to average mean value of EMS and SA in both wheat genotypes. Similar findings also reported by Arisandy *et al.* [28] and Nirmal-Raj *et al.* [29] in maize, seed vigour indices reduced under PEG induced drought.

Survival Rate % [SR]: In HD-3226 genotype, survival rate percentage in control-wild type was 99.33% in water and 99.00% in 15% PEG (Table 4). In case of EMS, it range from 87.67% (EMS 0.75%) to 98.52% (EMS 0.5%) and 89.06% (EMS 1%) to 93.17% (EMS 0.75%) in water and 15% PEG treatments, respectively. While SA decreased survival rate percentage as comparison to EMS, it range from 86.28% (SA 0.02%) to 87.78% (SA 0.08%) and 86.39% (SA 0.04%) to 87.78% (SA 0.02%) in water and 15% PEG treatments, respectively. The average mean value across all treatments in water was 95.01% (EMS) and 87.23% (SA) while it was 91.01% (EMS) and 87.28% (SA) in PEG. In HI-1620 genotype, survival rate percentage in control-wild type was 98.67% in water and 98.01% in 15% PEG. In case of EMS, it range from 89.05% (EMS 0.75%) to 96.89% (EMS 1%) and 87.03% (EMS 0.75%) to 91.17% (EMS

0.25%) in water and 15% PEG, respectively. SA decreased survival rate percentage in compare to EMS, it range from 90.11% (SA 0.04%) to 90.39% (SA 0.02% and SA 0.08%) and 88.13% (SA 0.02%) to 91.44% (SA 0.08%) in water and 15% PEG treatments, respectively. The average mean value across all treatments in water was 92.04% (EMS) and 90.29% (SA) while it was 89.22% (EMS) and 89.56% (SA) in PEG. The survival rate percentage decreased in PEG as comparison to water in both EMS and SA treatments according to average mean value of EMS and SA in both wheat genotypes. According to Natrajan and Shivshankar [30] and Sato and Gaul [21], reduction in seedling survival can be attributed to cytogenetic damage and physiological disturbances caused by mutagen treatment thus, seedling survivability may be hindrance caused by EMS and SA on different metabolic pathways of cells. Similar findings also reported by Rachovska and Dimova [31] in wheat, Khan *et al.* [32] in mungbean, Ilbas *et al.* [33] in barley and Kumar and Srivastava, [34] in Sesbania. Rajoriya *et al.* [24] also reported that plant survival decreased with increase in concentration of mutagens.

Morphological Mutation Frequency [MF%]: In present study, several morphological mutants showing variations in characters like altered plant height, leaf architecture, growth habit, tillers per plant, spike length and morphology etc. were isolated in M₂ generation of wheat genotypes. In HD-3226 genotype, morphological mutation frequency of EMS range from 3.38% (EMS 0.5%) to 4.70% (EMS 0.75%) in water and 3.44% (EMS 0.25%) to 5.33% (EMS 0.75%) in water and 15% PEG treatments, respectively (Table 5). While SA increased morphological mutation frequency as comparison to EMS, it range from 3.63% (SA 0.04%) to 4.84% (SA 0.08%) and 3.66% (SA 0.08%) to 6.28% (SA 0.04%) in water and 15% PEG treatments, respectively. The average mean value across all treatments in water was 4.14% (EMS) and 4.42% (SA) while it was 4.20% (EMS) and 4.88% (SA) in PEG. In HI-1620 genotype, morphological mutation frequency of EMS range from 2.81% (EMS 0.25%) to 4.68% (EMS 0.75%) in water and 3.04% (EMS 0.5%) to 4.92% (EMS1%) in water and 15% PEG treatments, respectively. While SA increased morphological mutation frequency as comparison to EMS, it range from 3.39% (SA 0.02%) to 4.56% (SA 0.08%) and 4.03% (SA 0.04%) to 5.95% (SA 0.02%) in water and 15% PEG treatments, respectively. The average mean value across all treatments in water was 4.05%

(EMS) and 4.17% (SA) while it was 4.13% (EMS) and 4.84% (SA) in PEG. The morphological mutation frequency increased in PEG as comparison to water in both EMS and SA treatments according to average mean value of EMS and SA in both wheat genotypes. Tokar [35] reported that morphological mutants are useful in gene mapping and phylogenetic studies

of crops. Khursheed *et al.*, [36] reported highest morphological mutation frequency in gamma rays and lowest with combined treatments in faba beans. Similarly, Raina and Khan, [37] also reported the highest mutation frequency with combination of gamma rays and sodium azide (5.86%) in cowpea.

Table 3. Seed Vigour Index of HD-3226 and HI-1620 wheat genotype after Ethyl Methane Sulphonate (EMS) and Sodium azide (SA) treatments in water and 15% PEG for *in-vitro* screening at seedling stage

Treatment	HD-3226			HI-1620		
	Water	PEG	Mean	Water	PEG	Mean
Control	32.86	15.44	24.15	19.20	18.87	19.03
EMS 0.25%	29.13	19.17	24.15	27.97	16.03	22.00
EMS 0.5%	22.43	18.41	20.42	28.61	16.75	22.68
EMS 0.75%	27.35	21.32	24.33	23.16	17.39	20.27
EMS 1%	25.24	14.92	20.08	23.47	14.02	18.75
SA 0.02%	16.69	14.25	15.47	17.50	14.29	15.90
SA 0.04%	13.31	15.63	14.47	16.81	14.67	15.74
SA 0.08%	18.89	10.74	14.81	16.23	14.15	15.19
EMS Mean	26.03	18.45	22.24	25.80	16.04	20.92
SA Mean	16.29	13.54	14.91	16.84	14.37	15.60
Total Mean	23.24	16.23		21.62	15.77	
	Factor (A)	Factor (B)	Factor (AxB)	Factor (A)	Factor (B)	Factor (AxB)
C.D.	1.43	0.72	2.02	1.92	0.96	2.72
SE(d)	0.70	0.35	0.99	0.94	0.47	1.33
SE(m)	0.49	0.25	0.70	0.66	0.33	0.94

Table 4. Survival rate percentage of HD-3226 and HI-1620 wheat genotype after Ethyl Methane Sulphonate (EMS) and Sodium azide (SA) treatments in water and 15% PEG for *in-vitro* screening at seedling stage

Treatment	HD-3226			HI-1620		
	Water	PEG	Mean	Water	PEG	Mean
Control	99.33	99.00	99.16	98.67	98.01	98.33
EMS 0.25%	97.97	91.45	94.69	90.44	91.17	90.80
EMS 0.5%	98.52	90.39	94.45	91.78	88.33	90.07
EMS 0.75%	87.67	93.17	90.42	89.05	87.03	88.03
EMS 1%	95.89	89.06	92.47	96.89	90.38	93.67
SA 0.02%	86.28	87.78	87.03	90.39	88.13	89.22
SA 0.04%	87.63	86.39	87.02	90.11	89.11	89.62
SA 0.08%	87.78	87.67	87.73	90.39	91.44	90.91
EMS Mean	95.01	91.01	93.01	92.04	89.22	90.63
SA Mean	87.23	87.28	87.25	90.29	89.56	89.92
Total Mean	92.69	90.63		92.27	90.45	
	Factor (A)	Factor (B)	Factor (AxB)	Factor (A)	Factor (B)	Factor (AxB)
C.D.	3.94	1.96	5.59	2.77	1.36	N/A
SE(d)	1.92	0.95	2.74	1.39	0.64	1.89
SE(m)	1.35	0.67	1.92	0.94	0.47	1.39

Table 5. Morphological mutation frequency of HD-3226 and HI-1620 wheat genotype after Ethyl Methane Sulphonate (EMS) and Sodium azide (SA) treatments in water and 15% PEG for *in-vitro* screening at seedling stage

Treatment	HD- 3226			HI- 1620		
	water	PEG	Mean	water	PEG	Mean
EMS 0.25%	4.32	3.44	3.88	2.81	4.56	3.68
EMS 0.5%	3.38	3.47	3.42	4.14	3.04	3.59
EMS 0.75%	4.70	5.53	5.11	4.68	4.03	4.36
EMS- 1%	4.18	4.65	4.41	4.57	4.92	4.74
SA- 0.02%	4.79	4.69	4.74	3.39	5.95	4.67
SA- 0.04%	3.63	6.28	4.95	4.54	4.03	4.29
SA- 0.08%	4.84	3.66	4.25	4.56	4.54	4.55
EMS Mean	4.14	4.2	4.17	4.05	4.13	4.09
SA Mean	4.42	4.88	4.68	4.17	4.84	4.50
Total Mean	4.266	4.535		4.105	4.442	
C.D.	1.026	N/A	1.45	0.832	N/A	1.176
SE(d)	0.496	0.265	0.702	0.402	0.215	0.569
SE(m)	0.351	0.188	0.496	0.285	0.152	0.402

4. CONCLUSION

Water is essential for seed germination, seedling growth, vegetative period of crop, flowering at translocation of minerals and nutrition incorporate throughout the plants, from root to leaf and vice versa in the plants [38]. Water stress is a major limiting factor for crop production and estimated the 50% of the global wheat production is affected by water deficit conditions [39]. Crops produced in countries with frequent drought can become more drought-resistant through mutation breeding. In vitro screening method using PEG has been proved to be very effective method for studying the effect of water stress on seed germination and seedling growth characters [40, 41, 42] and simple cost effective method to screen large set of germplasm within very less time period and accurately [43]. However, artificial induction of drought using PEG is dependent on concentration and varies with crop and genotype. In present investigation, osmotic stress significantly reduced seed germination, shoot length, root length, seed vigour index, survival rate percentage compared to the control-wild type. SA decreased germination of seeds, seedling length, seed vigour index and survival rate percentage while it increased morphological mutation frequency percentage as comparison to EMS according to average mean value of EMS and SA in both wheat genotypes genotype under water and 15% PEG stress condition. Considering the present data, variety HD-3226 was drought susceptible, while variety HI-1620 showed greater drought stress tolerance, showing better germination under osmotic stress.

ACKNOWLEDGEMENT

The authors gratefully acknowledge the institution for providing all inputs for completing trials and Haryana State Seed Certification Agency, Panchkula, Haryana for providing seeds of wheat genotypes used in study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Dubcovsky J, Dvorak J. Genome plasticity a key factor in the success of polyploid wheat under domestication. *Science*. 2007;316:1862–1866.
2. Mahpara S, Zainab A, Ullah R, Kausar S, Bilal M, Latif MI. The impact of PEGinduced drought stress on seed germination and seedling growth of different bread wheat (*Triticum aestivum* L.) genotypes. *PLoS ONE*. 2022;17(2): 0262937.
3. Kumar P, Kumar P, Vashali, Singh SK, Tomar A. Evaluation of drought resistance indices and grain yield in basmati and non-basmati rice under water stress at reproductive stage. *The Pharma Innovation Journal*. 2022;11(9):2081-2085.
4. Manoj NV, Chaudhary HK, Singh K. Screening of potential wheat genotypes for drought tolerance using polyethylene glycol. *Himachal Journal of Agricultural Research*. 2020; 46 (2): 130-135.

5. Javadinejad S, Dara R, Jafary F. Analysis and prioritization the effective factors on increasing farmers resilience under climate change and drought. *Agric. Res.* 2021;10: 497–513.
6. Vuković R, Camagajevac I.Š, Vuković A, Šunić K, Begović L, Mlinarić S, Sekulić R, Sabo N, Španić V. Physiological, Biochemical and Molecular Response of Different Winter Wheat Varieties under Drought Stress at Germination and Seedling Growth Stage. *Antioxidants.* 2022; 11: 693.
7. Khakwani, Abdul A, Dennett MD, Munir M. Early growth response of six wheat varieties under artificial osmotic stress condition. *Pak. J. Agri. Sci.* 2011; 48(2):119-123.
8. Braun HJ, Pfeiffer WH, Pollmer WG. Environments for selecting widely adapted spring wheat. *Crop Sci.*1992; 32:1420-1427.
9. Sharma V, Kumar A, Chaudhary A, Mishra A, Rawat S, Shami V, Kaushik P. Response of wheat genotypes to drought stress stimulated by PEG. *Stresses.* 2022; 2(1): 26-51.
10. Kadhimi AA, Alhasnawi AN, Isahak A, Ashraf MF, Mohamad A, Mohtar, WWY. Gamma radiosensitivity study on MRQ74 and MR269, two elite varieties of rice (*Oryza Sativa L.*). *Life Sci J.* 2016; 13(2):85-91.
11. Kumar P, Prasad B D, Sahni S. In vitro screening of gamma rays induced mutant population in rice for peg-induced drought stress. *Journal of Pharmacognosy and Phytochemistry.* 2018; 7(2): 3274-3277.
12. Barshile J. Chemical mutagenesis induced small leaf mutant in chickpea (*Cicer arietinum L.*). *International Journal of Researches in Biosciences and Agriculture Technology.* 2016;1(01).
13. Tabatabai SMT, Goshasbi F, Bakhshi B. Evaluation of the effect of polyethylene glycol (PEG) on germination and morphological characteristics of bread wheat. *Cereal Research Communications.* 2022; pp. 1-7.
14. Bressan RA, Hasegawa PM, Handa AK. Resistance of cultured higher plant cells to polyethylene glycol-induced water stress. *Plant Sci Lett.* 1981; 21:23-30.
15. Latif M, Bukhari SAH, Alrajhi AA, Alotaibi FS, Ahmad M, Shahzad AN, Mattar MA. Inducing drought tolerance in wheat through exopolysaccharide producing rhizobacteria. *Agronomy.* 2022; 12(5): 1140.
16. Abro AA, Akher SA, Memon S, Huda MN, Abro SA, Jahan SN. Influence of polyethylene glycol (PEG 6000) generate osmotic stress on seed germination of different wheat (*Triticum aestivum L.*) Genotypes. 2021;
17. Batool M, El-Badri AM, Wang Z, Mohamed IA, Yang H, AX, Zhou G. Rapeseed Morpho-Physio-Biochemical Responses to Drought Stress Induced by PEG-6000. *Agronomy.* 2022;12(3): 579.
18. Basha MH, Mehta AK. Screening of oat (*avenasativa l.*) mutant lines for drought tolerance using poly ethylene glycol-6000 at seedling stage. *Progressive Research – An International Journal.* 2016; 11 (VIII): 5561-5569.
19. El-Siddig MA, Baenziger S, Dweikat I, El Hussein A. Preliminary screening for water stress tolerance and genetic diversity in wheat (*Triticum aestivum L.*) cultivars from Sudan. *Journal of Genetic Engineering and Biotechnology.* 2013; 11: 87-94.
20. Abdul-Baki AA, Anderson JD. Vigour determination of soybean seed by multiple criteria. *Crop Sci.* 1973;13: 630-633.
21. Gustafsson A. (1940). The mutation system of the chlorophyll apparatus. *Lund Univ.*
22. Sato M, Gaul H. effect of ethylene methane sulphonate on the fertility of barley. *Radiation Botany.* 1967; 7: 7–15.
23. Chrispeeds MJ, Varner JE. Gibberelic acid induced synthesis and release of α -amylase and ribonuclease by isolated barley aleurons layers. *Plant Physiol.*1976; 42: 346–406.
24. Ananthaswamy HM, Vakil VK, Shrinivas A. Biological and physiological changes in gamma irradiated wheat during germination. *Radiation Botany.* 1971;11:1-12.
25. Rajoriya CM, Ahmad R, Rawat RV, Jat BL. Studies on induction of mutation in Fenugreek (*Trigonella fonum-graecum*). *International Journal for Research in applied science & engineering Technology.* 2016; 4(5): 333-375.
26. Fraser TE, Silk WK, Rost TL. Effects of low water potential on cortical cell length in growing regions of maize roots. *Plant Physiol.* 1990;93:648-651.
27. Memon S, Abro AA, Jakhro MI, Farid A, Habib M, Ahmed M, Bhutto LA, Memon SA, Farooq M. Polyethylene glycol

- mediated osmotic stress impacts on growth and biochemical aspects of wheat under artificial osmotic stress condition. Journal of Innovative Sciences. 2023; 9(1): 44-50.
28. Arisandy P, Bayuardi SW, Azrai M. Evaluation of drought tolerance in maize hybrids using stress tolerance indices. 2017; Int J Agron Agri R. 11: 46-54.
 29. Nirmal Raj R, Gokulakrishnan J, Prakash M. Assessing drought tolerance using PEG6000 and molecular screening by SSR markers in maize (*Zea mays* L.) hybrids. Maydica electronic publication. 2019; 64(9): 1-7.
 30. Natrajan AT, Shivshankar G. Studies on modification of mutations responses of barley seeds to ethyle methane sulphonate on the fertility of barley. Radiation Botany. 1965; 7: 7-15.
 31. Rachovska G, Dimova D. Effect of sodium azide and gamma rays on M1 quantitative characteristics of the productivity and their connection with M2 mutation changes in winter common wheat. Rasteniiev dni-nauki. 2000; 37: 413– 419.
 32. Khan S, Wani MR, Kumar P. Induced genetic variability for quantitative traits in *Vigna radiate* (L.) Wilczek. Pak J Bot. 2004; 36: 845–850.
 33. Ilbas EY, Eroglu H. Effects of the application of different concentrations of nan3 for different times on the morphological and cytogenetic characteristics of barley (*Hordeum vulgare* L.) seedlings. J Integr Plant Biol. 2005; 47: 1101–1106.
 34. Kumar G, Srivastava N. Efficiency and Effectiveness of Gamma Rays and Sodium Azide in *Sesbania cannabina* Poir. Cytologia. 2013; 78(1): 1–10
 35. Toker C. A note on the evolution of kabuli chickpeas as shown by induced mutations in (*Cicer reticulatum* Ladizinsky.) Genet. Resources Crop Evol. 2009; 56: 7–12.
 36. Khursheed S, Raina A, Parveen K, Khan S. Induced phenotypic diversity in the mutagenized populations of fababean using physical and chemical mutagenesis. J. Saudi Society Agric. Sci. 2019; 18: 113–119.
 37. Raina A, Khan S. Mutagenic effectiveness and efficiency of gamma rays and sodium azide in M2 generation of Cowpea [*Vigna unguiculata* (L.) Walp.] 2020. DOI: <https://doi.org/10.1101/2020.03.09.983486>.
 38. Kijne JW. Abiotic stress and water scarcity: identifying and resolving conflicts from plant level to global level. Field Crops Research. 2006; 97(1):3-18.
 39. Wang W, Vinocur B, Altman A. Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. Planta. 2003; 218 (1):1-4.
 40. Hadas A. Water uptake and germination of leguminous seeds under changing external water potential in osmoticum solution. J Exp Bot. 1976; 27:480-489.
 41. VandenBerg L, Zeng YJ. Response of South African indigenous grass species to drought stress induced by polyethylene glycol (PEG) 6000. Afr J Bot. 2006; 72:284-286
 42. Radhouane L. Response of Tunisian autochthonous pearl millet (*Pennisetum glaucum* (L.) R. Br.) to drought stress induced by polyethylene glycol (PEG) 6000. Afric J Biotechnol. 2007; 6:1102-1105.
 43. Kulkarni M, Deshpande U. In Vitro Screening of Tomato Genotypes for Drought Resistance Using Polyethylene Glycol. Afric J Biotechnol. 2007; 6(6):691-696.

© 2023 Rani et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/106924>