



The Role of Nutrition Interventions and Mothers Knowledge on Infant and Young Child Feeding: Mapping Study in Njombe and Geita Regions, Tanzania

Hassan Tearish Berenge^a
and Happiness Samuel Muhimbula^{b*}

^a *Sokoine University of Agriculture, Department of Policy Planning and Management, P.O Box-3035, Morogoro, Tanzania.*

^b *Sokoine University of Agriculture, Department of Human Nutrition and Consumer Science, P.O Box-3006, Morogoro, Tanzania.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/EJNFS/2023/v15i101349

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106762>

Original Research Article

Received: 24/07/2023

Accepted: 02/10/2023

Published: 11/10/2023

ABSTRACT

Children under the age of five are at risk of undernutrition particularly in developing countries. It was discovered that mothers' nutrition knowledge was one of the most influential factors in determining whether or not children under the age of five had an optimal nutritional status. The purpose of this study was to examine the roles of nutrition interventions and maternal knowledge regarding IYCF practices and child nutrition status in regions with high rates of stunting. This was a cross-sectional study conducted in the Tanzanian regions of Njombe and Geita from a community-based perspective and examining current and recently ended (within previous 3 months) nutrition

*Corresponding author: Email: happy.issa@sua.ac.tz;

interventions. A sample of 150 mothers with their children were selected purposefully to participate in the study. Women and their children younger than five years old were chosen purposively to participate in the study. The Chi-square test ($p < 0.05$) was used to compare IYCF practices, maternal knowledge with nutrition status, and differences between areas with few and many nutrition-related interventions. According to the results of this study, mothers' IYCF practices ($p = 0.014$) and their children's nutritional status ($p = 0.048$) improved significantly when nutrition interventions were easily accessible and readily usable. $P = 0.005$ revealed that 90.5% of teenage mothers (15–20 years old) had inadequate nutrition knowledge. Approximately, 68 % [49.7-86.3] of stunted infants in Njombe district were male, and there were few interventions in place, whereas the Bukombe district had a stunted infant rate of 38.9 % Multicomponent nutrition interventions are essential to strengthen programmes that address poor maternal knowledge, improve IYCF practices and nutrition status in regions with high chronic malnutrition.

Keywords: Intervention; nutrition status; stunting; infant feeding practices; undernutrition.

1. INTRODUCTION

Child undernutrition remains a global health problem with the goal that no country is harmless (GNR, 2018a). Undernutrition has destructive effects commonly in developing countries compared to developed countries [1]. Tanzania is one of the countries with a high rate of undernutrition, with approximately 31.8 percent of children under the age of five stunted [2].

WHO recommends IYCF indicators as the assessment guideline for acquiring suitable feeding practices information based on children's age groups [3]. Indicators are early initiation of breastfeeding (BF) within 1 hour after birth; exclusive BF without any other foods or drinks under 6 months; continued BF at age of 12-14.9 months; introduction of solid, semi-solid or soft foods at 6-8 months; minimum dietary diversity, meal frequency and acceptable diet at 6-23.9 months; and consumption of iron-rich/ iron-fortified food at age of 6-23.9 months [4]. The optimal feeding practices can be met through achieving all the recommendations.

Poor maternal nutritional knowledge, on the other hand, is one of the drivers of improper Infant and Young Child Feeding (IYCF) practices that lead to child malnutrition [5]. The regions with the high prevalence rate of chronic malnutrition in Tanzania had poor maternal nutrition knowledge with suboptimal IYCF practices [6]. Findings from other cohort studies observed that inadequate IYCF practices, maternal education, poor care, diseases, poor access to health services, and socio-economic status are among the major causes of undernutrition [7,8,9]. Maternal nutrition knowledge is likely to affect child feeding practices which both influence the nutrition status of children aged (0-56) months of age [10,11,12].

Government and non-governmental organizations are now focusing on the role of nutritional-sensitive programs that will raise healthier maternal and children's nutrition status [6,13,14,15]. According to the Tanzania Health and Demographic Survey (TDHS-MIS) of 2015/2016, various efforts have been made to reduce child malnutrition since 1992, when about 50% of children aged under five years were stunted compared to 31.8% in the current report. Furthermore, the Global Nutrition Report of 2018 revealed that Tanzania was among the countries with a high unacceptable burden of child malnutrition and its progress is currently simply not good enough. Due to the fact that multisectoral approaches to nutrition are required to achieve sustainable improvements in nutritional status, a great deal of effort is already being expended to achieve global and national strategic goals; therefore, it is time to determine which areas and fields of nutrition must be prioritized in order to meet the target (SDG 2.2). From that general point of view, this study hypothesized the potential favourable effects of nutrition interventions on maternal nutritional knowledge, IYCF practices, and child nutrition status.

In light of the limited existing evidence pertaining to strategies for expediting the reduction of child chronic malnutrition, the main focus of this study was to examine the potential effects of nutrition interventions on strengthening maternal nutrition knowledge, with the subsequent goal of improving the nutritional status. The Excel-based Scaled-Up Nutrition Project Monitoring Tool (SUN-PMT) was used for mapping and assessing current nutritional interventions by computing the coverage while determine their effect on maternal nutrition knowledge to the IYCF practices (Waiswa *et al.*, 2019).

2. METHODOLOGY

2.1 Description of Study Areas

The assessment and mapping of nutrition interventions were concentrated on the regions with a high rate of malnutrition. The regions and their respective districts were selected purposively based on the high prevalence rate of malnutrition. Njombe and Geita are among the mostly nutritional vulnerable regions in Tanzania with 49% and 41% rates of chronic malnutrition, respectively [11]. About 60 % of study participants, were farmers in both regions. The study areas showed to have different ethnicity and while variation in other socio-demographic features was minimal to justify the outcomes. The household survey was conducted in two representative districts such as Njombe Rural and Bukombe with coverage areas of 3448 km² and 10 842 km² respectively [17,18].

2.2 Study Design

A cross-section study was adopted for assessment of the efforts made by nutrition interventions on infant and young child feeding practices and the nutrition status of children less than 5 years of age. For the large extent and nature of this study, the cross-section is comparatively less time-consuming and inexpensive during the survey, using easy survey procedures and giving occurrence rates of various features [19].

2.3 Sampling Procedure and Study4 Study Population and Sampling Procedure

In this study, a multi-stage sampling method was used to obtain eligible respondents within the study areas. The two regions (Geita and Njombe) with a high rate of malnutrition were purposefully selected. The same was true for one district from each of the selected regions (Bukombe and Njombe DCs). A list of eligible households in selected villages was made, and if the total number of households in a village was higher than the sample size needed for a certain area, the interval of households visited was found by

dividing the total number of households by the sample size needed in that village.

Women or caretakers with children younger than five years old living in rural areas of the Bukombe and Njombe Rural Districts and were part of nutrition-related interventions were selected for study. Some of the nutrition project officers who work in the study areas and regional or district nutrition officers were also part of the study. There are 20 594 women of childbearing age in Njombe Rural and 51 153 in Bukombe. There is a 60% chance that a woman in Njombe Rural and an 86% chance that a woman in Bukombe will have a child under the age of five (URT, 2013).

2.4 The Sample Size

Accordingly, the Green formula was used to select the most statistically significant samples of household surveys possible. The number of predictors, *p*, determines the value of the constant *n*. Although the sample size is corrected as follows, it is expected that the number of predictors in this study will not exceed 12, based on the objectives and key questions.

$$\text{In this case, } N = 50 + 8(12) = 146.$$

A minimum sample size of 20 percent is sufficient for social science studies, according to Simonsohn *et al.* (2011), while the same authors presented at a conference suggested using a sample size of $N > 50$. [20]. The sample has grown to 150 after an additional four people were recruited and distributed equally among the two districts. The population size of Njombe District and Bukombe District dictated that the sample size was 65 and 85, respectively. In addition, other scientists believe that a sample size of 100 respondents or more is sufficient for statistically significant analysis [21,22]

2.5 Methods of Data Collection

2.5.1 Data collection and interviews

Maternal and child demographics, anthropometric assessment and mapping of project with pro nutrition interventions were done

Table 1. Average Score measurement scale

Code	Very Poor Knowledge (1)	Poor Knowledge (2)	Fair Knowledge (3)	Good Knowledge (4)	Excellent Knowledge (5)
Average Score (AS) value	0.00 – 1.80	1.81 – 2.60	2.61 – 3.40	3.41 – 4.20	4.21 – 5.00

Example of coding adopted on this study: AS ≤ 3.00 = 0 which means "No/Poor knowledge, while AS ≥ 3.10 means "Knowledgeable".

through face-to-face interviews and physical measurements.

2.5.2 Child's health and mother's socio-economic and demographic features

A pretested structured questionnaire was administered to eligible mothers with children aged under 5 years as a useful tool for recording quantitative information such as: mother's age, marital status, education, occupation, and maternal nutritional knowledge; child's age, sex, delivery place, and birth weight; and health or/and nutritional services availability and willingness assessment.

2.5.3 Maternal knowledge on infant and young child feeding (IYCF) practices (Average score measurement)

WHO recommends IYCF indicators as the assessment guideline for acquiring suitable feeding practices information based on children's age groups [3]. Indicators are early initiation of breastfeeding (BF) within 1 hour after birth; exclusive BF without any other foods or drinks under 6 months; continued BF at age of 12-14.9 months; introduction of solid, semi-solid or soft foods at 6-8 months; minimum dietary diversity, meal frequency and acceptable diet at 6-23.9 months; and consumption of iron-rich/ iron-fortified food at age of 6-23.9 months [4]. Individual dietary diversity score data, on the other hand, was transformed from a 24-hour recall assessment tool designed for complementary feeding practices [23].

A semi-structure questionnaire was used to obtain data on maternal nutrition knowledge. The questions were developed from FAO guideline for assessing nutrition related knowledge and attitudes, practices (KAP) [24]. A total question asked to mothers were 17 including those on early breastfeeding practices, complementary feeding practices and the risk associated with poor feeding practices. For each question correctly answered, 1 point was scored whereas wrong or I dont know answers was scored 0. The average score was adopted for assessing maternal nutrition knowledge. Then the average score was related and measured on the scale for further analysis.

2.5.4 Infant and Young Child Nutrition Status (Anthropometric)

Anthropometric measurements were used to assess the nutritional status of children. Weight

and length height were recorded for children using standard procedures [25]. The age of the children was recorded from their clinical cards/records. Infant and young children's weights were measured using a Seca 874 weighing scales wearing minimal clothing. Infant weight was measured using the "tared weighing" method while held by their mother. Infant recumbent length and height were measured using the UNICEF Portable length board and height measuring system SET-2 (S0114540) following standard procedures [25]. All anthropometric equipment was calibrated prior to use. Length /height -for-age z-score (LAZ/HAZ), weight-for-age z-score (WAZ), and weight-for-length / height z-score (WLZ /WHZ) were calculated using ENA for SMART Software (Software for Emergency Nutrition Assessment) [26]. Stunting, underweight and wasting were defined as z-scores below minus 2 standard deviations (SD) of the median values of the reference data.

2.5.5 Mapping of nutritional related interventions

The SUN-PMT tool was used to identify project-level interventions that have the potential to improve maternal nutritional knowledge, IYCF and nutritional status. Three approaches to data collection were employed in the mapping exercise; first, the direct one-to-one interview with project staff at the project office. Second, phone interviews with the project staff (especially if it was impossible to meet the staff in their office). Third, search for supplementary information from the project's websites where this was available. Fourth, the researcher read through the documents provided by the project staff which was available to supplement information collected through interviews. The assessment looked at details as to how the interventions improved maternal nutritional knowledge. The comparison of child nutritional status before (baseline data) and after (based on the status of existing interventions) by their association with maternal nutritional knowledge and IYCF practices was done to have outcome information [27]. The outcome data were obtained from the Tanzania Demographic Health Survey and Malaria Indicator Survey report of 2015, Tanzania National Nutrition Survey report of 2019, publishable and existing evaluation project reports as well as data and information from the regional nutrition officer, District nutrition officer, or authorized Project officer.

2.6 Data Processing and Analysis

The Statistical Package for Social Science (IBM SPSS 21) was used to enter raw data from structured and tested questionnaires. Despite the fact that anthropometric data were initially entered into software known as Emergency Nutrition Assessment (ENA) for Standardized Monitoring and Assessment of Relief Transitions (SMART) 2011, HAZ, WAZ, and WHZ were transformed from ENA for SMART software to IBM SPSS Statistics 21 for data cleaning and further analysis under WHO flags. A p-value of less or equal to 0.05 was considered a statistically significant difference between variables. Socio-economic and demographic features variables, IYCF, and child nutrition status indicators were both descriptively computed. Continuous data were checked for normality, tested and presented as mean and standard deviation; categorical data were computed and presented as Pearson Chi-square significance value; and anthropometric data (HAZ, WAZ, and WHZ) were also presented as the proportion of events that occurred with a 95% confidence interval (% [CI 95%]). The WHO standards classify stunted, underweight, and wasting when a z-score value is smaller than a negative 2 standard deviation of the median values of height for age, weight for age, and weight for height, respectively.

$$\text{Therefore, CI} = P \pm Z * (P(1-P)/n)^{0.5}$$

Where by p = sample proportion, Z = value from standard normal distribution of 95% N = Sample size and n = number of positive cases.

Lastly, mapping activity was additionally done through the Scaled-Up Nutrition Project Monitoring Tool (SUN-PMT) for appraisals and justifications of the effort made by nutritional interventions on the role of maternal nutritional knowledge in child nutritional status [28].

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Socio-economic and demographic characteristics of infants and mothers

The findings from Table 2 indicated that more than half (56.7%) of the children were females. The majority of children were an average of

23.96 months of age and had an average body weight of 3.21 kg. Almost 86% of the children completed the vaccination per the recommended age timeframe. About 88.7% of respondents were married or living together with their male partners, while only 7.3% were never married or living together with their male partners, and the rest were divorced or separated. The majority of respondents (88.7%) had a primary level of education and 81.3% of respondents had a 21-35-year-old age group. About 84.7% of respondents were farmers. The majority (86.7%) were delivered at health facilities, where about 69.3% took less than 30 minutes to reach the nearest health facility.

3.1.2 Mapping of existing nutrition interventions in Njombe and Geita regions

The study showed the implementing agents of nutrition activities by detailing what, who, when, where, how, and to what extent they were implementing nutrition interventions. The assessment was based on interventions aiming to improve maternal nutritional knowledge together with IYCF practices components. This study observed only two multicomponent nutrition interventions (ASRP-Tubadilishe Project and Kizazi kipya Project) with an average coverage area of 66.7% in Njombe. However, only ASRP had a component of maternal knowledge on IYCF practices. Also, findings from this study observed that more than half (52.3%) of respondents had low knowledge related to IYCF practices (Table 5). The result showed that the ASRP role could not be enough to reach its own target of 75% being impacted with acceptable IYCF practices. However, an increase in the stunting rate from 44% to 53.6% (Table 3) of children under five years demonstrates the need for increasing efforts for maternal nutrition knowledge for the betterment of child nutrition status.

Findings from (Table 4) observed four multicomponent nutrition interventions with maternal nutritional knowledge out of five executed in Geita. The projects in the Geita region (ASTUTE, Mtoto Mwerevu, USAID Boresha Afya, and Kizazi Kipya) had an optimistic effect on the community with the reason that they may have contributed to the decrease in proportion of stunting level by 7.1% while in the Njombe region it increased by 9.5% from 2015 to 2020 [6](URT, 2018).

Table 2. Socio-economic and demographic characteristics of infants and mothers

Category	Sub-category	Bukombe district	Njombe district	Total
		n=85	n=65	
Infant gender	Male	40(47.1)	25(38.5)	65(43.3)
	Female	45(52.9)	40(61.5)	85(56.7)
Infant age (months)	Mean (SD)	21.35(13.76)	27.37(13.85)	23.96(14.08)
Infant birth weight (kg)	Mean (SD)	3.24(0.42)	3.17(0.48)	3.21(0.45)
Infant vaccination status	Completed	42(49.4)	44(67.7)	86(57.3)
	Not completed	4(4.7)	1(1.5)	5(3.3)
	< 18 months	39(45.9)	20(30.8)	59(39.3)
Mother's marital status	Married	77(90.6)	56(86.2)	133(88.7)
	Divorced/Separated	4(4.7)	2(3.1)	6(4)
	Never Married	4(4.7)	7(10.8)	11(7.3)
Maternal age group	15-20 years	15(17.6)	7(10.8)	22(14.7)
	21-35 years	68(80.0)	54(83.1)	122(81.3)
	36-49 years	2(2.4)	4(6.2)	6(4.0)
Maternal education	Primary	51(60)	46(70.8)	97(64.7)
	Secondary	17(20)	15(23.1)	32(21.3)
	Technical/Vocational	3(3.5)	0(0.0)	3(2)
Maternal occupation	None	14(16.5)	4(6.2)	18(12)
	Farmer	68(80)	59(90.8)	127(84.7)
	Domestic help	15(17.6)	2(3.1)	17(11.3)
Maternal delivered place	Public servant &Others	2(2.4)	4(6.2)	6(4.0)
	Health facility	67(78.8)	63(96.9)	130(86.7)
	Home delivery	18(21.2)	2(3.1)	20(13.3)
Time to reach the nearest health facility	1-30 minutes	55(64.7)	49(75.4)	104(69.3)
	31-60 minutes	21(24.7)	16(24.6)	37(24.7)
	>61 minutes	9(10.6)	0(0.0)	9(6)

Note: Number in bracket is percentages, Continuous data being presented by Mean (SD) while SD means Standard Deviation

Table 3. Mapping and evaluation of nutrition projects in Njombe district as per 31 January 2020 (qualitative information)

Project Title	Implementing rganization	Covered Area(s)	Project Objectives	Outcome Evaluation done
Accelerating Stunting Reduction Program (ASRP) (Tubadilisha Project) (2013-2020)	Consortium of Doctors with Africa-(CUAMM) and Tanzania Home Economics Association (TAHEA)	The whole part of Njombe region.	To reduce the prevalence of stunting among children under five years from 44% in 2015 (TDHS) to 35% in 2020 (i.e. 3.4% average annual reduction rate), Recent survey stunting rate is 53.6 (2018/2019 TNNS)	<ol style="list-style-type: none"> 1. Increased proportion of pregnant women and mothers/caregivers of children under two years old who practice key pro-nutrition behaviours including IYCF, Health, WASH, and CCD (From 5% in 2014 to 75% in 2020) 2. Increased availability of diverse nutrient-rich foods at the household level in Njombe Regions (Mean household dietary diversity score, from 6 in 2013 to 10 in 2020) from year 1 to year 3. 3. Increased coverage of integrated management of severe acute malnutrition (IMSAM) for less than 5 children and improvement of overall SAM services governance. (Outcome two for Y4&Y5). 4. Strengthening the health system to prevent and respond to violence, abuse, neglect, and exploitation of children in the council
Kizazi Kipya Project	Community Concern of Orphans and Development Association (COCODA) and PACT.	Njombe Council Njombe Council.	Town and District	<ol style="list-style-type: none"> 1. To enable orphans and vulnerable children age 0-18 years together with adolescents aged 18-19 years to access complete HIV-related services and other essentials for improved health, nutrition, education, protection, livelihoods, and psychosocial well-being. <p>No evaluation report available during the period of data collection.</p>

Table 4. Mapping and evaluation of nutrition projects in Bukombe district as per 31 January 2020

Project Title	Implementing Organization	Covered Area(s)	Project Objectives	Outcome Evaluation done
Addressing Stunting in Tanzania Early (ASTUTE). (Mtoto Mwerevu) (2016-2020)	IMA WORLD HEALTH	17 wards in Geita Region	<ol style="list-style-type: none"> To reduce stunting to children with under five age. Consolidation a multi -sectoral reply to nutrition. Ornamental best care practices for infant, young child and maternal nutrition; water, sanitation and hygiene; an early childhood development. Increasing the knowledge of pregnant women, caregivers, household and community decision makers. 	<ol style="list-style-type: none"> From 2016 to 2019 there was decrease about 7.1% level of stunting from 46% to 38.9%.
Mtoto mwerevu July 2018 – Feb 2020	Mwangaza SBCO for Bukombe Dc only.	Geita region	<ol style="list-style-type: none"> Community groups formulation and educate them on nutritional knowledge. Capacity building on maternal, infant and young child nutrition practices; sanitation and hygiene; supplementary foods; child care and growth; and gender issues. 	<ol style="list-style-type: none"> There is a clearly understand of nutritional related knowledge in the community. Only 83% of the project's target was reached by an intervention (75% out of 90%). The intervention was targeted at three community groups in each ward. There was slight participation of male in the project activities, though gender norms and gender discrimination.
USAID Boresha Afya	JHPIEGO	17 wards in Geita such as	<ol style="list-style-type: none"> Maternal, Infant and Young Child Nutrition. Growth Monitoring. Management of acute malnutrition. 	<ol style="list-style-type: none"> For the integration of acute malnutrition, the project trained 34 health workers for implementation. Two hospitals in the region deliver services for the treatment of acute malnutrition. Training of nutrition-related knowledge to health workers and CHW toward facilities and communities' levels.
Kizazi Kipya	NELICO	Geita Dc	<ol style="list-style-type: none"> Nutrition counseling to mother and 	<ol style="list-style-type: none"> *Services delivery through

Project Title	Implementing Organization	Covered Area(s)	Project Objectives	Outcome Evaluation done
Red Cross	RED CROSS Society	Mbogwe Dc	child. 2. Referral and linkage to malnourished children. 1. Not available during surveyed period	community health worker by 00% from baseline to date. 1. Provided support of Ready to use therapeutic food to child with under five of age in the District.

Note: Outcome data for Kizazi Kipya Project was not found from NELICO

Table 5. Maternal nutritional knowledge (MNK) variation between age groups and within Bukombe and Njombe districts

Category	Sub-category	Maternal nutritional knowledge		X ²	p-value
		Poor/ lower	Good/ higher		
Mother's age group	14- 20 years	19 (90.5)	2 (9.5)	7.747	0.005
	21- 49 years	76 (58.9)	53 (41.1)		
District of respondent	Bukombe	61 (71.8)	24 (28.2)	6.005	0.014
	Njombe	34 (52.3)	31 (47.7)		

Note: Number in bracket is percentages

Table 6. Prevalence of stunting, underweighting and wasting in children with under five years of age by sex and age group in Bukombe and Njombe districts

Category	Sub-category	Bukombe district			Njombe district			p-value
		N	n	% [CI 95%]	N	n	% [CI 95%]	
Stunting								
Child sex	male	40	15 ^a	37.5 [22.5-52.5]	25	17 ^a	68.0 [49.7-86.3]	0.048 ^a
	female	45	17	37.8 [23.6-52.0]	40	18	45.0 [29.6-60.4]	
				<i>p-value</i>			<i>p-value</i>	
Child age	0-5.9 months	12	4	33.3 [6.6-60.0]	4	3	75.0 [34.6-115.4]	0.242 ^b
	6-11.9 months	24	13	54.2 [34.3-74.1]	13	6	46.1 [19.0-73.2]	0.772 ^c
	12-23.9 months	44	14	31.8 [18.0-45.6]	38	21	55.2 [39.4-71.0]	
	24-59.9 months	5	1	20.0 [-15.0-55.1]	10	5	50.0 [19.0-81.0]	
Underweighting								
Child sex	male	40	5	12.5 [2.3-22.7]	25	3	12.0 [-0.7-24.7]	0.972 ^a
	female	45	4	8.9 [0.6-17.2]	40	4	10.0 [0.7-19.3]	
				<i>p-value</i>			<i>p-value</i>	
Child age	0-5.9 months	12	0	-	4	0	-	0.189 ^b
	6-11.9 months	24	5	20.8 [4.6-37.0]	13	2	15.4 [-4.2-35.0]	0.531 ^c

Category	Sub-category	Bukombe district			Njombe district			p-value
		N	n	% [CI 95%]	N	n	% [CI 95%]	
Wasting	12-23.9 months	44	4	9.1 [0.6-17.6]	38	5	13.2 [2.5-23.9]	
	24-59.9 months	5	0	-	10	0	-	
Child sex	male	40	6	15.0 [4.9-26.1]	25	1	4.0 [-3.7-11.7]	0.017 ^a
	female	45	4	8.9 [0.6-17.2]	40	0	-	

Note: ^a Chi square test was used for finding significant difference between Bukombe and Njombe districts; ^{b,c} Chi square test was used for finding significant difference between sex and age groups in Bukombe and Njombe districts respectively. P-value ≤ 0.5 was considered as statistical significance

Table 7. Association of maternal nutrition knowledge among mothers' age group, IYCF practices and child nutritional status

Category	Sub category	Maternal nutritional knowledge (MNK)				Total (p-value)
		Bukombe district		Njombe district		
		Poor/Lower	Good/Higher	Poor/Lower	Good/Higher	
Mothers' age group	15-20 years	14(100.0)	0(0.0)	5(71.4)	2(28.6)	0.005
	21-49 years	47(66.2)	24(33.8)	29(50.0)	29(50.0)	
	<i>p-value</i>	0.010		0.284		
Child feeding Practices						
Early initiation of breastfeeding	disqualified	15(28.8)	8(15.4)	9(31.0)	9(31.0)	0.715
	qualified	21(40.4)	8(15.4)	4(13.8)	7(24.1)	
	<i>p-value</i>	0.577		0.474		
Min. dietary diversity	disqualified	45(61.6)	17(23.3)	30(48.4)	27(43.5)	0.655
	qualified	8(11.0)	3(4.1)	3(4.8)	2(3.2)	
	<i>p-value</i>	0.992		0.752		
Child nutritional status						
Stunting	stunted	26(35.6)	6(7.1)	22(33.8)	13(20.0)	0.054
	normal	35(41.2)	18(21.2)	12(18.5)	18(27.7)	
	<i>p-value</i>	0.131		0.066		
Underweight	underweighted	7(8.2)	2(2.4)	6(9.2)	1(1.5)	0.116
	normal	54(63.2)	22(25.9)	28(43.1)	30(46.2)	
	<i>p-value</i>	0.672		0.061		
Wasting	wasted	8(9.4)	2(2.4)	1(1.5)	0(0.0)	0.186
	normal	53(81.5)	22(25.9)	33(50.8)	31(47.7)	
	<i>p-value</i>	0.538		0.336		

Note: Number in bracket is percentages

3.1.3 Maternal nutritional knowledge (MNK) variation between age groups and within Njombe and Bukombe district

The finding from study (Table 5) showed that about 90.5% of adolescent mothers aged 14 to 20 years had lower maternal nutritional knowledge compared to those mothers aged 21 to 49 years who had 59% prevalence rate of poor or lower nutritional knowledge, though their difference was statistically significant ($p=0.005$). The variation of maternal nutritional knowledge within the districts of respondent had statistical meaningful ($p= 0.014$) with prevalence rate of 72% and 52% of mothers with poor or lower nutritional knowledge in Bukombe and Njombe districts respectively.

The variation observed on maternal nutritional knowledge within districts also showed the level and effort made by existing nutritional interventions with the elements of maternal nutritional knowledge together with IYCF practices. Despite the large numbers of nutrition intervention (ASTUTE, Mtoto mwerevu, USAID Boresha Afya etc.) in Bukombe district, mothers with reproductive age still had little knowledge related to child nutrition compared with mothers in Njombe district.

3.1.4 Nutrition status of children aged 0-59 months

The findings for infant and young child nutritional status was indicated from Table 6 where by stunting was significantly different between Bukombe and Njombe districts ($p = 0.048$). In Njombe district, 68% of male infants and young children were more stunted than female (45%). Related studies indicated that the infants and young children lived in rural areas and those who are male were more impacted with stunting [29]. Njombe district has large proportional average of stunted infants between 0-11.9 months of age. The other results presented by age categories report that infant between 0-11.9 months were harmful impacted with stunting than other age groups [8,30]. The proportion of stunting in all male infants and young children in Njombe district was between 49.7% and 86.3% (95% C.I). 12.5% and 15.0% of Male infants and young children in Bukombe district were underweight and wasting, respectively. Bukombe and Njombe districts had statistically significant differences for wasted infants and young children ($p = 0.017$). Also, there was no statistically significant difference in nutrition status for infants and young children between sex and age groups. The findings highlighted a high frequency of infants

and young children with low HAZ measurement in Njombe district; less frequency of WAZ and WHZ measurements in both Bukombe and Njombe districts were nearly correlated to recent national data, whereas 38.9% and 53.6% of infants and young children were stunted in Bukombe and Njombe districts respectively (URT, 2018).

3.1.5 Association of maternal nutrition knowledge on IYCF practices and nutrition status

Table 7 highlighted that mothers within regions of high-rate malnutrition had poor knowledge related to child nutrition. About 85.7% of adolescent mothers had little nutritional knowledge. There was a statistically significant difference between mothers with or without nutritional knowledge and the mother's age group in Bukombe district ($p = 0.010$), in addition ($p = 0.005$) significant difference of maternal nutrition in Bukombe and Njombe districts. The findings showed that maternal nutrition knowledge had no statistically significant difference with IYCF practice indicators ($p = 0.751$ and 0.655) and stunting, underweighting and wasting ($p = 0.058$, 0.116 and 0.186 ; Table 7) in Bukombe and Njombe districts.

3.2 Discussion

This study assesses the role of nutritional interventions scheduled maternal knowledge on IYCF practices within the regions of high rate of undernutrition and identifies the number of projects available, their coverage areas with their outcomes (either process or final outcomes). Mothers' IYCF practices knowledge is the fundamental principle for mothers to improve and comply with acceptable IYCF practices. Improved maternal knowledge of IYCF practices was assumed as the main for reduced if not eliminated child (0-59 months) undernutrition within the regions of high rate of malnutrition.

3.2.1 Socio-economic and demographic characteristics of infants and mothers

The findings were much closed with TDHS-MIS report of 2015/2016 here indicated that majority of children were born with acceptable birth place and weight, also almost more than half of respondents had primary level of education which may also have an impact to maternal nutritional knowledge hence child nutrition status. Based on the results (Table 2), most mothers (86.7%) gave birth at health facilities, and 69.3% of them got there in less than 30 minutes. This means that the study areas had ready-to-use

health services, including services related to nutrition for mothers and babies, which are important for improving the health and nutrition of children.

3.2.2 Mapping of existing nutrition interventions in Bukombe and Njombe districts

In spite of the effort made for improving nutrition outcomes there was little sensitive focus on civilizing maternal knowledge on infant and young child feeding practices as essential elements needed to pick up nutrition status. Other studies indicated that there is a need of amplification together with improves of nutrition specific and highly responsive nutrition action such as maternal knowledge and IYCF practices so as to sustain the outcome [31,32,33,34] (Turner et al., 2018; CW and UNICEF 2015). This was due to fact that mother with poor nutritional knowledge had underprivileged implementation of infant and young child feeding practices led to deprived child nutrition status [1,10,35,36,37]. Results from (Table 2, 3 and 4) also recognized the contribution of clinical services (PNC and ANC) which offer the essential opportunities of maternal nutrition knowledge transformation for better achievement of child nutrition status. However, the study observed little effort done within Njombe regions with small numbers of nutritional related interventions together with their coverage areas compared to Geita region.

3.2.3 Maternal nutritional knowledge (MNK) variation between age groups and within Bukombe and Njombe districts

The study revealed that mothers of reproductive age within regions of high prevalence rates of stunting still had little knowledge related to IYCF practices and child nutrition status. Also, there was little experience and skills in child nutrition-related knowledge for adolescent mothers compared to adults. The variation between age groups indicates the perception of the need to expand health services and nutrition-related knowledge at the earliest age. Adolescents are among the vulnerable groups that deserve special treatment in obtaining sexual and reproductive health services (SRHS), particularly knowledge and services related to maternal and infant nutrition [38]. Lower sexual and reproductive health services (SRHS) together with poor maternal nutritional knowledge of adolescent girls are among the determinants of unwanted pregnancy, which leads to the high risk

of inaccuracy in caring for the infant's health within the first 1000 days of life. Other studies have argued that mothers with limited access to SRHS have poor knowledge of IYCF practices, hence unacceptable child nutritional status [39,40].

3.2.4 Nutrition status of children aged 0-59 months

The study observed a significant reduction in stunting prevalence among under-5 children in Bukombe resulting from recommended IYCF practices due to nutrition interventions that rely on an improvement in maternal nutritional knowledge (Table 5 and 6). Table 3 shows some positive outcomes, namely an increase of 70% in practice key pro-nutrition behaviours including IYCF, in Njombe district. Despite of this, Table 6 shows a higher prevalence, although not statistically significant, of stunting children in Njombe district, compared with Bukombe district. When observing the association of maternal nutrition knowledge with IYCF practices indicators and child nutrition status, by districts (Table 7). The findings showed that maternal nutrition knowledge had no statistically significant difference with IYCF practices indicators and anthropometric indices (Table 7) in Bukombe and Njombe districts. While, other factors such as seasonal variation in food accessibility; social-economic factors, and little availability of animal-source foods may have contributed to poor nutrition status, other researchers revealed that optimal child nutrition status is largely dependent on how well mothers/caregivers have knowledge of IYCF practices [7,33,41,42,43] (Additionally, maternal nutritional knowledge had a higher average among major ethnic groups in Njombe district than in Bukombe district but stunting within major ethnic groups in Njombe was higher than the child stunted rate within major ethnic groups in Geita. Young children within ethnic groups in the Bukombe district had a greater chance to acquire a minimum acceptable diet as per WHO recommendations compared to those in the Njombe district. While other factors contributing to suboptimal feeding practices, further investigation on ethnicity and child nutrition are needed.

3.2.5 Association of maternal nutrition knowledge on IYCF practices and nutrition status

This study showed that mothers who didn't know much about nutrition were more likely to engage

in unacceptable IYCF practises and give their children bad nutrition. Mothers may not always know how important variety and balance are in a child's diet or how much and what kinds of food kids need to grow and be healthy. Since a mother's main job for the first five years of her child's life is to take care of him or her, it's important to know what to eat, especially when there aren't many resources for food and health. The findings seem similar to studies with the same group [10,11,44-50].

4. CONCLUSION AND RECOMMENDATIONS

4.1 Conclusion

The study estimated the importance of nutrition interventions on maternal knowledge of IYCF practices with an impact on child nutritional status within regions of high malnutrition rates in Tanzania. There were some initiatives made to improve child nutritional status within the most vulnerable regions, but the efforts were not good enough. Analysis from this study observed little effort guided by a small number of multicomponent nutrition interventions to attain nutrition status through maternal nutritional knowledge in Njombe. Still, there was poor maternal nutrition knowledge, but their variations reflected the numbers and readiness of nutrition projects to achieve IYCF practices together with acceptable child nutritional status. From the study view point, areas with less efforts had a worse increase in stunting rate from 44% to 54%, while areas with greater efforts and readiness had a significant decrease in stunting rate by 7% for the period of 2015-2020. Also, infants aged 0–11.9 months were more stunted than other age groups, and this emphasizes the need to observe maternal health and nutritional status while pregnant.

4.2 Recommendations

As per findings from this study and conclusion the followings are suggestions for improving maternal nutritional knowledge on IYCF practices for healthier child nutritional status.

- (i) Knowledge related to IYCF practices should expand and strengthen intentionally that the community, particularly women need a better understanding of what, when, how, and why to feed themselves and their infants to comply with WHO guidelines. This can be taken into consideration by all stakeholders such as

government authorities, and non-government organizations.

- (ii) The study observed that the prevalence of stunted children was very high for mothers with poor nutrition knowledge and had the worst effect on adolescent girls (15–20 years). This therefore speculates that the needs of governments and non-government organizations to initiate and strengthen the programs of nutrition knowledge to the earliest stages of learning may be at primary or secondary levels of education. There is probably a chance that some infants were born stunted, and this also may direct future program interventions to focus mainly on maternal health and child nutritional status.
- (iii) Mutual efforts from public entities and donors are needed to address, expand the coverage, and strengthen multi-component nutrition initiatives to improve their efficiency, effectiveness, impact, and sustainability on maternal nutrition knowledge.

CONSENT

As per international standard or university standard, respondents' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

Ethical clearance was obtained from Research Ethics Committees of Sokoine University of Agriculture, Morogoro Tanzania. Additionally, Njombe and Geita regions with District Administration and local leaders gave the permission to conduct the study in their respective areas. The permission to use village health centres for data collection was obtained from District Medical Officers (DMO) of the respective districts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. WHO/UNICEF/ WB. Levels and Trends in Child Malnutrition. World Bank Group, Washington DC. 2019:16.
2. URT. Tanzania National Nutrition Survey 2018. United National International

- Children's Emergency Fund, New York. 2019:144.
3. Aguayo VM, Kang Y, Dzed L, Campbell RK, Joshi V, Waid J, West KP. Infant and young child feeding practices and nutritional status in Bhutan. *Maternal and Child Nutrition*. 2018;14: 1–6.
 4. Menon P, Bamezai A, Subandoro A, Ayoya MA, Aguayo V. Age-appropriate infant and young child feeding practices are associated with child nutrition in India: Insights from nationally representative data. *Maternal and Child Nutrition*. 2015; 11(1):73–87.
 5. Mbogori T, Murimi M. Effects of a nutrition education intervention on maternal nutrition knowledge, dietary intake and nutritional status among food insecure households in Kenya. *International Journal of Community Medicine And Public Health*. 2019;6(5):1831.
 6. UNICEF. *Approach to Scaling up Nutrition for Mothers and Their Children*. United National International Children's Emergency Fund, New York. 2015:21.
 7. Abubakar A, Uriyo J, Msuya SE, Swai M, Stray-Pedersen B. Prevalence and risk factors for poor nutritional status among children in the Kilimanjaro Region of Tanzania. *International Journal of Environmental Research and Public Health*. 2012;9(10):3506 –3518.
 8. Muhimbula H, Kinabo J, O'Sullivan A. Determinants of infant nutrition status in rural farming households before and after harvest. *Maternal and Child Nutrition* 2019; 15(3):1–13.
 9. Fadare O, Amare M, Mavrotas G, Akerele D, Ogunniyi A. Mother's nutrition-related knowledge and child nutrition outcomes: Empirical evidence from Nigeria. *PLoS One*. 2019;14(2):1–17.
 10. El-nmer F, Salama AA, Elhawary D. Nutritional knowledge, attitude, and practice of parents and its impact on growth of their children. *Menoufia Medical Journal*. 2014;27:612–616.
 11. URT. Tanzania 2015-16 Demographic Health Survey and Malaria Indicator Survey 2015-2016. Ministry of Health, Community Development, Gender, Ederly and Children Dar es Salaam, Ministry of Health Zanzibar, National Bureau of Statistics Dar es Salaam, Office of Chief Government Statistician Zanzibar, ICF Rockville, Maryland USA. 2015: 260.
 12. Jemide JO, Ene-obong HN, Edet E, Udoh EE. Association of maternal nutrition knowledge and child feeding practices with nutritional status of children in Calabar South local government area, Cross River State, Nigeria. *International Journal of Home Science*. 2016;2(1):293–298.
 13. URT. Basic Demographic and Socio-Economic Profile. National Bureau Statistics, Dar es Salaam, Tanzania. 2012a:245.
 14. URT. Nutrition Policy Mapping for Tanzania 2012 Acknowledgement. Save the Children, Dar es Salaam. 2012b:56.
 15. Fanzo J, Hawkes C, Unomkesmalee E, Afshin A, Allemandi L, Assey O. *Global Nutrition Report: Shining a Light to Spur Action on Nutrition*. World Bank, Washington DC. 2018;165.
 16. Doudou MH, Ouedraogo O, Ouaro B, Bidault N, Reinhardt K. Mapping nutrition interventions, a key analytical tool for informing the multisectoral planning process: example from Burkina Faso. *Food and Nutrition Bulletin*. 2018;;39(3):449-464.
 17. URT. National Sample Census of Agriculture 2002/2003. Regional Report: Iringa Region. National Bureau of Statistics, Ministry of agriculture and Food Security, Ministry of Water and Livestock Development, Ministry of Cooperatives and Marketing, Presidents Office, Regional Administration and Local Government, Ministry of Finance and Economic. Zanzibar. 2007:344.
 18. URT. Njombe District Council Socio-Economic Profile. National Bureau of Statistics, Dar es Salaam. 2016b:155.
 19. Bell A, Jones K. Explaining fixed effects: random effects modeling of time-series. *Cross-Sectional and Panel Data*. 2015: 3(1):133–153.
 20. Simmons JP, Nelson LD, Simonsohn U. *Life After P-Hacking*. New Orleans, Los Angles. USA. 2013:210.
 21. Beleites C, Neugebauer U, Bocklitz T, Kra C, Popp J. Sample size planning for classification models. *Chemometrics in Analytical Chemistry*. 2013:760:25–33.
 22. Mundfrom DJ, Shaw DG, Ke TL. Minimum sample size recommendations for conducting factor analyses. *International journal of testing*. 2005 Jun 1;5(2):159-68.
 23. UNICEF. *Programming Guide Infant and Young Child Feeding*. Nutrition Section. United National International Children's Emergency Fund, New York. 2011:173pp.

24. Mariás Y, Glasauer P. Guidelines for assessing nutrition-related knowledge, attitudes and practices. Food and Agriculture Organization of the United Nations (FAO); 2014c.
25. Cogill B. Anthropometric Indicators Measurement Guide. Washington, DC: Food and Nutrition Technical Assistance (FANTA) Project, FHI 360; 2003.
26. De Onis M, Onyango AW, Borghi E, Garza C, Yang H, WHO Multicentre Growth Reference Study Group. Comparison of the World Health Organization (WHO) Child Growth Standards and the National Center for Health Statistics/WHO international growth reference: implications for child health programmes. *Public health nutrition*. 2006 Oct;9(7):942-7.
27. FANRPAN. Agriculture to Nutrition: Evaluation of Integrated Agriculture and Nutrition-Sensitive Interventions for the African Chicken Genetic Gains Programme in Tanzania - Baseline Data. Food and agriculture Organization, Rome, Italy. 2017:119.
28. Gertler PJ, Martinez S, Premand P, Rawlings LB, Vermeersch CM. Impact Evaluation in Practice. World Bank, Washington DC. 2016:267.
29. Keino S, Plasqui G, Etyang G, Van Den Borne B. Determinants of stunting and overweight among young children and adolescents in sub-Saharan Africa. *Food and Nutrition Bulletin*. 2014;35(2):167-178.
30. Ali Z, Saaka M, Adams A, Kamwininaang SK, Abizari A. The effect of maternal and child factors on stunting, wasting and underweight among preschool children in Northern Ghana. *BioMed Central Nutrition* 2017;3:1-13.
31. Deborah A, Kavita S, Elisabeth S, Lesley O, Tara K, Joyceline K, Malisa Y. Reducing Malnutrition in Tanzania: Estimates to support nutrition advocacy: Reducing malnutrition in Tanzania: estimates to support nutrition advocacy Tanzania; 2014. Available:<https://www.fantaproject.org/sites/default/files/resources/Tanzania.Profiles.pdf>] site visited on 10/5/2021.
32. IRS. Overcoming the Challenges of Under-Nutrition in Tanzania Through 2021: Reflecting on Trends, Questions, and Future Scenarios for Urgent Nutrition Action; 2017. Available:<https://www.actionagainsthunger.org.uk/publication/overcoming-challenges-undernutrition-tanzania-through>] -2021 site visited on 9/3/2021.
33. Msuya J. Nutrition improvement projects in Tanzania: Appropriate choice of institutions matters. *Electronic Journal*. 2019;12:1-40.
34. World Bank. Investment Climate Reforms: An Independent Evaluation of World Bank Group Support to Reforms of Business Regulations. The World Bank; 2015 Nov 2.
35. Acheampong I, Haldeman, L. Are nutrition knowledge, attitudes, and beliefs associated with obesity among low-income hispanic and African American women caretakers?; 2013. Available:<https://doi.org/10.1155/2013/1239012013> site visited on 10/5/2021.
36. Asoba GN, Ule I, Sumbele N, Anchang-kimbi JK, Metuge S, Teh RN. Influence of infant feeding practices on the occurrence of malnutrition, malaria and anaemia in children 5 years in the Mount Cameroon area: A cross sectional study. *Ploce One* 2019:1-17.
37. Mosimah CI. Impact of the Global Strategy for Infant and Young Child Feeding on Nutrition and Child Survival Indicators as Illustrated by the World Breastfeeding Trends Initiative Tool; 2015 Available:<http://corescholar.libraries.wright.edu/mph> site visited on 10/5/2021.
38. Chilinda I, Hourahane G, Pindani M, Chitsulo C, Maluwa A. Attitude of health care providers towards adolescent sexual and reproductive health services in developing countries: a systematic review. *Health*. 2014 Jul 28;2014.
39. Sedgh G, Finer LB, Bankole A, Eilers MA, Singh S. Adolescent pregnancy, birth, and abortion rates across countries: Levels and recent trends. *Journal of Adolescent Health*. 2015;56(2):223-230.
40. Niyeha D, Mpembeni R, Charwe D, Epimark S, Malima K, Aveika A, Niyeha D. Explaining progress Towards Millennium Development Goal 4 for child survival in Tanzania. *Journal of Global Health*. 2018;8:1-12
41. Kaminski J, Christiaensen L, Gilbert CL. Seasonality in local food markets and consumption: Evidence from Tanzania. *Oxford Economic Papers*. 2016;68(3):736-757.

42. Mistry SK, Hossain MB, Arora A. Maternal nutrition counselling is associated with reduced stunting prevalence and improved feeding practices in early childhood: A post-program comparison study. *Nutrition Journal*. 2019;18(1):1–9.
43. Rakotomanana H, Hildebrand D, Gates GE, Thomas DG, Fawbush F, Stoecker BJ. Maternal knowledge, attitudes, and practices of complementary feeding and child undernutrition in the Vakinankaratra Region of Madagascar: A mixed-methods study. *Current Developments in Nutrition* 2020;4(11):1–11
44. Agize A, Jara D, Dejenu G. Level of knowledge and practice of mothers on minimum dietary diversity practices and associated factors for 6-23-month-old children in Adea Woreda, Oromia; 2017. Available:<https://doi.org/10.1155/2017/7204562>] site visited on 10/5/2021.
45. Saaka M. Relationship between mothers' nutritional knowledge in childcare practices and the growth of children living in impoverished rural communities. *Journal of Health, Population and Nutrition*. 2014; 32(2):237–248.
46. WHO. Global nutrition shining a light to spur action on nutrition; 2018> Available:<https://doi.org/http://dx.doi.org/10.2499/9780896295643>] site visited on 20/6/2021.
47. WHO. Comprehensive implementation plan on maternal, infant and young child nutrition; 2014. Available:www.who.int] site visited on 20/6/2021.
48. Worldwide. Bringing Nutrition Actions to Scale in Iringa, Njombe and Mbeya Regions of Tanzania in-Depth Analysis of the Factors Associated with Stunting. Centre for Research on the Epidemiology of Disasters, Brussels, Belgium. 2015; 51.
49. Yakubu I, Salisu WJ. Determinants of adolescent pregnancy in sub-Saharan Africa: A systematic review. *Reproductive Health*. 2018;15(15): 1–11.
50. Simmons JP, Nelson LD, Simonsohn U. Palse –positive psychology undisclosed flexibility in data collection and analysis allows presentating anything as significant. *Psychological Science*. 2011;22(11):1359 – 1366.

© 2023 Berenge and Muhimbula; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/106762>