



Effect of Foliar Application of Various Concentrations of NAA and GA₃ on Fruiting, Yield and Quality Attributes of Ber cv. Banarasi Karaka, India

**Sheel Priya^a, A. K. Dwivedi^a, V. K. Tripathi^a
and Somendra Verma^{a*}**

^a Department of Fruit Science, C.S Azad University of Agriculture and Technology, Kanpur (Uttar Pradesh), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2023/v42i484329

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110091>

Original Research Article

Received: 15/10/2023

Accepted: 20/12/2023

Published: 24/12/2023

ABSTRACT

To study the effect of foliar application of different concentrations of NAA and GA₃ on fruiting, yield and quality of ber, an investigation was carried out in the Garden, Department of Fruit Science, C. S. Azad University of Agriculture and Technology, Kanpur during the year 2021-22 on ber cv. Banarasi Karaka. Results of the investigation revealed that the significantly main effect among the spray of various concentrations of NAA 30 ppm results, maximum number of fruit set (165.50), minimum fruit drop (85.15%), maximum fruit retention (14.45%), fruit weight (18.32g), yield (44.02 kg/tree), volume (16.17cc), length (4.36cm), diameter (2.56cm), pulp weight (16.12g), total soluble solids (15.48°Brix) and total sugars (10.32%) against control (water) treatment. Spray with various concentrations of GA₃ showed significant results, having maximized number of fruit sets (160.00), minimum fruit drop

*Corresponding author: E-mail: somendraverma56@gmail.com;

(89.16%), maximum fruit retention (10.16%), weight (15.95g), yield (36.78kg/tree), volume (16.17cc), length (4.36cm), diameter (2.56cm), pulp weight (16.12g), total soluble solids (15.48°Brix) and total sugars (10.32%) were recorded with spray of 30ppm GA₃ against control (water spray) treatment. Interaction effect among various concentration of NAA and GA₃ has significantly maximized with fruit retention (16.24%), volume (16.70cc), length (4.47cm), diameter (2.61cm), pulp weight (17.55g) and total sugars (10.59%), while non-significantly results showed highest number of fruit set (167.00), minimum fruit drop (83.76%), maximum fruit weight (19.40g), fruit yield (45.70 kg/tree) and total soluble solids (16.62°Brix) with application of NAA and GA₃ @ 30ppm against control (water spray) treatment. This study revealed that for realizing higher fruiting, fruit yield and quality of ber plants should be spread with GA₃ and NAA @ 30ppm under north Indian plains of Uttar Pradesh.

Keywords: GA₃; NAA; Ber; Fruiting; Yield; Physico-chemical characteristics.

1. INTRODUCTION

The ber (*Ziziphus mauritiana* Lamk.) is a vigorous growing, spreading tree with almost vine-like drooping branches, that belongs to the family Rhamnaceae. This species is evergreen and leaves are densely tomentose on their undersurface. The ber fruits are borne in the axil of leaves on the young growing shoots of the current year. Hence, regular annual pruning is necessary to induce healthy growth which will provide a maximum fruit-bearing area on the tree. With wide commercialization, many physiological problems related to flowers and fruit drop, embryo abortion, poor flowering and fruit setting, abnormal and small size fruits, etc. were observed which caused huge losses to the growers. In order to minimize these problems, over the years many experiments and advancements have been done, and among them, usage of NAA and GA₃ is one of the most adopted hormones and is utilized to improve flowering, fruiting, yield and quality of fruit.

PGRs (Plant Growth Regulators) are involved in a variety of physiological processes, including vegetative propagation, induction of seed lessens, increased fruit set, prevention of pre-harvest fruit drop, blooming regulation, fruit size inhibition, and flower and fruit thinning. NAA (Auxin) improved the quality of numerous fruits by increasing fruit sets, decreasing fruit drops, and increasing fruit sets. Auxin concentration in the plant is increased to prevent abscission. Fruit drop is also prevented by NAA (Auxin), which strengthens the pedicle. Fruit drop is prevented by a high auxin level in the abscission zone.

Gibberellins have mostly been used to manipulate a variety of physiological events and are economically employed to increase the quality of fruit in crops such as berries, grapes, citrus,

cherries, and apples, etc. In fruit crops, three physiological phenomena have been highlighted: rachis cell elongation, blossom thinning, and berry growth. Apart from that GA₃ is the most effective in breaking dormancy and causing rapid germination of seeds. Gibberellin influences bolting by stimulating cell division and cell elongation in the sub-apical meristems. Gibberellins have been found very reliable in producing parthenocarpy. Keeping these in view, the present experiment was carried out in the north Indian plains of Uttar Pradesh.

2. MATERIALS AND METHODS

The experiment was carried out in the Garden, Department of Fruit Science, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur (U.P.) during 2021-2022. The experiment comprised 16 treatments combined with four concentrations each NAA of (0, 10, 20 and 30 ppm) and GA₃ (0, 10, 20 and 30 ppm) under a completely randomized block design (factorial) with three replications. The experimental orchard consisted of 48-year-old well-established, healthy and uniform trees of ber cv. 'Banarasi Karaka' spaced at 10×10 meters distance. Foliar applications of the treatments are uniformly sprayed all over the tagged branch in the morning hours with the help of a fine nozzle foot sprayer. The observations were recorded from each treatment of all three replications viz., initial fruit sets, fruit drop (%), retention (%), length (cm), diameter (cm), volume (cc), weight (g), yield (kg/tree), pulp weight (g), total soluble solids (°Brix) and total sugars (%). Physical parameters were analyzed using standard methodology and chemical parameters using methods as suggested in AOAC [1]. Statically data was analysed according to the method suggested by Panse and Sukhantme [2].

3. RESULTS AND DISCUSSION

3.1 Fruiting Attributes

Application of NAA and GA₃ significantly influenced fruiting attributes viz., Initial fruit set, drop (%), and retention (%) in ber (Table 1). The main effect among various concentrations of NAA maximized initial fruit set (165.50) was recorded when the plant was treated with 30 ppm NAA (N₃) closely followed by 160.50 with a spray of 20 ppm NAA (N₂), whereas, a minimum fruit set (151.75) was found in plants kept under control (N₀) treatment. The enhancement of the initial fruit set was indicated in a range from 151.75 to 165.50 and the improvement of the fruit set was increased by 0.09 % and 0.05 % due to sprays of NAA @ 30 ppm. These results corroborate the

finding of Tiwari *et al.*, [3] in aonla, Badal and Tripathi [4] in guava. The initial fruit set was significantly influenced by different concentrations of gibberellic acid. Foliar feeding of 30ppm of GA₃ recorded maximum initial fruit set (160.00) followed by 159.25 in plants treated with 20ppm of GA₃ (G₂), while minimum fruit set (156.75) was recorded in plants kept under control (G₀) treatment. It probably might be due to providing the right concentration of GA₃ during investigation causing enhancement of vegetative growth of the plants and GA₃ hastening the production of more photosynthesis towards the fruit-bearing area which contributed to increased initial fruit set in plants. These findings are collaborated with the reports of Tripathi and Shukla in strawberry [5] Anushi *et al.*, [6] in mango.

Table 1. Effect of NAA and GA₃ and their interaction on fruit drop, retention and growth attributes of Ber

Treatments	Initial fruit set	Fruit drop (%)	Fruit retention (%)	Fruit weight (g)	Fruit volume (cc)
N ₀	151.75	94.16	5.84	11.95	11.92
N ₁	156.25	92.17	7.83	13.97	13.25
N ₂	160.50	89.70	10.30	16.00	14.72
N ₃	165.50	85.55	14.45	18.32	16.17
S.E(m)±	0.05	0.28	0.23	0.28	0.08
CD at 5%	1.45	0.83	0.67	0.83	0.24
G ₀	156.75	91.35	8.65	14.30	13.45
G ₁	158.00	90.87	9.13	14.72	13.85
G ₂	159.25	90.20	9.80	15.27	14.22
G ₃	160.00	89.16	10.84	15.95	14.55
S.E(m)±	0.50	0.28	0.23	0.28	0.08
CD at 5%	1.45	0.83	0.67	0.83	0.24
N ₀ G ₀	150	94.88	5.12	11.10	11.40
N ₀ G ₁	151	94.52	5.48	11.70	11.70
N ₀ G ₂	153	93.97	6.03	12.20	12.20
N ₀ G ₃	153	93.28	6.72	12.80	12.40
N ₁ G ₀	154	92.82	7.18	13.50	12.60
N ₁ G ₁	156	92.38	7.62	13.60	13.10
N ₁ G ₂	157	92.01	7.99	14.10	13.50
N ₁ G ₃	158	91.48	8.52	14.70	13.80
N ₂ G ₀	159	90.94	9.06	15.20	14.20
N ₂ G ₁	160	90.08	9.92	15.80	14.50
N ₂ G ₂	161	89.66	10.34	16.10	14.90
N ₂ G ₃	162	88.15	11.85	16.90	15.30
N ₃ G ₀	164	86.77	13.23	17.40	15.60
N ₃ G ₁	165.1	86.50	13.50	17.80	16.10
N ₃ G ₂	166	85.19	14.81	18.70	16.30
N ₃ G ₃	167	83.76	16.24	19.40	16.70
S.E(m)±	1.01	0.57	0.46	0.57	0.17
CD at 5%	NS	NS	1.34	NS	0.48

Table 2. Effect of NAA and GA₃ and their interaction on yield and quality attributes of Ber

Treatments	Fruit length (cm)	Fruit diameter (cm)	Pulp weight (g)	T.S.S. (°Brix)	Total sugars (%)
N ₀	3.17	2.15	7.04	12.39	8.39
N ₁	3.53	2.35	10.00	13.42	8.96
N ₂	4.00	2.56	12.90	14.42	9.62
N ₃	4.36	2.56	16.12	15.48	10.32
S.E(m)±	0.05	0.05	0.12	0.18	0.05
CD at 5%	0.14	0.15	0.36	0.54	0.15
G ₀	3.62	2.35	10.39	13.59	9.06
G ₁	3.72	2.40	11.08	13.74	9.24
G ₂	3.82	2.43	11.85	13.93	9.43
G ₃	3.90	2.43	12.73	14.45	9.56
S.E(m)±	0.05	0.05	0.12	0.18	0.05
CD at 5%	0.14	0.15	0.36	0.54	0.15
N ₀ G ₀	3.07	2.11	5.75	12.15	8.11
N ₀ G ₁	3.13	2.13	6.70	12.24	8.32
N ₀ G ₂	3.21	2.16	7.49	12.38	8.52
N ₀ G ₃	3.28	2.19	8.24	12.80	8.62
N ₁ G ₀	3.36	2.22	9.17	13.12	8.75
N ₁ G ₁	3.48	2.36	9.55	13.28	8.90
N ₁ G ₂	3.60	2.39	10.24	13.57	9.04
N ₁ G ₃	3.71	2.42	11.04	13.72	9.18
N ₂ G ₀	3.82	2.56	11.79	14.17	9.36
N ₂ G ₁	3.94	2.58	12.59	14.31	9.51
N ₂ G ₂	4.07	2.62	13.11	14.54	9.75
N ₂ G ₃	4.17	2.49	14.12	14.68	9.87
N ₃ G ₀	4.26	2.52	14.86	14.93	10.03
N ₃ G ₁	4.34	2.55	15.48	15.14	10.25
N ₃ G ₂	4.40	2.58	16.59	15.24	10.42
N ₃ G ₃	4.47	2.61	17.55	16.62	10.59
S.E(m)±	0.10	0.10	0.25	0.37	0.10
CD at 5%	0.28	0.30	0.72	NS	0.30

An interaction effect among various concentrations of NAA and GA₃ was found non-significant. Maximum fruit set (167) was observed with a spray of NAA 30ppm and GA₃ 30ppm (N₃G₃ treatment) followed by 166 with 20ppm NAA with 30ppm GA₃ (N₂G₂). The minimum initial fruit set (150) was observed under control (N₀ G₀) treatment. These findings are collaborated with the reports of Bankar and Prasad [7] in ber.

Foliar application of NAA and GA₃ gradually minimized fruit drops and increased fruit retention as compared to control. Application of NAA @ 30ppm significantly reduced fruit drop (85.55%) and increased fruit retention (14.45%) followed by NAA @ 20 ppm noting 89.70% of fruit drop and 10.30% fruit retention. Significantly maximum fruit drop (94.16%) with minimum fruit retention (5.84%) was recorded in plants kept under control (N₀). Reduction in fruit drop was ascertained due to sprays of NAA treatments and decreased by 9.12 and 4.71% over control (N₀).

When the auxin content of the fruits becomes low, fruit drop occurs due to the formation of an abscission layer and retention of fruit are ultimately decreased. The exogenous application of NAA might have increased the concentration of auxin in plants which possibly induced to reduction of fruit drop and fruit retention. These findings are in accordance with the reports of Kumar and Tripathi [8] in strawberry, Lal *et al.*, [9] in kinnow, Tripathi and Viveka Nand [10] in aonla, Badal and Tripathi [4] in guava, Tiwari *et al.*, [3] in aonla.

Gibberellic acid also positively and consistently influenced fruit retention and drop in ber and plants treated with 30ppm GA₃ significantly minimized (89.16%) fruit drop with increased retention (10.84%) and closely followed by GA₃ 20 ppm (90.20% and 9.98% respectively). Significantly maximum fruit drop (91.35%) and minimum retention (8.65%) was obtained in plants kept under control (G₀). As a result, fruit

drop was reduced by 2.38 and 1.24% less than control (G_0) with increased fruit retention as reported by 13.31 to 25.35%. Application of GA_3 may have enhanced auxin production, which has prevented fruit drop in the current study. The increase in fruit retention and reduced fruit drop might be due to the effectiveness of different chemicals as well as GA_3 on the metabolic activity of the plant and improved source-sink relationship which favorably influenced the metabolic status resulting in better check of fruit drop and enhancing retention of the more number of fruits on the plants. These findings are supported by Bhadauria *et al.*, [11] in aonla and Anushi *et al.*, [6] in mango, and Yadav and Chaturvedi [12] in ber. Interaction between NAA and GA_3 was found to be significant and the combined treatment of $N_3 \times G_3$ expressed a significantly minimum 83.76 % fruit drop and maximum fruit retention (16.24%) followed by $N_3 G_2$ (85.19% and 14.81%, respectively). Significantly maximum noted under controlled (N_0G_0) recording 94.88 % fruit drop and 5.12% of fruit retention. These findings are found to be in accordance with reports of Chaurasiya *et al.*, [13] in ber.

3.2 Physical Attributes

The physical attributes *viz.*, fruit length (cm), diameter (cm) and volume (cc) of ber was significantly influenced by NAA treatment and plant treated with 30ppm of NAA proved most effective and maximized the fruit length (4.36cm), diameter (2.56cm) and fruit volume (16.17cc) during experimental period, whereas minimum fruit length (3.17cm), diameter (2.15cm), fruit volume (11.92cc) were recorded in treatment produced from plants kept under control (N_0). The enhancement range of fluctuated fruit length from 3.17 to 4.36cm, diameter from 2.15 to 2.56cm and volume from 11.92 to 16.17cc, respectively were recorded over the control. The superiority in fruit length, fruit diameter and fruit volume were increased due to NAA treatment. It could be attributed to its role in cell division, cell elongation and decreased intracellular space in monocarpic cells, all of which could have improved plant health and resulted in healthier and larger fruit. These findings are in line with reports of Tripathi *et al.*, [14] in mango, Badal and Tripathi [15] in guava and Tiwari *et al.*, in aonla, Tripathi and Viveka Nand in aonla [10].

Gibberellic acid also affected positively in terms of fruit length, diameter and volume and treatment with 30ppm of GA_3 (G_3) significantly

enhanced fruit length (3.90cm), diameter (2.43cm) and volume (14.55cc). Fruits produced from untreated plants (G_0) showed minimum values for these parameters *i.e.*, 3.62cm, 2.35cm and 13.45cc, respectively. The improvement range varies from 3.62 to 3.90cm for fruit length, 2.35 to 2.43cm for fruit diameter and 13.45 to 14.55cc for volume over the control during the respective year of investigation. The increase in fruit size caused by GA_3 treatment could be attributed to a considerable increase in cell division and cell elongation. These discoveries are also linked to active photosynthesis in the plant, with photosynthesis being translocated to the fruits, presumably resulting in larger fruits. Shukla *et al.*, [16] Tripathi *et al.*, [17] in aonla, Dubey *et al.*, [18] Tripathi and Shukla, [19] in strawberry. The interaction effect of NAA \times GA_3 was found non-significant with respect to fruit length *i.e.*, at 30ppm of NAA \times GA_3 ($N_3 \times G_3$) produced maximum fruit length (4.47cm) as compared to control ($N_0 G_0$) in which minimum fruit length (3.07cm) was observed, whereas, fruit diameter and volume were have significant positive effect, showing maximum fruit diameter (2.61cm) and volume (16.70cc) as compared to control ($N_0 G_0$) which showing the minimum value of (2.11cm and 11.40cc, respectively). These findings are consistent with Tripathi *et al.*, [14] in mango.

Application of NAA and GA_3 significantly decreases the length of stone and diameter of stone in ber. Plants treated with 30ppm NAA (N_3) significantly minimized the length of stone and diameter of stone (1.53 and 0.84cm, respectively) as compared to control (N_0) where these were found maximum (2.63 and 1.13cm, respectively). These findings are in agreement with the reports of Singh *et al.*, [20] in ber and Rathod *et al.*, [21] in aonla.

Gibberellin consistently and positively minimizes stone length and stone diameter in ber. Its 30ppm GA_3 (G_3) concentration proved most effective (1.89 and 0.94cm, respectively) as compared to the control (G_0) which proved maximum stone length and diameter (2.18 and 1.13cm, respectively) during the present investigation. These results may have been associated with the active performance of photosynthesis in the plant and they were translocated to the stones which caused to decrease in stone size. These findings are in line with reports of Tripathi *et al.*, [17] in aonla. The interactive effect of NAA associated with GA_3 did not differ significantly but further improvement

was observed over mean values and combined treatment of $N_3 \times G_3$ recorded minimum stone length (1.42cm), whereas maximum stone length (2.79cm) was recorded under control (N_0G_0). In respect of stone diameter was found positively significant and plants treated with 30ppm @ NAA and 30ppm of GA_3 (N_3G_3) significantly minimized diameter of stone (0.80cm). The maximum diameter of stone (1.17cm) was recorded under control (N_0G_0).

3.3 Yield Attributes

Foliar feeding of NAA and GA_3 significantly enhanced fruit weight, pulp weight and fruit yield of ber {Fig. 1 (a) and (b)}. The fruit weight (18.32g), pulp weight (16.12g) and fruit yield (44.02 kg/tree) was enhanced significantly when the plant was treated with 30ppm NAA as compared to the rest of all other treatments, whereas, the poorest values of fruit weight (11.95g), pulp weight (7.04g) and fruit yield (22.82 kg/tree) were found in plants kept under control (N_0). In the current study, the growth regulator NAA may have boosted the synthesis of additional photosynthates and their translocation to the fruits, resulting in higher fruit and pulp weight which ultimately enhanced the yield. These results are in agreement with the respects of Singh et al., [22] in mango, Tiwari et

al., [3] in aonla, Kumar and Tripathi [8] in strawberry, Tripathi and Viveka Nand [10] in aonla.

All the yield attributory parameters i.e., fruit weight, pulp weight and fruit yield were improved with different doses of gibberellic acid and 30ppm of GA_3 emphatically improved fruit weight (15.95g), pulp weight (12.73g) and fruit yield (36.78 kg/tree) as compared to control (G_0) which show minimum fruit weight (14.30g), pulp weight (10.39g) and fruit yield (31.75 kg/tree). These findings are in support with the reports of Tripathi et al., [17] in aonla, Dubey et al., [18] in strawberry.

The interaction effect between NAA x GA_3 was recorded as non-significant in terms of fruit weight, pulp weight and fruit yield and 30ppm of NAA and 30ppm of GA_3 ($N_3 G_3$) treated plants produced maximum fruit weight (19.40g), pulp weight (17.55g) and fruit yield (45.70 kg/tree) closely followed with treatment N_3G_2 which expressed fruit weight of 18.70g, pulp weight of 16.59g and fruit yield of 44.49 kg/tree. Significantly minimum fruit weight (11.10 g), pulp weight (5.75g) and fruit yield (20.78 kg/tree) were recorded under control (N_0G_0). These findings are in line with reports of Kale et al., [23] in ber.




Fig. 1(a). Effect of NAA and gibberellic acid on yield (kg/tree)




Fig. 1(b). Effect of NAA, gibberellic acid and their interaction on yield (kg/tree)

3.4 Chemical Quality Parameters

NAA and GA₃ significantly enhanced the chemical quality of fruits *i.e.*, TSS and total sugars when plants were treated with different concentrations of these hormones. Significantly maximum TSS (15.48⁰Brix) and total sugar (10.32%) was recorded under 30ppm of NAA (N₃) treated plants followed by treatment of 20ppm NAA (N₂) which exhibited TSS of 14.43⁰Brix and total sugars of 9.62%. The plants kept under the control exhibited a minimum TSS (12.39⁰Brix) and total sugars (8.39%). The application of the growth regulators (NAA) may have produced a redirection of more solid metabolites towards developing fruits, boosting amylase activity, and therefore increasing total soluble solid content by converting starch into simple sugar. Kumar and Tripathi [8] in strawberry, Badal and Tripathi [15] in guava.

Different concentrations of GA₃ also improve TSS and total sugars content in fruits and its 30ppm dose of GA₃ (G₃) was found most effective in respect of TSS and total sugars, which exhibited a maximum value of TSS (14.45⁰Brix) and total sugars (9.56%) followed by 20ppm of GA₃ (G₂), which exhibited values of 13.93⁰Brix and 9.43%, respectively. Significantly minimum TSS (13.59⁰Brix) and total sugars (9.06%) were recorded under control (G₀). GA₃ may cause an increase in the inactivation of the amylase enzyme, which is responsible for the conversion of starch to sugar and is linked to an increase in TSS and total sugars content of fruit.

These findings collaborated with the findings of Verma *et al.*, [24] in strawberry, Shukla *et al.*, [16] Tripathi *et al.*, [17] in aonla and Anushi *et al.*, [6] in mango.

The interaction effect between NAA x GA₃ produced significant effect on TSS and total sugar content and 30ppm of NAA combined with 30ppm GA₃ (N₃ x G₃) was found most effective exhibited maximum TSS (16.62⁰Brix) and total sugars (10.59%) followed by N₃G₂ recorded TSS of (15.24⁰Brix) and total sugars content *i.e.*, 10.42% over its control (N₀G₀), which showed minimum values 12.15⁰Brix and 8.11% respectively. These findings are in line with the reports of Ram *et al.*, [25] in ber.

4. CONCLUSION

This study revealed that for realizing higher fruiting, fruit yield and quality of ber plants should be spread with GA₃ and NAA @30ppm under north Indian plains of Uttar Pradesh, India.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. AOAC. Official Methods of Analysis. Association of Official Analytical Chemists, Washington DC; 1980.

2. Panse VG, Sukhatme PV. Statistical methods for agricultural workers, 4thEdn., New Delhi; 1985.
3. Tiwari P, Tripathi VK, Singh A. Effect of foliar application of plant bio-regulators and micronutrients on fruit retention, yield and quality attributes of aonla. Progressive Research-An International Journal, 12 (Special-IV). 2017;2565-2568.
4. Badal DS, Tripathi VK. Influence of foliar feeding of NAA and Boron on growth, flowering, fruiting, and yield of winter season guava (*Psidium guajava* L.) cv. L-49. Biological Forum-An International Journal. 2021a;13(3):387-391.
5. Tripathi VK, Shukla PK. Influence of plant bio-regulators and micronutrients on flowering and yield of strawberry cv. Chandler. Annals of Horticulture. 2008; 1(1):45-48.
6. Anushi, Tripathi VK, Awasthi V, Yashasvi GN. Impact of pre-harvest application of plant bio-regulators and micronutrient on fruit retention, yield and quality of mango (*Mangifera Indica* L.). Frontiers in Crop Improvement, 9 (Special Issue-III). 2021; 1026-1030.
7. Bankar GS, Prasad RN. Effect of gibberellic acid and NAA on fruit set and quality of fruit in ber cv. Gola. Progressive Horticulture. 1990;22(1-4):60-62.
8. Kumar R, Tripathi VK. Influence of NAA, GA₃ and Boric Acid on growth, yield and quality of strawberry cv. Chandler. Progressive Horticulture. 2009;41(1):113-115.
9. Lal D, Tripathi VK, Nayyer MA, Kumar S, Ahmed M, Siddiqui MW. Pre-harvest spray of Gibberellic Acid, NAA, and Calcium Nitrate on fruit retention, yield and quality of Kinnow Mandarin. Environment & Ecology. 2016;34(4C):2288-2292.
10. Tripathi VK, Viveka Nand. Effect of foliar application of Boron, Zinc and NAA on fruit retention, yield and quality attributes of aonla. Progressive Horticulture. 2022; 54(1):76-81.
11. Bhadauria AS, Tripathi VK, Singh A, Gupta S. Effect of foliar application of plant bio-regulators and micronutrients on fruit retention, yield and quality attributes of Aonla. Progressive Research-An International Journal. 2018;13(3):216-219.
12. Yadav DN, Chaturvedi OP. Influence of GA₃ and trace elements on fruit drop, growth and quality of Ber (*Zizyphus mauritiana* Lamk.) cv. Banarasi Karaka. Farm Science Journal. 2005; 14:27-28.
13. Chaurasiya, Gangadhar R, Sharma A, Tiwari S, Goyal G, Bhadauria AS, Singh AP, Yadav A. Influence of foliar application of GA₃ with and without NAA on fruit drop, growth, yield and quality of ber (*Zizyphus mauritiana* Lamk.). International Journal Current Microbiol Applied Science. 2019; 8:45-56.
14. Tripathi VK, Singh A, Bhadauria AS, Gupta S. Influence of GA₃ and Naphthalene Acetic Acid alone and in combination on fruit drop, yield and quality of mango cv. Amrapali. Progressive Research-An International Journal. 2019;14(1):10-13.
15. Badal DS, Tripathi VK. Effect of foliar application of NAA and Boron on physico-chemical parameters of winter season guava (*Psidium guajava* L.) cv. Lucknow-49. The Pharma Innovation Journal. 2021b;10(9):928-932.
16. Shukla HS, Kumar V, Tripathi VK. Effect of Gibberellic Acid and Boron on development and quality of aonla fruits 'Banarasi'. Acta Horticulturae. 2011;890: 375-380.
17. Tripathi VK, Pandey SS, Kumar A, Dubey V, Tiwari P. influence of foliar application of gibberellic acid, calcium and boron on fruit drop, yield and quality attributes of aonla (*Embllica officinalis*) cv. NA7. Indian Journal of Agricultural Sciences. 2018; 88(11):1784-8.
18. Dubey, Vishal, Meena ML, Tripathi VK. Effect of plant bio-regulators and micronutrient on vegetative growth, yield and quality of strawberry cv. Chandler. Progressive Research-An International Journal. 2017;12(3):330-332.
19. Tripathi VK, Shukla PK. Influence of plant bio-regulators, boric acid and zinc sulphate on yield and fruit characters of strawberry cv. Chandler. Progressive Horticulture. 2007;39(2):154-158.
20. Singh K, Randhawa, and Singh K. Effect of growth regulators and fungicides on fruit drop, yield and quality of fruit in ber cv. Umran. Journal Research, PAU. 2001; 38(3-4):181-185.
21. Rathod RK, Ramdevputra MV, Jadeja SR, Parmar LS, Jivani LL. Effect of foliar application of micronutrients and growth regulator on fruit yield of Aonla (*Embllica officinalis* Gaertn.) cv. Gujarat Aonla-1. Journal of Pharmacognosy and Phytochemistry. 2019;8(5):133-137.

22. Singh NP, Malhi CS, Sharma RC. Effect of plant bioregulators (PBRs) on flowering, fruit yield and quality in mango cv. Dashehari. Horticultural Journal. 2005; 18:10-12.
23. Kale VS, Dod VN, Adpawar RN, Bharad SG. Effect of PGR on fruit characters and quality of ber. Crop Research. 2000; 20(2):327-333.
24. Verma S, Dwivedi AK, Tripathi VK. Effect of gibberellic acid and boron on growth, yield and yield attributory traits in strawberry (*Fragaria x ananassa Duch.*) under North Gangetic Plains. Biological Forum-An International Journal. 2021; 13(3):262-267.
25. Ram RB, Pandey, Sharad and Akhilesh K. Effect of plant growth regulators on fruit retention, physico-chemical parameters and yield of ber cv. Banarasi Karaka. Biochemical and cellular Archive. 2005; 5(2):229-232.

© 2023 Priya et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/110091>