



Effect of Different Levels of Cinder and Vermicompost on Germination and Seedling Growth of Papaya (*Carica papaya* c.v Red Lady) its Treatment with GA₃

Rishika Rai ^{a++*}, Samir E. Topno ^{a#} and Vijay Bahadur ^{at}

^a Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj-211007 (U. P.), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i71024>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118563>

Original Research Article

Received: 14/04/2024
Accepted: 18/06/2024
Published: 20/06/2024

ABSTRACT

An experiment was conducted to find out suitable growth hormone and growing medium for germination and seedling growth of papaya. The treatments included soaking the seed of papaya variety Red lady in GA₃(200ppm) for 24.00 hours except for control. The perforated polythene bags of 4 x 5 cm size (150 gauges thickness) were used for sowing. The bags were filled with a mixture

⁺⁺ M. Sc Scholar;

[#] Assistant Professor;

[†] Associate Professor and Head;

*Corresponding author: E-mail: rishikarai653@gmail.com;

Cite as: Rai, Rishika, Samir E. Topno, and Vijay Bahadur. 2024. "Effect of Different Levels of Cinder and Vermicompost on Germination and Seedling Growth of Papaya (*Carica Papaya* c.V Red Lady) Its Treatment With GA₃". *Journal of Advances in Biology & Biotechnology* 27 (7):641-47. <https://doi.org/10.9734/jabb/2024/v27i71024>.

of Soil, Vermicompost, Cinder and Cocopeat in the ratio of 1:1:1, 1:2:1, 1:1:2 and 2:1:1. Experiment was laid out in Completely randomized design nine treatment with three replication. Findings indicate that treatment T₉ Soil +Cocopeat +Vermicompost +Cinder (1:1:1:1) +GA₃ 200ppm performed best in terms of growth parameters viz., germination (77.37), days taken for completion of germination (15.27), seedling height (25.00 cm), number of leaves per seedling (13.53 cm), diameter of seedling (cm) (1.03), Leaf area (cm²) (97.64cm), Chlorophyll content (SPAD Value) (54.97), Length of longest tap root (4.52 cm) and Survival (%) (83.27) under Prayagraj agro-climatic condition.

Keywords: Papaya; seedling growth; different levels effect of cinder; tropical fruit.

1. INTRODUCTION

The papaya is a tropical fruit belonging to the Caricaceae family that possesses both medicinal and nutritional qualities. a fast-growing, continuously fruiting tropical fruit crop that is evergreen. This tropical fruit, which is part of the Caricaceae family, is incredibly healthful, reviving, and tasty. The papaya, which is native to tropical America, is often referred to as pawpaw or papaw. Soil-based and organic-based growing media are the two main types available for use in container nurseries. In contrast to soil-based media that mostly consists of field soil, organic-based media—which is composed of organic materials mixed with inorganic ingredients—such as compost, peat, coconut coir, or other organic materials—promotes superior root development. For their growing media, nurseries in temperate regions have access to a large variety of commercial goods, such as premixed mixtures of peat moss, vermiculite, and perlite. However, most nurseries in tropical regions lack simple and economical access to these resources, and even nurseries in temperate regions are working to substitute more locally sourced and sustainably produced materials for some of these elements. Growers frequently make their own medium in the tropics using materials that are readily available nearby [1]. Gibberellic acid is essential to plants, particularly to fruit crops like papaya and grapes. Plant cell elongation and expansion, particularly in horticultural crops [1]. One potential use for gibberellic acid is as a seed dormant breaker [2-4]. A highly powerful hormone called gibberellic acid is used to initiate fruit seed germination, particularly in papaya seeds. It is frequently utilized as a hormone in the growing business to promote the development of larger bundles, particularly Thompson seedless grapes and papaya [1] and Mishra et al., [5]. Gibberellic acid (also known as Gibberellin A₃, GA, and GA₃) is a hormone that is present in both fungus and plants. Early in seed germination, GA₃ seems to

primarily stimulate the gluconeogenic enzymes' activity. Cow urine contains about 1.21% N₂, 0.01% P₂O₅ and 1.35% K₂O (Subramaniam, 2005) with micronutrients Fe, Mn, Zn and Cu (Yawalkar, 1996). Media is a type of substrate that gives developing plants the necessary nutrients and structural support. Most of the propagation media used in nurseries to grow horticultural plants are either organic or inorganic. Typically grown from seeds, papayas pique the interest of scientists because they contain viscous sarcotesta that inhibits germination Desai et al., [6]. Thus, the purpose of this study was to investigate the effects of varying concentrations of cinder, vermicompost, and cocopeat on papaya seedling germination and growth, as well as to determine the culture costs of various treatments.

2. MATERIALS AND METHODS

The experiment was conducted at the Department of Horticulture, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Prayagraj, during the year 2023. The seed of papaya variety, Red Lady was purchased from the authentic sources in Prayagraj. The treatments included soaking the seed of papaya variety Red lady in GA₃(200ppm) for 24.00 hrs duration along with a control. Experiment was laid out in Completely randomized design nine treatment with three replication. The treatment combination viz., T₀ Soil, T₁ Soil+Vermicompost +Cinder (1:1:1)+ GA₃ 200ppm, T₂ Soil+Vermicompost +Cinder (1:2:1) + GA₃ 200ppm, T₃ Soil+Vermicompost +Cinder (1:1:2) + GA₃ 200ppm, T₄Soil+Vermicompost +Cinder (2:1:1) + GA₃ 200ppm, T₅ Cocopeat+Vermicompost +Cinder (1:1:1) + GA₃ 200ppm, T₆Cocopeat +Vermicompost +Cinder (1:2:1) + GA₃ 200ppm, T₇Cocopeat +Vermicompost +Cinder (1:1:2) + GA₃ 200ppm, and T₈ Cocopeat +Vermicompost +Cinder (2:1:1) + GA₃ 200ppm and T₉Soil+Cocopeat

+Vermicompost +Cinder (1:1:1:1) + GA₃ 200ppm. The polythene bags (4 x 5 cm 150 gauge) were purchased from local market of Prayagraj. The perforated polythene bags of 4 x 5 cm size (150 gauges thickness) were used for sowing. The bags were filled with a mixture of Soil, Vermicompost, Cinder and Cocopeat in the ratio of 1:1:1, 1:2:1, 1:1:2 and 2:1:1. Before sowing the seeds in the polythene bags, soil were treated with fungicide, i.e. copper oxychloride @ 2%. Each polythene bag had two seeds put in it. Water cane was regularly irrigated softly in the morning and evening. At twenty-four hours, treated papaya seeds were planted in polythene bags that had been correctly filled, tagged, and arranged according to the plan. Papaya crops are typically irrigated every 8 to 10 days in the winter and every 4 to 5 days in the summer, depending on the soil and climate variations. Water-cane irrigation was used to keep the right moisture content in seedlings planted in polythene bags. The pre-sowing-dip technique was used to treat the seeds. Growth regulators in the appropriate concentrations were added in sufficient amounts to the designated beakers.

3. RESULTS AND DISCUSSION

The information is displayed in Table 1. It was evident that the application of varying concentrations of cinder and vermicompost, along with seed treatment with GA₃, had a noteworthy impact on papaya seed germination and seedling growth. When gibberellic acid is applied to the embryo, it stimulates the manufacture of hydrolyzing enzymes, which in turn promotes the growth of the embryo and increases germination. Similar finding were reported by Bharche et al., [7], Babu et al., [8] in papaya and Mishra et al., [5]. It could be because the medium that include organic manures have organic acid in them. Consequently, higher germination percentages and minimum days to germination may have been aided by increased moisture availability and certain acids. Similar results were reported by Prasana et al., [9] in mango. Whereas, the maximum days taken for completion of germination (15.27) was recorded in T₉ Soil+Cocopeat +Vermicompost +Cinder (1:1:1:1) + GA₃ 200ppm followed by T₈ Cocopeat +Vermicompost +Cinder (2:1:1) + GA₃ 200ppm, T₁ Soil+Vermicompost +Cinder (1:1:1)+ GA₃ 200ppm and T₂ Soil+Vermicompost +Cinder (1:2:1) + GA₃ 200ppm. However, T₀ (Soil) reported the minimal number of days required for germination to be completed, which was 15.27.

When gibberellic acid is applied to the embryo, it stimulates the manufacture of hydrolyzing enzymes, which in turn promotes the growth of the embryo and increases germination. Similar finding were reported by Bharche et al., [7], Babu et al., [8] in papaya and Mishra et al., [5]. It could be because the medium that include organic manures have organic acid in them. Consequently, higher germination percentages and minimum days to germination may have been aided by increased moisture availability and certain acids. Similar results were reported by Prasana et al., [9] in mango. Whereas maximum seedling height was observed in T₉ Soil+Cocopeat +Vermicompost +Cinder (1:1:1:1) + GA₃ 200ppm with 25.00 cm, followed by T₈ Cocopeat +Vermicompost +Cinder (2:1:1) + GA₃ 200ppm, T₃ Soil+Vermicompost +Cinder (1:1:2) + GA₃ 200ppm, T₂ Soil+Vermicompost +Cinder (1:2:1) + GA₃ 200ppm and T₁ Soil+Vermicompost +Cinder (1:1:1)+ GA₃ 200ppm. While the minimum seedling height 18.77 cm respectively were observed in T₀ (Soil). The increased height observed in GA₃ 200 ppm seeds could be explained by the possibility that the papaya seedling's internal levels of GA₃ synthesis were insufficient, and that the external application of GA₃ increased cell multiplication and cell elongation, which improved plant growth Mishra et al., [5]. Growing medium not only acts as a growing place but also as a source of nutrient for plant growth. Media composition used influences the quality of seedling Wilson et al., [10]. Humic acids (vermicompost) applied in the medium increased plant height, leaf area and dry weight of peppers, tomatoes and marigold Arancon et al., [11]. While as maximum Number of leaves per seedling was observed in T₉ Soil+Cocopeat +Vermicompost +Cinder (1:1:1:1) + GA₃200ppm with 13.53 cm, followed by T₈ Cocopeat +Vermicompost +Cinder (2:1:1) + GA₃ 200ppm, T₃ Soil+Vermicompost +Cinder (1:1:2) + GA₃ 200ppm, T₂ Soil+Vermicompost +Cinder (1:2:1) + GA₃ 200ppm and T₁ Soil+Vermicompost +Cinder (1:1:1)+ GA₃ 200ppm. While the minimum Number of leaves per seedling 9.22 respectively were observed in T₀ (Soil). The explanation for the increase in leaves could be that GA₃ stimulates the plant's physiological processes and chemicals' stimulatory effects to produce new leaves more quickly. Mishra & associates [5]. Growing media serves as a space for plants to grow as well as a supply of nutrients. The type of media utilized affects the seedling's quality, according to Wilson et al. [10]. When added to the medium, humic acids (vermicompost) improved the height, leaf

Table 1. Effect of different levels of cinder, vermicompost and GA₃ on germination and seedling growth of Papaya

Treatments No.	Germination (%)	Days taken for completion of germination	Seedling height (cm)			Number of leaves per seedling			Diameter of seedling (cm)			Leaf area (cm ²)		
			30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS	30 DAS	60 DAS	90 DAS
T ₀	47.05	22.94	6.71	12.18	18.77	4.03	6.08	9.22	0.15	0.27	0.50	15.33	28.77	56.51
T ₁	74.08	16.40	9.88	16.34	22.84	6.13	9.19	12.39	0.26	0.53	0.97	36.33	46.45	81.77
T ₂	75.72	16.63	9.31	16.80	22.73	6.38	9.49	12.52	0.28	0.53	0.79	32.67	48.54	82.47
T ₃	68.00	17.52	9.95	15.69	21.67	6.07	8.75	12.28	0.24	0.50	0.77	31.38	51.56	85.00
T ₄	58.00	18.23	7.67	13.77	19.81	4.96	7.49	10.71	0.18	0.35	0.57	28.39	51.79	85.43
T ₅	54.51	19.51	8.01	13.84	19.83	5.24	7.33	10.73	0.18	0.33	0.57	28.53	52.15	80.07
T ₆	51.33	19.21	7.45	14.00	20.36	5.18	7.19	10.53	0.17	0.32	0.54	33.97	54.08	83.44
T ₇	61.17	17.89	9.65	15.54	21.76	5.18	8.20	11.87	0.23	0.45	0.62	31.64	51.15	83.82
T ₈	71.38	16.41	9.91	17.10	22.04	6.39	9.77	13.15	0.31	0.56	1.00	33.53	55.74	96.82
T ₉	77.37	15.27	10.66	17.74	25.00	6.58	9.87	13.53	0.33	0.60	1.03	35.05	59.17	97.64
F-Test	S	S	S	S	S	S	S	S	S	S	S	S	S	S
S.Ed.(±)	2.16	0.65	0.21	0.61	0.60	0.14	0.19	0.27	0.01	0.01	0.03	0.80	1.79	1.43
C.D. at 0.5%	4.54	1.38	0.44	1.28	1.27	0.29	0.40	0.57	0.02	0.03	0.07	1.69	3.78	3.00
CV	4.15	4.49	2.92	4.89	3.45	3.06	2.80	2.84	6.97	4.47	5.69	3.22	4.41	2.10

Table 2. Effect of different levels of cinder, vermicompost and GA₃ on germination and seedling growth of papaya

Treatments No.	Chlorophyll content (SPAD Value)	Length of longest tap root (cm)	Survival (%)
T ₀	44.71	4.52	58.64
T ₁	48.02	5.49	80.69
T ₂	48.44	5.38	86.15
T ₃	50.62	5.26	75.64
T ₄	51.65	5.16	66.30
T ₅	47.67	5.48	61.19
T ₆	47.63	5.51	59.54
T ₇	49.87	6.14	71.60
T ₈	53.85	6.16	82.36
T ₉	54.97	6.50	83.27
F-Test	S	S	S
S.Ed.(±)	0.80	0.19	0.99
C.D. at 0.5%	1.69	0.41	2.09
CV	1.98	4.32	1.68

area, and dry weight of tomatoes, peppers, and marigolds [11]. The maximum diameter of seedling (cm) was observed in T₉ Soil+Cocopeat +Vermicompost +Cinder (1:1:1:1) + GA₃ 200ppm with 1.03 cm, followed by T₈ Cocopeat +Vermicompost +Cinder (2:1:1) + GA₃ 200ppm, T₃ Soil+Vermicompost +Cinder (1:1:2) + GA₃ 200ppm, T₂ Soil+Vermicompost +Cinder (1:2:1) + GA₃ 200ppm and T₁ Soil+Vermicompost +Cinder (1:1:1)+ GA₃ 200ppm. While the minimum diameter of seedling (cm) 0.50 respectively were observed in T₀ (Soil). The administration of GA₃ may have caused the plant's diameter to expand because it increased the somatic absorption of nutrients, which led to cell elongation and a rise in plant height [12]. Similar, results were also recorded by Dhankar and Singh [13] in aonla and Bharche et al., [7] in papaya Mishra et al., [5]. Growing media serves as a space for plants to grow as well as a supply of nutrients. The type of media utilized affects the seedling's quality Wilson et al., [10]. When applied to the medium, humic acids (vermicompost) improved the height, leaf area, and dry weight of peppers, tomatoes, and marigold plants Arancon et al., [11]. The maximum Leaf area (cm²) was observed in T₉ Soil+Cocopeat +Vermicompost +Cinder (1:1:1:1) + GA₃ 200ppm with 97.64 cm, followed by Cocopeat +Vermicompost +Cinder (2:1:1) + GA₃ 200ppm, Cocopeat +Vermicompost +Cinder (1:2:1) + GA₃ 200ppm and Cocopeat +Vermicompost +Cinder (1:1:2) + GA₃ 200ppm. While the minimum Leaf area (cm²) 56.51 respectively were observed in T₀ (Soil). The explanation for the increase in leaves could be that GA₃ stimulates the plant's physiological

processes and chemicals' stimulatory effects to produce new leaves more quickly. Mishra et al., [10]. Growing media serves as a space for plants to grow as well as a supply of nutrients. The type of media utilized affects the seedling's quality Wilson et al., [10]. When applied to the medium, humic acids (vermicompost) improved the height, leaf area, and dry weight of peppers, tomatoes, and marigold plants. Whereas, the maximum Chlorophyll content (SPAD Value) (54.97) was recorded in T₉ Soil+Cocopeat +Vermicompost +Cinder (1:1:1:1) + GA₃ 200ppm followed by T₈ Cocopeat +Vermicompost +Cinder (2:1:1) + GA₃ 200ppm, T₄ Soil+Vermicompost +Cinder (2:1:1) + GA₃ 200ppm and T₃ Soil+Vermicompost +Cinder (1:1:2) + GA₃ 200ppm. While the minimum Chlorophyll content (SPAD Value) 44.71 respectively were recorded in T₀ (Soil). This appears to be the result of water and nutrients being mobilized and transported at a faster rate, which may have encouraged the synthesis of more photosynthetic products and moved them to different plant parts, potentially leading to improved seedling growth and an increase in fresh and dry weight. These results are consistent with the papaya research conducted by Lay et al. (2015). The different levels of GA₃ did not affect significantly to increase chlorophyll content. Whereas, the maximum Length of longest tap root (cm) (6.50) was recorded in T₉ Soil+Cocopeat +Vermicompost +Cinder (1:1:1:1) +GA₃ 200ppm followed by T₈ Cocopeat +Vermicompost +Cinder (2:1:1) +GA₃ 200ppm, T₂ Soil+Vermicompost +Cinder (1:2:1) +GA₃ 200ppm and T₁ Soil+Vermicompost +Cinder (1:1:1)+GA₃ 200ppm. While the minimum Length

of longest tap root (cm) 4.52 respectively were recorded in T₀ (Soil). This could be because GA3 promotes nutrient uptake by the somatic cells, which leads to cell elongation and an increase in the length and quantity of secondary roots in the tap root Mishra et al., [5]. Whereas, the maximum Survival (%) (83.27) was recorded in T₉ Soil+Cocopeat +Vermicompost +Cinder (1:1:1:1) +GA3 200ppm followed by T₈ Cocopeat +Vermicompost +Cinder (2:1:1) +GA3 200ppm, T₈ Cocopeat +Vermicompost +Cinder (2:1:1) +GA3 200ppm, T₂ Soil+Vermicompost +Cinder (1:2:1) +GA3 200ppm and T₁ Soil+Vermicompost +Cinder (1:1:1)+GA3 200ppm. While the minimum Survival (%) 58.64 respectively were recorded in T₀ (Soil). This could be because GA3 caused the seed coat to weaken, which allowed radicals and plumules to emerge and have a favorable impact on the initiation of roots and shoots. In addition to this, GA3 promotes cell elongation and expansion, which improves root and shoot growth and increases seedling survival [14].

4. CONCLUSION

Based on the above finding, it is concluded that treatment T₉ Soil+Cocopeat +Vermicompost +Cinder (1:1:1:1) + GA3 200ppm performed best in terms of germination percentage (77.37%), days taken for completion of germination (15.27), and growth parameters were seedling height (25.00 cm), number of leaves per seedling (13.53), diameter of seedling (1.03cm), Leaf area (97.64cm²), Chlorophyll content (SPAD Value) (54.97), Length of longest tap root (4.52cm) and Survival percentage (83.27%) under Prayagraj agro-climatic condition.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bharadwaj RL. Effect of growing media on seed germination and seedling growth of

papaya cv. 'Red lady'. African Journals of Plant Sciences. 2014;8(4):178–184.

2. Nejatizadeh F. Effect of vermicompost and nitrogen fertilizer on the growth and production of aloe vera. Int. J. Plant Soil Sci. 2024, Mar 9;36(4):337-45. [cited 2024 Jun. 2] Available: <https://journalijpss.com/index.php/IJPSS/article/view/4486>

3. Vyas D, Sonkar P, Kanpure RN, Ausari PK, Ninama N, Suresh. Effect of soil propagation media and bio-fertilizers on seedling germination and seedling vigour in aonla (*Emblica officinalis* Gaertn.). Int. J. Environ. Clim. Change. 2023, Sep 25;13(10):3962-73. [cited 2024 Jun. 2] Available: <https://journalijecc.com/index.php/IJECC/article/view/3070>

4. Blouin M, Barrere J, Meyer N, Lartigue S, Barot S, Mathieu J. Vermicompost significantly affects plant growth. A meta-analysis. Agronomy for Sustainable Development. 2019, Aug;39:1-5.

5. Mishra Unmesh, Vijay Bahadur, Prasad VM, Pushendra Verty, Ashutosh Kumar Singh, Saket Mishra, Swaroop N. Influence of GA3 and growing media on growth and seedling establishment of papaya (*L. Carica papaya.*) cv. pusa nanha. Int.J.Curr.Microbiol.App.Sci. 2017;6(11):415-422.

6. Desai Amit, Panchal Bharat, Trivedi Ashwin, Prajapati Dinesh. Studies on seed germination and seedling growth of papaya (*Carica papaya* L.) CV. madhubindu as influenced by media, GA3 and cow urine under net house condition. Journal of Pharmacog Nosy and Phytochemistry. 2017;6(4):1448-1451

7. Barche SK, Singh K, Singh DB. Response of seed treatment on germination, growth, survivability and economics of different cultivars of papaya (*Carica papaya* L.). Acta Horticulture. 2010;851:279-281.

8. Dhinesh Babu K, Patel RK, Singh A, Yadav DS, De LC, Deka BC. Seed germination, seedling growth and vigour of papaya under north East Indian condition. Acta Horticulture. 2010;851:299-306.

9. Parasana JS, Leua HN, Ray NR. Effect of different growing medias mixtures on germination and seedling growth of mango (*Mangifera indica* L.) cultivars under net house conditions. The Bioscan. 2014;8(3): 897-900.

10. Wilson SB, Stoffella PJ, Graetz DA. Use of compost as a media amendment for containerized production of two subtropical perennials. *Journal Environmental Horticulture*. 2011;19:37–42.
11. Arancon NQ, Lee S, Edwards CA, Atiyeh R. Effect of humic acids derived from cattle, food and paper – waste vermicompost on growth of green house plants. *Pedobiologia*. 2004;47: 741–744.
12. Feucht JR, Watson DP. The effect of gibberellins on internal tissues of pea (*Phaseolus vulgaris* L.). *American Journal of Botany*. 1958;45:520-522.
13. Dhankhar DS, Singh M. Seed germination and seedling growth in aonla (*Phyllanthus emblica* Linn.) as influenced by gibberellic acid and thiourea. *Field Crop Research*. 1996;12(3):363-366.
14. Rahangdale P. Effect of GA3 and date of sowing on seed germination, growth and survival of custard apple (*Annona squamosa* L.) seedlings. M.Sc. (Hort.) Thesis, J. N. Krishi Vishwa Vidyalaya, Jabalpur (M.P.). 2015;4849.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/118563>