



## **Assessment of Burnout in Healthcare Professionals of Pakistan Amid COVID-19 – A Cross-Sectional Study**

**Nisha Zahid<sup>1\*</sup>, Marium Syed<sup>1</sup>, Sharmake Hersi<sup>1</sup>, Syed Hasan Danish<sup>1</sup>  
and Farah Ahmed<sup>1</sup>**

<sup>1</sup>*Ziauddin Medical College, Ziauddin University, Karachi, Pakistan.*

### **Authors' contributions**

*This work was carried out in collaboration among all authors. The concept of this study, data analysis, drafting, and finalizing of the results were done by authors NZ and MS. The article was critically reviewed and finally approved by author SHD. While the data collection and analysis based work was performed with the help of author SH. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJTDH/2021/v42i530456

#### Editor(s):

- (1) Dr. Maria Aparecida Visconti, Instituto de Biociências, Universidade de São Paulo, Brazil.
- (2) Dr. Rameshwari Thakur, Muzaffarnagar Medical College, India.
- (3) Dr. Chan-Min Liu, Xuzhou Normal University, China.
- (4) Dr. Giuseppe Murdaca, University of Genoa, Italy.

#### Reviewers:

- (1) Ki-Young Kim, Kyung Hee University, Republic of Korea.
  - (2) Manisha Naithani, AIIMS Rishikesh, India.
  - (3) Khalid Al-Johani, King Abdulaziz University, Kingdom of Saudi Arabia.
- Complete Peer review History: <http://www.sdiarticle4.com/review-history/66682>

**Original Research Article**

**Received 04 March 2021**  
**Accepted 14 April 2021**  
**Published 21 April 2021**

### **ABSTRACT**

**Introduction:** According to WHO latest by March 19th 2021, it has now become a global threat with globally 122,665,852 confirmed cases and 2,707,206 deaths involving 220 countries.. NIH Pakistan has reported the national tally of COVID 19 cases in Pakistan to 543,214 and total deaths 11,683, making it an extremely serious issue in Pakistan like in the rest of the world. Physicians, consultants, paramedics and other staff working in hospitals are at risk due to the emergence of COVID-19 which also is responsible for severe burnout in the specialties who are working at the frontline. Burnout affects job performance, job satisfaction, interpersonal relationships, and vulnerability to illnesses [1]. This research aims at finding the burnout in healthcare professionals of Karachi, Pakistan due to the current situation of COVID-19 pandemic.

\*Corresponding author: E-mail: [dr.nishaa.shaikh@gmail.com](mailto:dr.nishaa.shaikh@gmail.com);

**Materials and Methods:** It is a cross-sectional study carried out in the tertiary care hospitals of Karachi, Pakistan from Feb'2020 – Apr' 2020 in which the questionnaire based on Malsch burnout inventory were distributed amongst the healthcare professionals including doctors and paramedics, working at the front in departments of emergency, covid special units and other areas. A scoring system of 1-4 was applied (strongly disagree-strongly agree).

**Results:** The results of this study showed that highest burnout was faced by healthcare professionals in the category of depersonalization and personal accomplishment amid COVID-19 while the association between the burnout and the demographic factors mentioned was also found to be significant. Moreover, the challenges faced by the healthcare professionals to the maximum owing to this pandemic were found to be lack of presence of personal protective equipment and fear to spread the infection to family.

**Conclusion:** The study concluded that the current pandemic situation of COVID-19 has been significantly associated with the burnout and stress amongst the health care professionals in tertiary hospitals of Karachi, Pakistan, particularly those who are working on the frontline.

*Keywords: COVID-19; healthcare professionals; burnout; infection.*

## 1. INTRODUCTION

It is a well-established fact that Burnout has effects on job performance, job satisfaction, interpersonal relationships, and vulnerability to illnesses [1]. Physicians and consultants working in the specialties at the front lines of care such as emergency medicine, general internal medicine, neurology, and family medicine) are at the highest risk of developing burnout because of the stressful nature of work. According to a study, more than half of the doctors and paramedics in United States are suffering substantial symptoms of exhaustion [2]. The similar finding are seen in United Kingdom, A study conducted in 2017 showed the prevalence of burnout in general practitioners and consultants to be 54% [3]. Healthcare professionals include doctors, nurses and paramedics, are taking important life-changing decisions every moment and that is why their mental and physical wellbeing is an outright obligation [4]. Various studies have been conducted throughout the world to gauge burnout in health care professionals. One such study conducted by Ehsan Zarei, showed that 52.9% of primary health care workers had burnout [5].

The emergence of the COVID-19 has been the hot cake since December 2019 and its consequences has led to fears, uncertainties, and anxiety among individuals worldwide and is by far the largest outbreak of atypical pneumonia since the severe acute respiratory syndrome (SARS) outbreak in 2003 [6]. At this time, there are no specific vaccines or treatments for COVID-19 [7]. According to WHO latest by March 19<sup>th</sup> 2021, it has now become a global threat with globally 122,665,852 confirmed cases and 2,707,206 deaths involving 220 countries.

NIH Pakistan has reported the national tally of COVID 19 cases in Pakistan to 543,214 and total deaths 11,683 hence it has become a serious issue in Pakistan like rest of the world (National Institute of Health Pakistan 2021).

As far as a pandemic is concerned, several studies have been led to evaluate burnout in health professionals during the outbreak of MERS (middle east respiratory syndrome and SARS (severe acute respiratory syndrome) [8,9]. Currently, there is no known information on the psychological impact and mental health of the healthcare professionals dealing with COVID-19 pandemic specifically however some studies have been conducted in different settings of Pakistan mentioning the increase in stress and depression amongst employees of hospitals fighting against this pandemic [10]. This is especially relevant with the ambiguity surrounding an outbreak of such supreme extent. As far as the research on COVID-19 is concerned, most of the studies focus on identifying the epidemiology and clinical characteristics of infected patients [11]. To the best of our knowledge, no study till date has been conducted to assess the burn out in healthcare professionals who are working day and night to fight against COVID-19. This research aims at finding the burnout in healthcare professionals of Karachi, Pakistan due to the current situation of COVID-19 pandemic.

## 2. MATERIALS AND METHODS

We carried out a cross-sectional study from February' 2020 to April'2020, at the tertiary care hospitals of Karachi. The health care workers

including doctors, nurses and paramedical staff working full time in COVID-19 special units of the hospital are included in our study. While All the other professional Health Care Workers who were not employed by the COVID-19 special units of the hospital and any other staff members including IT department, management and guards were excluded our study. The sample size was calculated through RAOSOF software keeping the population size of 2000 and confidence level of 95 % (n=323) and the sampling done through convenience sampling technique. The participants were informed about the purpose of the study and consent was taken prior to data collection. Ethical approval for the study was taken from the institutional ethics committee. The participants were asked to fill a questionnaire which was self-administered comprised of demographic data and a validated Maslach Burnout Inventory (MBI) taken from www.mindgarden.com. It is used to measure burnout specifically in people working in healthcare. MBI constitutes of 22 items covering three elements of burnout: emotional exhaustion EE, depersonalization DP and personal accomplishment PA. Emotional exhaustion refers to a state of feeling emotionally drained and worn out and indicates negative attitude towards work. The cut of values for EE is 0-17 for low level, 18-29 for medium level and more than 29 for high level burnout. Depersonalization is characterized by loss of interest in job and colleagues. The cut off values for depersonalization is 0-5 for low level, 6-11 for medium level and more than 12 for high-level burnout. Personal accomplishment is a dimension of burnout which refers to associated with the feeling of competence, high self-efficacy and sense of accomplishment reduced personal accomplishment indicates burnout. The cut off values for personal accomplishment includes 0-33 as high level burnout, 34-39 as medium level and more than 39 as low-level burnout. Thus, it is self-evident that high scores in sections a and b and low scores in section c are associated with high burnout [4]. A scoring system of 1-4 was applied, (strongly disagree-strongly agree). Data analysis was conducted on SPSS Version 21. Mean and standard deviation was calculated for numerical data. For categorical data we took out frequency and percentages. P value of less than 0.05 was taken as significant.


### 3. RESULTS

The maximum number of respondents i.e. n=217 (66.0%) were among age group of 25-39 years of

age and only n=9 (2.7%) were in 40-59. All participants were Pakistan nationals of which n=155 (47.4%) were males and n=167 (51.7%) were females. There was no big difference in percentages of singles and married n=135 (43.5%) and (n=183) 59% and participants who were divorced were only n=2 (0.6%) Most participants n=190 (57.8%) had higher education and only n=35 (10.6%) had secondary level education. Of 329 participants n=271 (82.4%) were doctors and remaining n=58 (17.6%) comprised of paramedics. Respondents employed in private work setup were n=162 (49.2%) and those employed in public work set up were n=167 (50.8%). Amid corona virus outbreak duty hours of participants were varying, n=176 (53.5%) were working for 10-12 hours while n=81 (24.6%) was working 12-24 hrs and n=33 (10%) of hcp working for more than 24 hrs. Hence the time they get to relax and sleep also vary between 6-8 hours n=233 (70.8%) and 4-6 hrs. n=71 (21.6%). Majority n=324 (98.5%) were allowed to meet family, however, maximum family hours varied between 6-8 hrs, n=153 (46.5) and 4-6 hours, n=129 (39.2). The n=274 (83.3%) of hcp used caffeine while n=51(15.5%) participants used cigarettes and n=4 (1.2%) takes medicines to relieve stress.

As the figure shows, our healthcare professionals are emotionally exhausted and displayed the characteristics of moderate n=181(55%) and high burnout n=146 (44.3%). The n=312 (94.8%) participants presented with individualities of high burnout in the factor of personal accomplishment while n=321 (97.5%) of them also exhibited high burnout when asked about aspects of depersonalization during these tough times of COVID-19.

According to the current scenario of COVID-19 we have asked our respondents about the biggest challenge in coping with the present situation and n=274 (83.3%) were worried about the nonavailability of personal protective equipment, n=203 (61.7%) were in stress because of little or no information on coronavirus transmission and treatment, while the biggest fear our healthcare professionals are going through with is, concern to spread the infection to their family and friends n=295 (89.9%) and the consequences covid 19 might cause on their health. And lastly, we also compared burnout amongst the individual health professionals with the demographic data provided and the results are shown in the table.



**Fig. 1. burnout for all three categories, EE(emotional exhaustion), PA(personal accomplishment) and DE(depersonalization)**

#### 4. DISCUSSION

This study was much needed to find out burnout in healthcare professional due to coronavirus disease 2019 (COVID-19) pandemic which has become one of the major health crises for this generation as it has affected the people across the globe [12]. Our healthcare professionals who are fighting with this pandemic on the frontline have been under constant stress which is a great influencing factor for job related burnout. As the table shows more than 70% of our doctors and paramedics both in private or public setup are working for about 10-24 hrs.to cope up with this pandemic. Long duty hours have always been key factor in generating stress, our results are in agreement to various studies including a study conducted in SaudiArabia that also concluded that the majority of residents work 60 or more hours per week, and there was a very high degree of burnout amongst residents [13]. Physician burnout is a significant problem in the medical profession, and work overload is the main contributor to it fatigue, stress, and accentuate resiliency [14]. Few health care workers chose not to meet their family (1.5%) and most of them were allowed to meet their family (98.5%) since COVID-19 is a community spread infection due to which the risk of infection is evenly distributed across the population hence health care workers feel isolated by their family because of their hospital work (40.1%) and this isolation also leads to stress and anxiety [15].

Fig. 1 shows burnout in all three categories i.e emotional exhaustion, personal accomplishment and depersonalization due to COVID-19 outbreak. As far as we know no study has reported burnout in healthcare professionals during the times of COVID-19. A study took place in Korea that assessed the burnout in nurses in the times of MERS-COV and resulted in that there is significant correlation between stress and burnout specially emotional exhaustion and depersonalization [16]. The flaws of health care systems all over the world are now mainly highlighted. Even the highly developed countries with an extraordinary smoothly working health care systems are unable to meet with the demands of controlling the situation in this pandemic including the number of patients in need of treatment, primarily, intensive care with endotracheal intubation and mechanical ventilation [17]. The challenges health professionals are facing in this pandemic has also played a great role in increasing anxiety and stress and the results suggest that most of the respondents think that little or no information about the COVID-19 is an enormous issue to deal with [18]. The unavoidable stress, fear and anxiety is reported to be insightful in healthcare professionals and other front line workers, due to the fact that very limited knowledge is available regarding COVID-19 hence, the emotional and physical surge has been increased (Harapan et al., 2020). Another study pointed out the stress and fear in surgical residents in tertiary care hospital of Pakistan [19]. A questionnaire based study was conducted to evaluate the

**Table 1. comparison of burnout with the demographical data**

	Low burnout		Moderate burnout		High burnout		p-value
	n	%	n	N	%		
<b>Age group</b>							
18-24	0	0.0%	13	13.0%	87	87.0%	.677
25-39	4	1.8%	23	10.6%	190	87.6%	
40-59	0	0.0%	0	0.0%	9	100%	
>60	0	0.0%	0	0.0%	3	100%	
<b>Gender</b>							
Males	4	2.5%	15	9.4%	140	88.1%	.86
Females	0	0.0%	21	12.4%	149	87.6%	
<b>Marital status</b>							
Single	1	0.7%	14	9.8%	128	89.5%	.864
Married	3	1.6%	22	12.0%	159	86.4%	
Divorced	0	0.0%	0	0.0%	2	100%	
<b>Working hours</b>							
8hrs	0	0.0%	5	12.8%	34	87.2%	.713
10-12hrs	3	1.7%	19	10.8%	154	87.5%	
12-24hrs	0	0.0%	10	12.3%	71	87.7%	
>24hrs	1	3.0%	2	6.1%	30	90.9%	
<b>Educational level</b>							
Matric	0	0.0%	5	38.5%	8	61.5%	.016
Secondary	0	0.0%	2	5.7%	33	94.3%	
University	2	2.2%	5	5.5%	84	92.3%	
Higher	2	1.1%	24	12.6%	164	86.3%	
<b>Designation</b>							
Doctor	4	1.5%	29	10.7%	238	87.8%	.626
Paramedical staff	0	0.0%	7	12.1%	51	87.9%	
<b>Work setup</b>							
Private	2	1.2%	22	13.6%	138	85.2%	.319
Public	2	1.2%	14	8.4%	151	90.4%	
<b>Sleeping hours</b>							
8-10hr	0	0.0%	0	0.0%	4	100%	.013
6-8hr	2	0.9%	28	12.0%	203	87.1%	
4-6hr	0	0.0%	8	11.3%	63	88.7%	
<4hr	2	9.5%	0	0.0%	19	90.5%	
<b>Meet_family</b>							
Yes	4	1.2%	36	11.1%	284	87.7%	.704
No	0	0.0%	0	0.0%	5	100%	
<b>Family hours</b>							
12-16hr	0	0.0%	0	0.0%	12	100%	.077
6-8hr	2	1.3%	25	16.3%	126	82.4%	
4-6hr	1	0.8%	7	5.4%	121	93.8%	
<4hr	1	2.9%	4	11.4%	30	85.7%	
<b>Stress reliever</b>							
Caffeine	4	1.5%	23	8.4%	247	90.1%	.000
Cigarette	0	0.0%	10	19.6%	41	80.4%	
Medicine	0	0.0%	3	75%	1	25%	

knowledge of healthcare professionals about COVID 19 which suggests that there were significant knowledge gaps present in health care professionals and it is critically to improve the knowledge and perceptions of HCWs [20]. We need to cope with this challenge by providing

informational sessions on how to deal with this pandemic and all latest researches should be shared. One such study was conducted which showed that all healthcare professionals, especially nurses who were constantly pressured to work in highly infected areas of the hospital

were given adequate support, clinical supervision and briefing and debriefing sessions to alleviate potential burnout among nurses and other healthcare workers [21].

Another challenge is availability of personal protective equipment. Medical personnel protection is of particular importance because of the risk of infecting other members of medical teams, including not only physicians, nurses or paramedics, but also other support personnel necessary to maintain the continuity of care for patients [22]. Based on the observations of other countries' pandemic outbreak, the maximum standing should be given to infection control preparedness in Pakistan's healthcare system [23]. The results also showed that 89.9% of respondents are horrified due to the spread of this infection to their family. A study conducted in India suggested that The health care professionals who are dealing with confirmed/suspected COVID19 patients are being quarantined and isolated and the rest of them are separated from their family concerning the fear of transmitting the disease to their own family members [24].

One of The major contributing factor for causing burnout in health care professionals is long working shifts also our frontline workers are not on high risk themselves but also their families are equally exposed [25]. There is lack of preparedness especially in Pakistan which is further highlighted due to Rapid spread of covid19. The public hospitals were dealing the situation in a traditional way which was something under required [26].

We conducted this study to evaluate the burnout amongst health care workers at the peak time of COVID-19 in Pakistan which is a strong point of this study, also that we modified the questionnaire (MBI) to find out the challenges which are health care workers were going through due to the pandemic. The limitations of the study includes that this study could have been done on a large scale to see a bigger picture.

## 5. CONCLUSION

The study concluded that the current pandemic situation of COVID-19 has particularly associated with the burnout and stress amongst our health care professionals who are working on the frontline which has shown to be associated with various factors including demographics, emotional exhaustion, personal accomplishment,

depersonalization and challenges to cope up with COVID-19 as mentioned in the study.

## CONSENT AND ETHICAL APPROVAL

As per international standard or university standard guideline participant consent and ethical approval has been collected and preserved by the authors.

## ACKNOWLEDGEMENT

We are thankful to Dr. Farah Ahmed (Associate professor, department of Community Medicine) for her constant help and valuable input regarding the whole idea of this study. We would also like to thank Dr.Akhtar Ali for his support and help about data analysis.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Chemali Z, Ezzeddine F, Gelaye B, Dossett M, Salameh J, Bizri M, Dubale B, Fricchione G. Burnout among healthcare providers in the complex environment of the Middle East: A systematic review. *BMC public health*. 2019;19:1337.
2. Dyrbye LN, Shanafelt TD, Sinsky CA, Cipriano PF, Bhatt J, Ommaya A, West CP, Meyers D. Burnout among health care professionals: A call to explore and address this underrecognized threat to safe, high-quality care. *NAM perspectives*; 2017.
3. Imo UO. Burnout and psychiatric morbidity among doctors in the UK: A systematic literature review of prevalence and associated factors. *BJPsych bulletin*. 2017;41:197-204.
4. Baruah A, Das S, Dutta A, Das B, Sharma T, Hazarika M. degree and factors of burnout among emergency healthcare workers in India. *International journal of scientific research (Ahmedabad, India)*. 2019;8:41.
5. Zarei E, Ahmadi F, Sial MS, Hwang J, Thu PA, Usman SM. Prevalence of burnout among primary health care staff and its

- predictors: A study in Iran. *International journal of environmental research and public health*. 2019;16:2249.
6. Ahorsu DK, Lin CY, Imani V, Saffari M, Griffiths MD, Pakpour AH. The fear of COVID-19 scale: Development and initial validation. *International Journal of Mental Health and Addiction*. 2020;1.
  7. Watkins J. Preventing a COVID-19 pandemic. *British Medical Journal Publishing Group*; 2020.
  8. Kim JS, Choi JS. Factors influencing emergency nurses' burnout during an outbreak of middle east respiratory syndrome coronavirus in Korea. *Asian nursing research*. 2016;10: 295-299.
  9. Maunder RG, Lancee WJ, Balderson KE, Bennett JP, Borgundvaag B, Evans S, Fernandes CM, et al. Long-term psychological and occupational effects of providing hospital healthcare during SARS outbreak. *Emerging infectious diseases*. 2006;12:1924.
  10. Bukhari N, Rasheed H, Nayyer B, Babar ZUD. *Pharmacists at the frontline beating the COVID-19 pandemic*. Springer; 2020.
  11. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, Ho RC. Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in china. *International Journal of Environmental Research and Public Health*. 2020;17:1729.
  12. Shanafelt T, Ripp J, Trockel M. Understanding and addressing sources of anxiety among health care professionals during the COVID-19 pandemic. *Jama*; 2020.
  13. Hameed TK, Masuadi E, Al Asmary NA, Al-Anzi FG, Al Dubayee MS. A study of resident duty hours and burnout in a sample of Saudi residents. *BMC medical education*. 2018;18:180.
  14. Patel RS, Bachu R, Adikey A, Malik M, Shah M.. Factors related to physician burnout and its consequences: a review. *Behavioral Sciences*. 2018;8: 98.
  15. Kumar S. Burnout and doctors: Prevalence, prevention and intervention. *Healthcare, Multidisciplinary Digital Publishing Institute*. 2016;37.
  16. Kim Y, Seo E, Seo Y, Dee V, Hong E. Effects of middle east respiratory syndrome coronavirus on post-traumatic stress disorder and burnout among registered nurses in South Korea; 2018.
  17. Chughtai AA, Khan W. Use of personal protective equipment to protect against respiratory infections in Pakistan: A systematic review. *Journal of infection and public health*; 2020.
  18. Nagesh S, Chakraborty S. The frontline health workforce amidst the COVID-19 crisis: Challenges and recommendations. *Journal of Global Health*. 2020;Saving10.
  19. Osama M, Zaheer F, Saeed H, Anees K, Jawed Q, Syed SH, Sheikh BA. Impact of COVID-19 on surgical residency programs in Pakistan; A residents' perspective. Do programs need formal restructuring to adjust with the " new normal"? A cross-sectional survey study. *International Journal of Surgery*; 2020.
  20. Bhagavathula AS, Aldhaleei WA, Rahmani J, Mahabadi MA, Bandari DK. Knowledge and perceptions of COVID-19 among health care workers: Cross-sectional study. *JMIR Public Health and Surveillance*. 2020;6:e19160.
  21. Lopez V, Chan KS, Wong YCJ. Nursing care of patients with severe acute respiratory syndrome in the intensive care unit: Case reports in Hong Kong. *International journal of nursing studies*. 2004;41:263-272.
  22. Smereka J, Szarpak L. The use of personal protective equipment in the COVID-19 pandemic era. *The American journal of emergency medicine*; 2020.
  23. Mukhtar S. Preparedness and proactive infection control measures of Pakistan during COVID-19 pandemic outbreak. *Research in Social & Administrative Pharmacy*; 2020.
  24. Li S, Wang Y, Xue J, Zhao N, Zhu T. The impact of COVID-19 epidemic declaration on psychological consequences: a study on active Weibo users. *International*

- journal of environmental research and public health. 2020;17:2032.
25. Sasangohar F, Jones SL, Masud FN, Vahidy FS, Kash BA. Provider burnout and fatigue during the COVID-19 pandemic: Lessons learned from a high-volume intensive care unit. Anesthesia and analgesia; 2020.
26. Sarwar S, Waheed R, Sarwar S, Khan A. COVID-19 challenges to Pakistan: Is GIS analysis useful to draw solutions? Science of The Total Environment. 2020;139089.

© 2021 Zahid et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle4.com/review-history/66682>