



Hepatocellular Injury Ameliorated by a Common African Food, *Parkia biglobosa*

O. I. N. Onyekachi^a, S. F. Orji^b, C. N. Ugwu^c, C. Igwenyi^b, C. L. Uche^d,
I. O. Abali^e, M. U. Nwobodo^b, C. E. Iwuoha^f, N. M. Chika-Igwenyi^b,
C. A. Onyeaghala^g, F. U. Agu^h and A. I. Airaodion^{i*}

^a Department of Medical Microbiology, Ebonyi State University, Abakaliki, Nigeria.

^b Department of Internal Medicine, Alex Ekwueme Federal University Teaching Hospital, Abakaliki, Ebonyi State, Nigeria.

^c Department of Internal Medicine, Ebonyi State University, Abakaliki, Nigeria.

^d Department of Haematology, Abia State University, Uturu, Nigeria.

^e Department of Surgery, Abia State University, Uturu, Nigeria.

^f Department of Community Medicine, Abia State University, Uturu, Nigeria.

^g Department of Internal Medicine, University of Port-Harcourt Teaching Hospital, Rivers State, Nigeria.

^h Department of Physiology, Gregory University, Uturu, Abia State, Nigeria.

ⁱ Department of Biochemistry, Federal University of Technology, Owerri, Imo State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRIMPS/2022/v11i4199

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92175>

Original Research Article

Received 12 July 2022
Accepted 22 September 2022
Published 30 September 2022

ABSTRACT

Background: *Parkia biglobosa* seed has been reported to possess hepatoprotective potential. Therefore, this study sought to investigate its ability in ameliorating KBrO₃-induced hepatotoxicity. **Methodology:** *P. biglobosa* was extracted with soxhlet extractor with 95% ethanol as the solvent. Twenty-four adult male Wistar rats were acclimatized under laboratory conditions and were randomly grouped into A, B, C and D. Group A was given distilled water orally. Animals in groups B, C and D were administered 100 mg/kg body weight of potassium bromate, but groups C and D were also treated with 100 and 200 mg/kg body weight of *P. biglobosa* respectively. Both

*Corresponding author: E-mail: augustineairaodion@yahoo.com;

potassium bromate and *P. biglobosa* were freshly prepared on daily basis and administered to rats by oral gavage. After 28 days of treatment, the animals were sacrificed under mild diethyl ether anaesthetization 24 hours after cessation of last treatment. Blood and liver tissue were collected.

Results: The findings demonstrated that, when compared to the control group, KBrO₃ caused a significant increase ($P<0.05$) in ALT, AST, LDH, ALP, total bilirubin (TB), conjugated bilirubin (CB), and unconjugated bilirubin (UB) levels, but decreased total protein, albumin and globulin in the serum of animals. In the liver cells, KBrO₃ reduced hepatic biomarkers. These perturbations were neutralized in the groups treated with 100 and 200 mg/kg body weight, respectively.

Conclusion: The result of the present study revealed that KBrO₃ is hepatotoxic at a dose of 100 mg/kg body weight. The result further suggests that *P. biglobosa* possesses hepatoprotective properties in rats *in vivo*. This study can be replicated in human trial.

Keywords: Hepatoprotective potential; *Parkia biglobosa* seed; potassium bromate.

1. INTRODUCTION

Liver disease is a primary cause of death worldwide and a major global issue [1]. In the UK, the standardized mortality rate grew by more than 400% between 1970 and 2010 [2]. Infectious, carcinogenic, immune-mediated, and toxic substances can all cause harm to the liver because of its specific structure and functions [3]. Alterations in the concentrations of liver enzymes such alanine transaminase (ALT), aspartate transaminase (AST), and alkaline phosphatase (ALP) are early indicators of hepatic damage [4,5]. ALT is found in the cytoplasm of liver cells and in extrahepatic sources like the skeletal and cardiac muscles, although its quantity in the liver is significantly larger [3]. Although it is not unique to hepatocyte membrane damage, an increase in the serum level of ALT is highly predictive of extrahepatic injury [5]. In contrast, AST is present in the cytoplasm of all tissues other than the bones, albeit when compared to other tissues, it is most abundant in the liver and skeletal muscle [6]. Thus, hepatocellular injury can be distinguished from skeletal muscle injury because, in the former, the rise in ALT is greater than that in AST, resulting in a lower AST/ALT ratio than that which is possible in skeletal muscle injury [7]. On the other hand, high levels of bilirubin and low levels of glutathione indicate the death of hepatocytes [8,9], while alkaline phosphatase (ALP) is a sign of cholestasis [10].

The creation of bread, cheese, and beer all employ potassium bromate (KBrO₃), a common food ingredient. Due to its capacity to enrich dough and increase its volume and flexibility, it is frequently used in the production of bread [11]. The International Agency for Research on Cancer (IARC) categorized this molecule as a potential carcinogen in 1999, according to Mahmud et al. [12], yet despite this classification,

the compound is still used industrially in some nations (including Nigeria). The well-known toxin and cancer-causing substance is still utilized in the production of bread in different cities of Nigeria [13,14]. The harmful effect of KBrO₃ was shown in a study by Bayomy et al. [15] which was supported by elevated levels of malonaldehyde in the liver tissue, AST and ALT serum enzymes, as well as glutathione (GSH) and superoxide dismutase (SOD) in the liver tissues.

Parkia peas, which belong to the Fabaceae family (subfamily Mimisoideae), are highly prevalent in tropical regions like Africa. The *Parkia biglobosa*, one of the several species in this genus, is well recognized for its usage in cooking. The different tribes in Nigeria have distinct names for the plant known as African locust bean. It is known as "Ogili" among the Ibo, "Dorawa" among the Hausa, and "Iru" among the Yoruba [16,17]. The African locust bean seeds are cleaned, cooked till soft, washed and dehulled, and then allowed to ferment for a few days to produce this culinary product. After being shaped into balls, the fermented product is allowed to dry [18]. Analgesic, antihypertensive, antifungal, anti-inflammatory, cardioprotective, and hepatoprotective applications of African locust bean seed have all been described [17,19-25]. Therefore, the purpose of this study is to further assess how African locust bean protects against potassium bromate-induced hepatic enzyme distortion.

2. MATERIALS AND METHODS

2.1 Collection and Extraction of *Parkia biglobosa*

Having purchased *P. biglobosa* (African locust bean) seeds from a local market in Ibadan,

western Nigeria, the seeds were identified by a botanist. These were sun-dried and ground into powder using a mechanical blender (Moulinex). The extraction of the food product was done using a soxhlet apparatus, with 95% ethanol as the solvent; the steps outlined by Airaodion et al. [26,27] were followed. About 25 g of the sample powder in 250ml of ethanol in a round bottom flask was added to the soxhlet extractor and condenser on a heating mantle. Heat generated by the heating mantle was channeled to the solvent until it began to evaporate; the vapour was passed through the apparatus to the condenser. The condensate dropped into a reservoir that housed the sample-containing thimble. When the solvent level reached the siphon and was poured back into the flask with a flat bottom, the cycle was resumed. The operation was given a total of 18 hours. With a yield of 2.55 g and a percentage yield of 10.20 percent, the ethanol was evaporated in a rotary evaporator at 35 °C at the end of the process. The extract was kept in the refrigerator for further analysis.

2.2 Animal Treatment

Twenty-four (24) mature male Wistar rats (*Rattus norvegicus*) weighing between 140 and 160 g were used in the experiment. They were acclimated in a laboratory setting for seven (7) days prior to the trial. The rats were housed in wire-mesh cages with free access to commercial rat food and water. The animals were kept in standard temperature and humidity conditions with 12-hour alternating cycles of light and dark. This inquiry was carried out in accordance with the Declaration of Helsinki and the guidelines established by the Committee for the Purpose of Control and Supervision of Experiments on Animals. Additionally, NIH policy was followed when doing animal experiments [28]. The rats were picked at random and grouped into groups A, B, C, and D. Group A (control group) received oral distilled water; groups C and D received 100 and 200 mg/kg body weight of *P. biglobosa*, respectively, while the animals in group B received 100 mg/kg body weight of potassium bromate. The rats in groups B and C received 100mg/kg body weight of *P. biglobosa* in addition. Fresh potassium bromate and *P. biglobosa* were administered to rats every day by oral gavage. The animals had 28 days of successive treatments before being killed. The animals were sacrificed while being gently sedated with diethyl ether twenty-four hours following the last treatment. Through a heart puncture, blood was taken.

2.3 Preparation of Liver Homogenate

The animals were dissected, and the liver was taken out. After that, the liver of each rat were weighed. The liver tissue was homogenized in 50 mMol/L Tris-HCl buffer using a Teflon homogenizer (pH 7.4). The combination was centrifuged at 5000 rpm for 15 minutes, and the supernatants were stored in the freezer until they were required for biochemical analysis.

2.4 Determination of Hepatic Biomarkers

Alkaline phosphatase (ALP), aspartate aminotransferase (AST), alanine aminotransferase (ALT) and lactate dehydrogenase (LDH) activities were analysed from the liver and serum homogenates following the method of Reitman and Frankel; Randox commercial Enzyme kits were used [28]. Other biomarkers, namely Total Protein (TP), Albumin (ALB), Globulin (GLO), Total Bilirubin (TB), Conjugated Bilirubin (CB), and Unconjugated Bilirubin (UB), were also assayed from liver and serum homogenates following procedures previously described by Airaodion et al. [25].

2.5 Statistical Analysis

The results are presented as the mean minus the standard deviation. Turkey's test was used after one-way analysis of variance (ANOVA) to determine how homogeneous the groups were. Graph Pad Prism Software Version 8.00 was used for all analyses, and p values below 0.05 were regarded as statistically significant.

3. RESULTS

As shown in Table 1, the findings demonstrated that, when compared to the control group, potassium bromate caused a significant increase ($p < 0.05$) in the ALT, AST, LDH, ALP, total bilirubin (TB), conjugated bilirubin (CB), and unconjugated bilirubin (UB) levels in the serum of animals treated with only 100 mg/kg body weight of potassium bromate. The serum levels of these biomarkers in those rats treated with potassium bromate, as well as 100 and 200 mg/kg body weight doses of *P. biglobosa* extract, however, decreased significantly in a dose-dependent manner. The serum levels of total protein (TP), albumin (ALB), and globulin (GLO), on the other hand, decreased significantly in the group of rats treated with only 100 mg/kg body weight of potassium bromate, when compared with those in the control group (Table 1). An additional

Table 1. Effect of *P. biglobosa* on hepatic biomarkers in the serum of potassium Bromate-induced rats

| Hepatic biomarkers | Control | KBrO ₃ only | KBrO ₃ + 100 mg/kg <i>P. biglobosa</i> | KBrO ₃ + 200 mg/kg <i>P. biglobosa</i> | p-value |
|--------------------|-------------|------------------------|---|---|---------|
| AST (U/L) | 54.27±3.34 | 86.25±4.37 | 79.56±4.52 | 60.27±3.85 | 0.02 |
| ALT (U/L) | 151.53±9.53 | 211.28±8.93 | 198.47±6.36 | 173.96±4.83 | 0.04 |
| LDH (U/L) | 7.43±0.82 | 13.79±2.83 | 10.42±1.42 | 7.96±2.04 | 0.03 |
| ALP (U/L) | 73.45±6.47 | 102.00±5.93 | 94.02±2.27 | 79.22±4.83 | 0.02 |
| TP (g/dL) | 7.89±1.03 | 4.89±1.23 | 5.67±0.84 | 7.09±0.59 | 0.01 |
| ALB (g/dL) | 5.05±0.92 | 2.91±0.18 | 3.35±0.27 | 4.34±0.44 | 0.04 |
| GLO (g/dL) | 2.84±0.22 | 1.98±0.25 | 2.32±0.17 | 2.75±0.21 | 0.03 |
| TB (g/dL) | 0.09±0.00 | 0.22±0.01 | 0.19±0.01 | 0.11±0.00 | 0.02 |
| CB (g/dL) | 0.04±0.00 | 0.09±0.00 | 0.09±0.00 | 0.05±0.00 | 0.02 |
| UB (g/dL) | 0.05±0.00 | 0.13±0.01 | 0.10±0.00 | 0.06±0.00 | 0.02 |

Values are presented as Mean±SD, where n = 6

Legend: AST = Aspartate Aminotransferase, ALT = Alanine Aminotransferase, LDH = Lactate Dehydrogenase, ALP = Alkaline Phosphatase, TP = Total Protein, ALB = Albumin, GLO = Globulin, TB = Total Bilirubin, CB = Conjugated Bilirubin, UB = Unconjugated Bilirubin

Table 2. Effect of *P. biglobosa* on hepatic biomarkers in the liver of potassium Bromate-induced rats

| Hepatic biomarkers | Control | KBrO ₃ only | KBrO ₃ + 100 mg/kg <i>P. biglobosa</i> | KBrO ₃ + 200 mg/kg <i>P. biglobosa</i> | p-value |
|--------------------|-------------|------------------------|---|---|---------|
| AST (U/L) | 48.46±2.27 | 29.75 ±1.88 | 36.67 ±1.89 | 45.44±3.01 | 0.04 |
| ALT (U/L) | 128.19±4.29 | 97.45±5.55 | 108.33±3.10 | 122.84±5.25 | 0.02 |
| LDH (U/L) | 9.35±2.11 | 7.84±2.22 | 8.17±1.18 | 9.00±0.96 | 0.01 |
| ALP (U/L) | 78.85±3.29 | 81.27±3.55 | 81.84±4.18 | 79.79±3.21 | 2.96 |
| TP (g/dL) | 9.35±1.85 | 7.27±1.44 | 7.93±1.26 | 8.88±1.19 | 0.05 |
| ALB (g/dL) | 6.99±1.21 | 5.04±0.66 | 5.37±1.06 | 6.29±1.37 | 0.05 |
| GLO (g/dL) | 2.36±0.28 | 2.23±0.11 | 2.56±0.44 | 2.59±0.28 | 1.96 |
| TB (g/dL) | 0.34±0.01 | 0.29±0.03 | 0.29±0.02 | 0.32±0.02 | 1.74 |
| CB (g/dL) | 0.21±0.02 | 0.18±0.00 | 0.19±0.04 | 0.19±0.03 | 3.02 |
| UB (g/dL) | 0.13±0.00 | 0.11±0.00 | 0.10±0.00 | 0.13±0.00 | 5.67 |

Values are presented as Mean±SD, where n = 6

Legend: AST = Aspartate Aminotransferase, ALT = Alanine Aminotransferase, LDH = Lactate Dehydrogenase, ALP = Alkaline Phosphatase, TP = Total Protein, ALB = Albumin, GLO = Globulin, TB = Total Bilirubin, CB = Conjugated Bilirubin, UB = Unconjugated Bilirubin

treatment with 100mg and 200mg/kg body weight of locust bean seed extract resulted in a significant increase in the serum levels of TP, ALB, and GLO, in a dose-dependent manner.

Table 2 shows that the activities of AST, ALT, and LDH as well as the levels of TP and ALB in the liver cells of those rats given 100 mg/kg body weight dose of potassium bromate was significantly decreased (p<0.05), when compared to the rats in the control group. The additional administration of 100 mg/kg and 200 mg/kg body weight doses of *P. biglobosa* extract attenuated this decrease in a dose-dependent manner. The changes observed in the levels of the other biomarkers in the liver cells were of no statistical significance (Table 2).

4. DISCUSSION

One of the main characteristics of liver injury is changes in particular liver enzymes released into the blood when the liver cells deteriorate [29]. The serum levels of the aminotransferases rapidly rise when liver cells suffer ischemic injury, either from a lack of blood or oxygen [30]. With the exposure of one group of rats to KBrO₃ alone, the levels of AST, ALT, and ALP sharply increased, suggesting liver cell necrosis brought about by oxygen deprivation (Table 1). Similar results were obtained in a study by Hassan et al. [31] where rats given potassium bromate displayed elevated ALT, AST, and ALP activity. Numerous risk factors, such as potassium bromate, alcohol, environmental contaminants,

irradiation, and medications cause oxidative stress on liver cells [25,32]. The use of natural antioxidants found in medicinal plants has the power to scavenge free radicals created by oxidative stress, hence lessening liver damage caused by reactive oxygen species (ROS) [32]. One of these therapeutic plants with antioxidant properties is *Parkia biglobosa*, which is an everyday diet in Africa [16,33]. When the African locust bean seed extract was administered to the rats in the present research, the increase in serum aminotransferase activity brought about by the administration of potassium bromate was attenuated; the extent of attenuation depended on the dose of *Parkia biglobosa* extract administered. This supports the view that *Parkia biglobosa* seed extract has hepatoprotective characteristics as reported by Saleh et al. [16].

Although it is mostly a liver enzyme, alkaline phosphatase (ALP) can also be present in bone. Its activity may be increased in physiological circumstances, such as during pregnancy, but it is increased in pathological circumstances, such as hepatic congestion brought about by right-sided heart failure [34]. Additionally, certain medications including glucocorticoids and anti-convulsant, as well as high-fat diets, may cause it to rise [35]. The finding of Hassan et al. [31] is in agreement with the rise in ALP activity in the serum of Wistar rats fed with 100mg/Kg body weight of KBrO_3 in the index study.

The increased activity of indicators like ALT, AST, and LDH in the liver tissue of animals without pretreatment with *P. biglobosa* may be the result of hepatocyte cellular necrosis, which increases cell permeability. Alteration in the level of Lactate dehydrogenase is a measure of cell damage, including liver toxicity and artery endothelial instability [36]. The considerable rise in LDH activity seen in rats treated with KBrO_3 alone may be indicative of the onset of cytolysis, which is a potential marker of membrane damage, including damage to blood vessel endothelial membranes. Reactive oxygen species (ROS) are produced in endothelial cells as a result of this endothelial membrane rupture, whether directly or indirectly, as was previously observed [37]. Free radicals attack unsaturated fatty acids in the membranes, causing membrane lipid peroxidation that reduces membrane fluidity, enzyme leakage, loss of receptor activity, and damage to membrane proteins that result in cell inactivation [38]. Oxidative stress results from a reduction in the antioxidant defense system when lipid peroxidation increases [39]. This

shows that the treatment of KBrO_3 may have compromised the liver membrane of the rats, allowing for increased levels of AST, ALT, and LDH in the liver as well as subsequent penetration. However, when mice were given 100 and 200 mg/kg body weight of *P. biglobosa* seed extract, this impact was lessened. According to a recent report, *P. biglobosa* seed lowers the production of ROS in Wistar rats' hearts, livers, and kidneys [34]. So, the antioxidant activity of *P. biglobosa* seed may be the cause of its protective effects against KBrO_3 -induced hepatotoxicity.

In this investigation, KBrO_3 dramatically decreased the serum concentrations of total protein, albumin, and globin when compared to the control group. According to Ekam and Udosen's report [40], a drop in serum protein during hepatotoxicity simply denotes the presence of para-proteins or a drop in antibody synthesis. This decrease is consistent with their findings. The level of total protein, albumin, and globulin may fall as a result of liver dysfunction, malnutrition and malabsorption, diarrhea, nephrosis, alpha-1-antitripsin deficiency, acute hemolytic anemia, hypogammaglobulinemia/agammaglobulinemia, loss through the urine in severe kidney disease, pregnancy, and other conditions. Long-term hepatic cell death lowers serum levels of total protein, albumin and globulin, and generates additional hepatic releases to worsen hepatic dysfunction [40]. Other hemeoproteins include cytochromes, catalase, peroxidase, tryptophan pyrrolase, and a small pool of free heme.

Bilirubin is the breakdown result of the heme moiety of hemeoglobin [41]. Hyperbilirubinaemia is caused by an increase in the amount of directly reacted bilirubin in the blood. This syndrome is toxic and can lead to jaundice, auditory impairment, and neurotoxicity with subsequent brain damage [41]. On the other hand, modest unconjugated hyperbilirubinaemia functions as a mild antioxidant and may provide defense against the development of tumors and cardiovascular diseases [42]. Low concentrations of directly reacting bilirubin can sometimes lead to cardiac issues as well as strokes in humans [42]. Under a variety of clinical circumstances, serum bilirubin levels are frequently elevated. The fact that bilirubin is linked to serum albumin throughout blood circulation lessens the possibility of toxicity considered to be brought on by free bilirubin [43]. Bilirubin binds to albumin with a high affinity, but is quickly and selectively

taken up by the liver, conjugated with glucuronate, and secreted into bile [42]. As a result, bilirubin is transformed in the liver into bilirubin glucuronic acid and eliminated along with bile. Comparing the KBrO₃-treated rats to the control animals revealed a statistically significant increase in the serum levels of total, conjugated, and unconjugated (indirect) bilirubin, which may indicate tissue injuries. However, in animals that received both KBrO₃ and *P. biglobosa* seed extract at the same time, this effect was reduced.

Folk medicine has long recognized the wide range of therapeutic uses of *P. biglobosa* seed, and scientific development has produced ample proof for the majority of these claims. The hepatoprotective potential of this plant has been further proven by the current *in vivo* study. Based on a significant increase (P<0.05) in hepatic biomarkers in the KBrO₃ group when compared to the control group, it was determined in this study that KBrO₃ intoxication in rats caused hepatocellular damage. Concurrent administration of *P. biglobosa* seed extract to rats alleviated KBrO₃-induced changes in rat serum hepatic biomarkers in a dose-dependent manner, indicating that this plant may have hepatoprotective properties. Our findings seem to support an earlier finding by Airaodion et al. [25] that *Parkia biglobosa* protects the liver from acute ethanol-induced oxidative stress in Wistar rats.

In comparison to the control group, Table 2 demonstrates that the administration of KBrO₃ decreased the activities of AST, ALT, LDH, and ALP as well as the concentrations of total protein, albumin, globulin, total bilirubin, conjugated, and unconjugated bilirubin in the liver. Since the liver is where these biomarkers are primarily produced, the findings of this study may suggest that KBrO₃ decreased either the synthesis of these parameters or increased their efflux from the liver thereby increasing their level in the blood. As a result of the *P. biglobosa* seed extract treatment, these perturbations were attenuated. The effect of *P. biglobosa* seed in this study could be due to its rich phytochemical composition and antioxidant potential [34].

5. CONCLUSION

The result of the present study revealed that KBrO₃ is hepatotoxic at a dose of 100 mg/kg body weight. The result further suggests that

P. biglobosa possesses hepatoprotective properties in rats *in vivo*. This study can be replicated in human trial.

CONSENT

It is not applicable.

ETHICAL APPROVAL

Animal Ethic committee approval has been taken to carry out this study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Wang H. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: A systematic analysis for the Global Burden of Disease Study. *Lancet*. 2016;388:1459–1544.
2. Williams R, Aspinall R, Bellis M. Addressing liver disease in the UK: A blueprint for attaining excellence in healthcare and reducing premature mortality from lifestyle issues of excess consumption of alcohol, obesity, and viral hepatitis. *Lancet*. 2014;384(9958):1953-1997.
3. Gwaltney-Brant SM. Nutraceuticals in hepatic diseases. In RC Gupta, *Nutraceuticals Efficacy, Safety, and Toxicity*. Academic Press. 2016;87-99. Available: <https://doi.org/10.1016/B978-0-12-802147-7.00007-3>
4. Kwo PY, Cohen SM, Lim JK. ACG Clinical Guideline: Evaluation of abnormal liver chemistries. *American Journal of Gastroenterology*. 2017;112:18–35.
5. Sookoian S, Pirola CJ. Liver enzymes, metabolomics and genomewide association studies: From systems biology to the personalized medicine. *World Journal of Gastroenterology*. 2015;21(3): 711-725.
6. Washington IM, Van Hoosier G. Clinical Biochemistry and Haematology. In MA Suckow, KA Stevens, RP Wilson (Eds) *The laboratory rabbit, guinea pig, hamster, and other rodents*, American College of

- Laboratory Animal Medicine. Academic Press. 2012;57-116.
Available:<https://doi.org/10.1016/B978-0-12-380920-9.00003-1>.
7. Aulbach AD, Amuzie CJ. Biomarkers in nonclinical drug development. In A comprehensive guide to toxicology in nonclinical drug development. Academic Press. 2017;447-471.
Available:<https://doi.org/10.1016/B978-0-12-803620-4.00017-7>
 8. Feldstein AE, Gores GJ. Apoptosis in alcoholic and nonalcoholic steatohepatitis. *Front Biosci.* 2005;10:3093–3099.
 9. Malhi H, Gores GJ. Cellular and molecular mechanisms of liver injury. *Gastroenterology.* 2008;134:1641–1654.
 10. Chen VL, Du X, Chen Y, Kuppa A, Handelman SK, Vohnoutka RB, Peyser PA, Palmer ND, Bielak LF, Halligan B, Speliotes EK. Genome-wide association study of serum liver enzymes implicates diverse metabolic and liver pathology. *Nature Communication.* 2021;12:816.
 11. Ahmad MK, Mahmood R. Protective effect of Taurine against Potassium Bromate-induced haemoglobin oxidation, oxidative stress, and impairment of antioxidant defense system in blood. *Environmental Toxicology.* 2014;31(3):304-313.
 12. Mahmud SS, Moni M, Imran AB, Foyez T. Analysis of the suspected cancer-causing Potassium Bromate additive in bread samples available on the market in and around Dhaka city in Bangladesh. *Food Sci Nutr.* 2021;9:3752-3757.
 13. Airaodion AI, Ewa O, Ogbuagu EO, Ogbuagu U, Agunbiade AP, Oloruntoba AP. Evaluation of potassium bromate in bread in Ibadan metropolis: Fifteen years after ban in Nigeria. *Asian Food Science Journal.* 2019;7(4):1–7.
 14. Airaodion AI, Awosanya OO, Ogbuagu EO, Ogbuagu U, Njoku OC, Esonu C, Airaodion EO. Assessment of bread in ogbomoso metropolis for the presence of potassium bromate. *Asian Journal of Research in Biochemistry.* 2019;4(2):1-6.
 15. Bayomy NA, Soliman GM, Abdelaziz EZ. Effect of Potassium Bromate on the liver of adult male albino rats, and a possible protective role of vitamin C: Histological, immunohistochemical, and biochemical study. *The Anatomical Record.* 2016;299(9):1256-1269.
 16. Saleh, MSM, Jalil J, Zainalabidin S, Asmadi AY, Mustafa NH, Kamisah Y. Genus *Parkia*: Phytochemical, medicinal uses, and pharmacological properties. *Int. J. Mol. Sci.* 2021;22:618.
 17. Ezirim EO, Uche CL, Abali IO, Iwuoha CE, Chika-Igwenyi NM, Onyeaghala CA, Orji SF, Ugwu CN, Ugwu NI, Igwenyi C, Airaodion AI. Therapeutic potential of *Parkia biglobosa* seed against Potassium Bromate-induced testicular toxicity. *International Journal of Research and Reports in Gynaecology.* 2022;5(3):78-89.
 18. Samyal S. Recent trends in alkaline fermented foods. In J. Singh, A. Vyas (Eds), *Advances in Dairy Microbial Products.* Woodhead Publishing. 2022;57-79.
Available:<https://doi.org/10.1016/B978-0-323-85793-2.00001-1>
 19. Okoye TC, Uzor PF, Onyeto CA, Okereke EK. Safe African medicinal plants for clinical studies. In V. Kuete (Ed), *Toxicological Survey of African Medicinal Plants.* Elsevier. 2014;535-555.
Available:<https://doi.org/10.1016/B978-0-12-800018-2.00018-2>.
 20. Yahaya U, Abubakar S, Salisu A. Antifungal activity of *Parkia biglobosa* extract on pathogenic *Candida albicans*. *Journal of Applied Sciences.* 2019;19:235-240.
 21. Ouedraogo N, Atchade C, Traore TK, Ouedraogo B, Laurent B, Kabre E, Yoda J, Kini FB, Lompo M, Ouedraogo M, Ouedraogo S. Anti-inflammatory activity of extracts from *Parkia biglobosa* (Jacq.) R.Br. Ex G.Don. (Fabaceae-Mimosoideae) Trunk bark. *Journal of Pharmacology and Toxicology.* 2021;16:1-8.
 22. Kouadio F, Kanko C, Juge M, Grimaud N, Jean A, N'Guessan YT, Petit JY. Analgesic and anti-inflammatory activities of an extract from *Parkia biglobosa* used in traditional medicine in the Ivory Coast. *Phytother Res.* 2000;14(8):635-7.
 23. Airaodion AI, Ogbuagu EO. Ameliorative effect of *Parkia biglobosa* (African Locust Bean) against Egg-yolk induced hypertension. *International Journal of Bio-Science and Bio-Technology.* 2020;12(5):2233-7849.
 24. Adi K, Metowogo K, Mouzou A, Lawson-Evi P, Eklu-Gadegbeku K, Agbonon A, Lamboni C, Essien K, Aklikokou K, Gbeassor M. Evaluation of cardioprotective effects of *Parkia biglobosa* (Jacq. Benth) Mimosaceae stem bark. *Journal of Applied Pharmaceutical Science.* 2013;3(2):60-64.

25. Airaodion AI, Ogbuagu EO, Ogbuagu U, Adeniji A, Agunbiade AP, Airaodion EO. Hepatoprotective effect of *Parkia biglobosa* on acute Ethanol-induced oxidative stress in Wistar rats. International Research Journal of Gastroenterology and Hepatology. 2019;2(1):1-11.
26. Airaodion AI, Ngwogu AC, Ekenjoku JA, Ngwogu KO. Hepatoprotective potency of ethanolic extract of *Garcinia kola* (Heckel) seed against acute ethanol-induced oxidative stress in Wistar rats. International Research Journal of Gastroenterology and Hepatology. 2020;3(2):1-10.
27. Airaodion AI, Ogbuagu EO, Ekenjoku JA, Ogbuagu U, Airaodion EO. Therapeutic effect of methanolic extract of *Telfairia occidentalis* leaves against acute ethanol-induced oxidative stress in Wistar rats. International Journal of Bio-Science and Bio-Technology. 2019;11(7):179-189.
28. National Research Council. United States Committee for the update of the Guide for the Care and Use of Laboratory Animals. Guide for the Care and Use of Laboratory Animals. 8th Ed., Washington (DC): National Academies Press (US); 2011. PMID: 21595115.
29. Reitman S, Frankel S. A colorimetric method for determination of serum glutamate oxaloacetate and glutamic pyruvate transaminase. American Journal of Clinical Pathology. 1957;28:56-58.
30. Contreras-Zentella ML, Hernández-Muñoz R. Is liver enzyme release really associated with cell necrosis induced by oxidant stress? Oxidative Medicine and Cellular Longevity. 2016;12:Article ID 3529149. Available:<https://dx.doi.org/10.1155/2016/3529149>.
31. Ghosh A, Onsager C, Mason A, Arriola L, Lee W, Mubayi A. The role of oxygen intake and liver enzyme on the dynamics of damaged hepatocytes: Implications to ischaemic liver injury via a mathematical model. PLoS ONE. 2021;16(4):e0230833.
32. Hassan I, Ebaid H, Alhazza IM, Al-Tamimi J. The alleviative effect of vitamin B2 on potassium bromated-induced hepatotoxicity in male rats. BioMed Res Int. 2017;9(5):23;31. Article ID 8274261. Available:<https://doi.org/10.1155/2020/8274261>.
33. Li S, Tan H, Wang N, Zhang Z, Lao L, Wang C, Feng Y. The role of oxidative stress and antioxidants in liver diseases. International Journal of Molecular Science. 2015;16:26087-26124.
34. Ugwu CN, Abali IO, Iwuoha CE, Chika-Igwenyi NM, Onyeaghala CA, Orji SF, Igwenyi C, Uche CL, Onyekachi OIN, Nwobodo MU, Airaodion AI. Ameliorative effect of *Parkia biglobosa* (African locust bean) seed against potassium bromate-induced oxidative stress. Merit Research Journal of Medicine and Medical Sciences. 2022;10(8):213-219.
35. Fernandez NJ, Kidney BA. Alkaline phosphatase beyond the liver. Vet Clin Pathol. 2007;36(3):223-33.
36. Airaodion AI, Ogbuagu EO, Ewa O, Ogbuagu U, Awosanya OO, Adekale OA. Ameliorative efficacy of methanolic extract of *Corchorus olitorius* leaves against Acute ethanol-induced oxidative stress in Wistar rats. Asian Journal of Biochemistry, Genetics and Molecular Biology. 2019; 7(6):1-9.
37. Airaodion AI, Akinmolayan JD, Ogbuagu EO, Esonu CE, Ogbuagu U. Preventive and therapeutic activities of methanolic extract of *Talinum triangulare* leaves against ethanol-induced oxidative stress in Wistar rats. International Journal of Bio-Science and Bio-Technology. 2019;11(7):85-96.
38. Airaodion AI, Ogbuagu U, Ekenjoku JA, Ogbuagu EO, Airaodion EO, Okoroukwu VN. Hepato-protective efficiency of ethanol leaf extract of *Moringa oleifera* against hydrocarbon exposure. International Journal of advances in Herbal and Alternative Medicine. 2019;03(01):32-41.
39. Airaodion AI, Ogbuagu EO, Ekenjoku JA, Ogbuagu U, Airaodion EO. Therapeutic effect of methanolic extract of *Telfairia occidentalis* leaves against acute ethanol-induced oxidative stress in Wistar rats. International Journal of Bio-Science and Bio-Technology. 2019;11(7):179-189.
40. Ekam VS, Udosen EO. Total protein, albumin and globulin levels following the administration of activity directed fractions of *Vernonia amygdalina* during acetaminophen induced hepatotoxicity in Wistar albino rats. Global Journal of Pure and Applied Sciences. 2012;18(1&2):25-29.
41. Airaodion AI, Ogbuagu EO. Effect of *Cyperus esculentus* L. (tiger nut) milk on hepatic and renal indices of Wistar rat.

- Asian Journal of Research in Nephrology. 2020;3(2):10-16.
42. Airaodion AI, Ene AC, Ogbuagu EO, Okoroukwu VN, Ekenjoku JA, Ogbuagu U. Biochemical changes associated with consumption (by rats) of “garri” processed by traditional and instant mechanical methods. Asian Journal of Biochemistry, Genetics and Molecular Biology. 2019;2(4):1-11.
43. Airaodion AI, Megwas AU, Ekenjoku JA, Ngwogu KO, Ngwogu AC. Nephro- and hepato-toxicity of common household insecticides used in Nigeria. International Research Journal of Gastroenterology and Hepatology. 2020;3(2):21-28.

© 2022 Onyekachi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/92175>