



Indirect Ophthalmoscopy: An Abandoned Workmanship

Hussain Ahmed Khaqan ^{a \circ} , Sabah Eric ^{b* $\#$} , Ateeq ur Rehman ^{a $\#$} ,
Hasnain Muhammad Buksh ^{a \dagger} , Muhammad Ali Haider ^{a \ddagger} and Aamna Jabran ^{a \yen}

^a Department of Ophthalmology, Post graduate Medical Institute, Ameer-ud-din Medical College, Lahore General Hospital, Lahore, Pakistan.

^b Government Dispensary Muhajar Goth, Mirpurkhas, Sindh Institute of Ophthalmology and Visual Sciences, Hyderabad, Pakistan.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/88405>

Short Research Article

Received 17 April 2022
Accepted 26 June 2022
Published 04 July 2022

ABSTRACT

Purpose: We sought to evaluate the application of indirect ophthalmoscopy in the routine practice of retinal surgeons.

Methods: A self-administered questionnaire was circulated among various ophthalmologists. Potential participants were vitreoretinal surgeons and those ophthalmologists who work in a vitreo retinal department in Asia, The United Kingdom and USA. The respondents were asked to mark either "YES" or "NO" for the given questions through google survey form.

Results: Responses were obtained from 68 ophthalmologists in total out of which 57(83.82%) were retinal surgeons and 11(16.18%) were not. 66 (97.06%) admitted using an indirect ophthalmoscope in their routine practice whereas, only 2 (2.94%) were not using it routinely. 43(63.24%) considered it to be an essential tool for examining retinal patients and 25(36.76%) do not consider it a compulsory part of the retinal examination.

^{\circ} Professor Ophthalmology,

^{$\#$} Consultant Ophthalmologist,

^{\dagger} Senior Registrar,

^{\ddagger} Assistant Professor,

^{\yen} Senior Registrar,

*Corresponding author: E-mail: teamkhaqan@gmail.com, drsabaheric@gmail.com;

Conclusion: Unfortunately, the majority of ophthalmologists seem to be naive to the imperative requirement of indirect ophthalmoscopy in order to examine vitreo retinal patients, notwithstanding the certitude that this skill has been considered mandatory in routine clinical practice of ophthalmologists.

Keywords: Indirect Ophthalmoscope; Vitreo retinal surgeons; ophthalmology; retinal detachment; fundus.

1. INTRODUCTION

The ophthalmoscopic examination is a significant clinical dexterity that permits ophthalmologists, peculiarly vitreo retinal surgeons to analyze, screen and diagnose numerous sight-threatening retinal pathologies. However, despite the necessity and ascendancy of indirect ophthalmoscopes this art seems to be not fully mastered by many [1].

In this era, it is observed that the indirect ophthalmoscope is an under-utilized tool in the clinical setting and that the majority of ophthalmologists lacked confidence in its use.

This prompts the loss of a significant demonstrative help that depends on a compact and easy to grasp-instrument [2].

The binocular indirect ophthalmoscope is an optical instrument [3] placed on the examiner's head, and is occasionally fixed to spectacles. It is used to inspect the posterior segment of the eye and produces a stereoscopic image with between 2x and 5x magnification. It is very beneficial for the diagnosis and treatment of retinal tears, holes, and detachments particularly when the periphery is intended to be observed. Fully dilated pupils are a prerequisite [4,5].

The examiner orientates his/her head to direct the light from the internal light source into the patient's eye in a dark room. A plus-powered condensing lens is held by the examiner at its focal length from the patient's eye for two reasons; 1.The lens 'condenses' light from the illumination system towards the patient's pupil. 2. Light reflected from the retina passes back through the lens creating a real, horizontally and laterally inverted image of the fundus situated between the lens and the examiner [6,7].

The viewing system of the indirect ophthalmoscope consists of a pair of low-

powered convex lenses. This design renders the examiner a stereoscopic view of the virtual image. A +20D lens is the standard lens for general examination contributing to 3x magnification and a field of view of approximately 45°. A +30D lens will offer 2xmagnification alongside a field of roughly 65° which is ordinarily used to examine the pediatric population and those with small pupils.

It is important for ophthalmologists and vitreo retinal surgeons to master the art of indirect ophthalmoscopy and make this a mandatory practice to avoid any unforeseen complications [8].

The aim of this survey was to evaluate how much importance is given to indirect fundoscopy by the majority of retinal surgeons and those ophthalmologists who are working in a vitreo retinal setting.

2. METHODS AND MATERIALS

A questionnaire was formulated in which the practicing ophthalmologists were asked 6 questions through Google Survey. The survey included particularly retinal surgeons from the UK, USA and Asia.

They were asked that how often they use an indirect ophthalmoscope in their clinical routine practice, how many retinal detachment patients are examined by indirect ophthalmoscope, is the indirect ophthalmoscopy done with or without indentation and how much importance does the indirect ophthalmoscopy holds in a retinal surgeon's life?

68 respondents took part in this survey and they were not asked to identify themselves by mentioning their names but the information was implored on their current position i.e; general ophthalmologists or vitreo retinal surgeons.

Table 1. Demographic study

Questions	Responses obtained from 68 ophthalmologists				
	100%	75%	50%	25%	0%
For how much of your retina patients do you perform indirect ophthalmoscopy	10(14.71%)	7(10.29%)	15(22.06%)	33(48.53%)	3(4.41%)
For how much % of your retinal detachment patients do you perform indirect ophthalmoscopy	13(18.84%)	5(7.25%)	8(11.59%)	20(40.58%)	15(21.74%)
how many % of your retina patients undergo indirect ophthalmoscopy with indentation	4(5.88%)	1(1.47%)	14(20.59%)	27(39.71%)	22(32.35%)
Do you use indirect ophthalmoscope in your routine practice	YES 66 (97.06%)		NO 2 (2.94%)		
Do you consider indirect ophthalmoscopy to be an essential tool for examining retinal patients	43(63.24%)		25(36.76%)		

Questionnaires were inspected with the response of all ophthalmologists grouped together. The degree of internal consistency within the whole group of experts was measured by Kendall's coefficient of concordance (W). All descriptive data were analyzed with a statistical software package (SPSS, Version 16.0 for Windows).

3. RESULTS AND DISCUSSION

Of the 80 Google survey forms distributed online, a total of 70 were returned (i.e. 87.5% response rate), however in 2 of these, the respondents had not marked one question. 68 correctly completed survey forms were therefore available for analysis.

Responses were obtained from 68 ophthalmologists in total out of which 57(83.82%) were retinal surgeons and 11(16.18%) were not.

Retinal vasculature has the quality to be viewed without invasive techniques and surprisingly, minor retinal microvascular abnormalities can be identified on indirect ophthalmoscopy which can be sight saving as well as life saving ; putting something aside for some patients.. In today's clinical practice barely any clinicians perform ophthalmoscopy, and numerous who do can't dependably recognize anomalies of the fundus [9-12]. Given the unfortunate capability of most doctors with ophthalmoscopy, the eventual fate

of the assessment of the visual fundus is unsure and presents an enormous clinical challenge.

4. CONCLUSION

The most significant results show that only 25% of retina patients are examined under indirect ophthalmoscope with or without indentation. This survey highlights that indirect ophthalmoscopy is an abandoned workmanship of the modern era [13] [14]. Notwithstanding the revolution in the field of ophthalmology, ophthalmologists should still rely primarily on using an indirect ophthalmoscope to assess the prognostic signs in a retina that may lead to retinal detachment and other sight-threatening complications.

Yet, clinicians still appear to avoid the indirect ophthalmoscope in practice, putting patient safety at risk.

CONSENT

As per international standard or university standard, respondents' written consent has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ophthalmoscopy simulation: Advances in training and practice for medical students and young ophthalmologists
2. Retinal drawing Valappil S, Jayan A - Kerala J Ophthalmol
3. Indirect Ophthalmoscopy 101 - American Academy of Ophthalmology
4. Available:https://journals.lww.com/internationalophthalmology/Citation/1978/01820/The_Importance_of_Indirect_Ophthalmoscopy.12.aspx
5. the Relative Importance of Direct and Indirect Ophthalmoscopic Examination In the Treatment OF Retinal Detachment - PMC
6. Understanding and caring for an indirect ophthalmoscope - PMC
7. Wide-Angle Viewing System versus Conventional Indirect Ophthalmoscopy for Scleral Buckling - PMC
8. A questionnaire-based assessment of Safe Eye Examination (SEE) technique - PMC
9. Available:<https://www.thepermanentjournal.org/issues/2015/fall/5944-book-review.htm>
10. IOn the Evolution of Binocular Ophthalmoscopy | Medical Devices and Equipment | JAMA Ophthalmology
11. The clinical relevance of visualising the peripheral retina – Science Direct
12. Questionnaire-based survey on the importance of quality of life measures in ophthalmic practice | Eye
13. The demise of direct ophthalmoscopy - PMC
14. Nonmydriatic Ocular Fundus Photography in the Emergency Department - PMC

© 2022 Khaqan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/88405>*