



## **Distribution of ABO and Rhesus Blood Groups among COVID-19 Positive Subjects in Rivers State, Nigeria**

**Onengiye Davies<sup>a\*</sup>, Serekara G. CHRISTIAN<sup>a</sup> and Stella U. Ken-Ezihuo<sup>a</sup>**

<sup>a</sup> *Department of Medical Laboratory Science, Rivers State University, Port Harcourt, Nigeria.*

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

#### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/90321>

**Original Research Article**

**Received 12 June 2022**  
**Accepted 06 August 2022**  
**Published 13 August 2022**

### **ABSTRACT**

Coronavirus-2019 (COVID-19) is a novel disease that is ravaging society today. It has been reported that individuals with some blood group antigens are more predisposed or susceptible to certain diseases. This study was carried out to determine the distribution of some blood group antigens among COVID-19-infected individuals. The study involved one hundred (100) subjects, comprising 75 males and 25 females, within the ages of 20 and 60 years, who were infected with COVID-19, and receiving treatment at the COVID-19 Isolation Centre. The bio-data of the subjects were obtained using a well-structured questionnaire. Five millilitres (5 ml) of blood collected aseptically from each of the subjects who gave informed consent were dispensed into K<sub>3</sub>EDTA bottles for the determination of the ABO and Rh blood groups. The bio-data show that 74% of the subjects were married while 26% were single. Also, 87% of the subjects had university degree or HND, 6% had OND while 7% had only secondary school education, indicating that majority of the subjects were educated. The results from the study showed that the distribution of ABO blood group among the subjects were 75% blood group O, 11% blood group A, 10% blood group B and 4% blood group AB. Also, 91% of the subjects were Rh positive while 9% were Rh negative. The results indicate that the blood group antigens found in the subjects were similar to the pattern reported in the population where the research was conducted. It is concluded that the prevalent blood group in this study was blood group O, while the most prevalent Rh group was Rh-D positive. It is recommended that blood groups be considered in the testing menus for the

\*Corresponding author: E-mail: [nengidavies25@gmail.com](mailto:nengidavies25@gmail.com);

management of COVID-19 infection, since some subjects may have such blood group antigens that can predispose to the infection.

*Keywords: Coronavirus; COVID-19; blood groups; ABO; rhesus; Port Harcourt.*

## 1. INTRODUCTION

Blood group systems can be defined as one or more antigens that are controlled at a single gene locus, or by two or more very closely linked homologous genes with little or no observable recombination between them [1]. In other words, a blood group system refers to a group of antigens that are encoded by alleles at a single gene locus or by a number of gene loci that are so closely linked that crossing over between them is very rare or does not occur [2]. These blood group antigens are either sugars or proteins that are attached to various components in the red blood cell membrane. For example, the antigens of the ABO blood group are sugars, while those of the Rhesus blood group are proteins [3].

The ABO blood group system, which was discovered by Landsteiner in 1900, was the first blood group ever to be discovered and still remains the most clinically significant of all the blood group systems [4]. Landsteiner classified the blood groups based on the antigens, A and B, that he identified on the surface of the blood of his subjects [5], and the expression of these antigens is determined by the ABO gene located on chromosome 9 (9q34.2) [6].

The Rh blood group system is the most complex, and perhaps the second most studied blood group system [7]. There are over 50 antigens in the human Rh blood group system but the principal Rhesus antigens of medical importance are D, C, E, c and e [8]. A person with Rh antigen is referred to as Rhesus positive while individuals lacking the antigen are Rhesus negative [9]. The Rh antigen was so named because a related antigen was first discovered in Rhesus monkeys [10].

Inheritance of Rh antigens is determined by a complex of two closely linked genes, namely RHD and RHCE, which are found on the short arm of chromosome 1(1p36.11) and positioned in opposite directions to each other [11]. One of the genes encodes the protein carrying D antigen (Rh D); the other encodes the protein carrying C or c and E or e antigen (Rh CE). In the Rh system different antigen combinations (or

haplotypes) are observed, with the following as the most common: Dce (Ro, Rho), DCe (R1, Rh1), DcE (R2, Rh2), DCE (Rz, Rz), ce (r,rh), cE (r", hr") [12].

The Coronaviridae is a family of enveloped, single-stranded, positive-sense RNA viruses with the largest genome among RNA viruses. The club-shaped spike (S) proteins on their surface make them look like a crown. Other structural proteins include the haemagglutinin esterase (HE) (only found in some of them), small membrane (E), membrane (M), nucleocapsid (N) and internal (I) protein [13]. The most common symptoms at the start of COVID-19 are fever, cough, and fatigue. Other symptoms include sputum production, headache, rhinorrhoea, sneezing, sore throat, haemoptysis, diarrhoea, dyspnoea, and lymphopenia, with an incubation period of about 5 days, and a time frame between the onset of COVID-19 symptoms and death is about 6 to 41 days [14].

The presence or absence of the red blood cell antigens has been implicated in the susceptibility and pathogenesis of some bacterial, viral and parasitic infections, such as *Vibrio cholera*, *Escherichia coli* [15], *Helicobacter pylori* [16], Norovirus [17], Rotavirus and Human Immunodeficiency Virus (HIV) [18].

This study seeks to determine the distribution of ABO and Rh blood group antigens among COVID-19 positive subjects, a piece of information that is scanty in our area of study.

## 2. METHODOLOGY

### 2.1 Study Design

A cross-sectional study design was used for this study, comprising subjects who were positive for COVID-19, and were on admission and isolation at the Rivers State COVID-19 Isolation Centre, Eleme Local Government Area, Rivers State, Nigeria, between December, 2021 and March, 2022.

### 2.2 Study Area

This study was carried out in the Rivers State COVID-19 Isolation Centre in Eleme local

government area (LGA), Rivers State, Nigeria. The facility was the only facility used in the testing and management of COVID-19 in the State.

Eleme Local Government Area is one of the twenty-three Local Government Areas Rivers State, South-South, Nigeria, with a population of 190,884 as at the 2006 National Census, and is located between longitude 7° and 7°15" East of the Meridian and latitudes 4°60" and 4°35" North of the Equator. The total area of Eleme is about 138 square kilometers (km<sup>2</sup>), and is bordered on the North by Obio/Akpor Local Government Area and Oyigbo Local Government Area, on the East by Tai Local Government Area, on the South by Ogu/Bolo Local Government Area and Okrika Local Government Area [19].

### 2.3 Study Population

This study involved a total of 100 subjects, comprising 75 male and 25 female subjects, who were between the ages of 20 and 60 years and who were tested and were confirmed positive for COVID 19. Convenient sample size was used to determine the sample size for this study,

because COVID-19 is a novel disease that is still infecting several people globally, and the prevalence in the study area was not yet established at the time of this study.

The subjects gave their informed consent before being recruited for this study. The demographic data of the subjects were obtained using a structured questionnaire.

### 2.4 Eligibility Criteria

#### 2.4.1 Inclusion criteria

Subjects within the age bracket of 20 to 60 years, who tested positive for COVID-19, confirmed for the infection, and admitted in the isolation center were used for this study.

#### 2.4.2 Exclusion criteria

Subjects who were unconscious or those who were experiencing severe difficulty in breathing as a result of COVID-19 were excluded from the study (because they were not able to give informed consent), as well as those who did not give their consent to participate in the study.




Fig. 1. Location of Eleme local government area [19]

## 2.5 Sample Collection

Five millilitres (5mls) of whole blood sample were collected from each subject by venipuncture technique and dispensed into ethylene diamine tetraacetic acid-anticoagulant bottles; the blood was properly mixed with the anticoagulant by several gentle inversions, which was followed by the analysis of ABO and Rh blood groups.

## 2.6 Sample Analysis

The Determination of ABO and RhD Blood Groups was done using PRO-MED for AB (Lot Number B 191007 and Lorne Laboratories Ltd, Lot Number A200101; Expiry Date: 09/2021).

**Method:** The slide method was used according to manufacturer's instruction.

**Principle:** The ABO and Rh blood grouping system is based on agglutination reaction. When red blood cells carrying one or both antigens are exposed to the corresponding antibodies, they interact with each other to form visible agglutination or clumping.

**Procedure:** A clean and dry glass slide was divided into four sections with a glass marking pencil. The sections were labelled as Anti-A, Anti-B and Anti-D to identify the antisera. Then, one drop of anti-A serum, one drop of anti-B serum, and one drop of anti-D serum were placed in the center of the corresponding sections of the slide. One drop of the blood sample to be tested was added to each antisera. Each mixture (antiserum and blood) was mixed together with the aid of separate sticks, and the contents were tilted and examined for agglutination after two minutes.

**Result Interpretation:** Positive (+) result: Little clumps of red cells were seen floating in a clear liquid, indicating a positive result.

Negative (-) result: Red cells were floating homogeneously in a uniform suspension, indicating a negative result.

Confirmation of Rh D negative results: The blood sample was used to prepare 5% red cell suspension. One drop of the anti-D antiserum was dispensed into a clean test tube, and then one drop of the cell suspension was added to it. The mixture was incubated at 37°C in a water bath for 30 minutes. Then the mixture was centrifuged, and then, using a Pasteur pipette,

the mixture was placed on a clean slide and viewed under the microscope using X10 objective. There was no agglutination, confirming the Rh D negative status.

## 2.7 Data Analysis

The data generated from the study were analysed using Statistical Package for Social Sciences (SPSS) (IBM SPSS Statistics version 23), and were expressed in tables. A comparison of the proportion of subjects with the different blood groups was done using Pearson's Chi-Square and p-values less than 0.05 were considered statistically significant.

## 3. RESULTS

The number of male subjects in this study was 75, while that of female was 25, giving a total of 100. Seventy-four of these subjects were married while 26 were single. Eighty-seven of these subjects had a university degree or HND from a polytechnic, 6 had OND while 7 had O/Level (SSCE).

**Table 1. Demographic Characteristics of Subjects**

Variable	Frequency (%)
Gender	
Male	75 (75)
Female	25 (25)
Marital Status	
Married	74 (74)
Single	26 (26)
Educational Qualification	
Degree/HND	87 (87)
OND/NCE	6 (6)
SSCE	7 (7)

**Table 2. Percentage Distribution of COVID-19 Positive Subjects Based on ABO Blood Groups**

Blood Group	Number of Subjects	Percentage (%)
A	11	11
B	10	10
AB	4	4
O	75	75
Total	100	100

$\chi^2$ -value= 134.480, p-value= <0.001

Table 2 describes the distribution of subjects according to ABO blood groups. Eleven (11) subjects had blood group A, 10 had blood group B, 4 had blood group AB while 75 had blood group O.

**Table 3. Percentage Distribution of COVID-19 Positive Subjects Based on Rh Blood Group**

From the analysis, ninety-one (91) subjects were Rh D positive while 9 subjects were Rh D negative.

Blood Group	Number of Subjects	Percentage (%)
Rh D Positive	91	91
Rh D Negative	9	9
Total	100	100

$\chi^2$ -value= 67.240, p-value= <0.001

**Table 4. Distribution of ABO Blood Groups of COVID-19 Positive Subjects Based on Gender**

The number of male subjects with blood group A, blood group B, blood group AB and blood group O were 6, 10, 3 and 56 respectively, while the number of female subjects with the A, B, AB and O blood groups were 5, 0, 1 and 19 respectively.

Blood Group	Number of Male Subjects (%)	Number of Female Subjects (%)
A	6 (8)	5(20)
B	10 (13)	0(0)
AB	3 (4)	1(4)
O	56 (75)	19(76)
Total	75 (100)	25(100)

**Table 5. Distribution of Rh-Blood Group of COVID-19 Positive Subjects Based on Gender**

The numbers of male subjects who are Rh-positive were 68, while those who are Rh-negative were 7, with percentages of 91 and 9 respectively. Similarly, the numbers of female subjects who are Rh-positive were 23, while those who are Rh-negative were 2, with percentages of 92 and 8, respectively.

Blood Group	Number of Subjects	Percentage (%)
Male		
Rh D -Positive	68	91
Rh D -Negative	7	9
Female		
Rh D -Positive	23	92
Rh D -Negative	2	8

#### 4. DISCUSSION

This study assessed the distribution of some blood group antigens among COVID-19 infected individuals. The results from this study indicated

that there were more male subjects who were positive for COVID-19 than female subjects. This result is in consonance with a previous report by Wenham et al. [20], which reported that more male than female were prone to COVID-19 infection. It is believed that this happens because women produce more interferon (type 1) than men; interferon is known to be vital in early response during COVID-19 infection [21]. Furthermore, estrogen, which is known to be produced in women, has also been reported to confer some protection against infections by way of increasing immune responses by T-cells, raising the production of antibodies, increasing neutrophil count as well as producing cytokine via monocytes and macrophages [22].

Data from the study also indicate that 74 subjects were married while 26 subjects were single. It has been reported that married people tend to seek remedies for COVID-19 because of the fear of infecting other family members [23]. This finding agrees with the work of another author [24].

All the subjects recruited for the study had academic achievements (SSCE, University degree or HND from a Polytechnic) and are thus literates. This indicates a high level of literacy among the subjects. This finding is in agreement with the work of another researcher [25], which reported a similar high percentage of educated subjects in their study. This finding is probably because education increases awareness of a disease [26]. Another study [27] had reported that the high educational attainment in their study may be a reflection of the awareness and understanding of the infection by the subjects.

The data from this study showed that a greater percentage of the subjects (75%) had blood group O. This finding could be attributed to the fact that blood group O is more prevalent (or common) in Nigeria [28]. However, the observed high frequency of blood group O in the study population does not reflect the susceptibility of the population to COVID 19 infection; the high prevalence of this blood group O could be an evolutionary adoption to confer resistance to diseases that affect our population [29]. It has been reported also that blood group O is associated with a decreased risk of COVID-19 infection [30]. SARS-COV-2 which includes the corona virus, has spike proteins which enable it to gain entry into and infect the host cell [31].

The findings of this study agree with that of another finding [32] who conducted a study that

evaluated the ABO blood group distribution of COVID-19 patients at the University of Abuja Teaching Hospital, stating that no relationship was found between COVID-19 and ABO blood group. On the contrary, a study [33] reported an association between the ABO blood group and COVID-19, stating that blood group O was less represented among the patients, while blood groups B and AB were significantly more represented. From this study, the COVID-19 positive subjects with blood group O constituted the largest fraction followed by blood groups A, B and then AB; this report is attributed to the distribution pattern of the ABO blood groups among the subjects. This report follows the trend of normal ABO blood group distribution (blood group O>A>B>AB). This normal ABO trend has been reported by several authors [34-36].

This study showed that 91% of the subjects were Rh D positive while 9% were Rh D negative. In other words, a significant majority of the subjects in this study were Rh D Positive. This finding is in agreement with the work of another researcher [37], who reported that 96.42% of their subjects were Rhesus positive. Rh D negative blood group has been reported to be at lower risk of viral infections as well as the severity of illness and death [38]. Other studies have indicated similar findings. A study [39] had reported that individuals with blood group A Rh D positive and blood group AB Rh D positive are more susceptible to COVID-19 infection than individuals with blood group O- whether Rh D positive or negative still had a significant lower risk of COVID-19 infection. In other words, individuals who are Rh D positive have increased susceptibility to COVID-19 infection. The mechanism by which Rh factor increases susceptibility to COVID-19 infection is not clearly understood, necessitating the need for further research to confirm the role of Rh blood group in the susceptibility of individuals to COVID-19 infection [40]. It has been proposed that the observation in this study may be due to possible interaction between the Rh factor and SARS-COV-2, but there is need for further research to confirm this [41]. In this study, the COVID-19 positive subjects who were Rh D-positive constituted the largest fraction; this report is attributed to the distribution pattern of the Rh D blood group among the subjects. This study has revealed a trend in the Rhesus distribution in Rivers State to be Rh D-positive > Rh D-negative in consonance with regular Rhesus blood group distribution in Nigeria [42,28].

## 5. CONCLUSION

The findings from this study revealed that more of the subjects had blood group O in the study locality, followed by blood groups A, B and AB. The results from this study also revealed that more of the subjects were Rh D positive in the study locality.

It is recommended that more research should be carried out to proffer a better understanding of the mechanism involved in the interactions of blood group antigens and COVID-19 infections in order to help in the treatment and management of the disease.

## CONSENT

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

## ETHICAL CONSIDERATION

Ethical approval for this study was obtained from the Research Ethics Committee of the Ministry of Health, Port-Harcourt, Rivers State, Nigeria.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Intharanut K, Nathalang O. Update on blood group terminology and molecular polymorphisms. *J Hematol Transfus Med.* 2018;28(4):459-5.
2. Daniels G, Reid ME. Blood groups: the past 50 years. *Transfusion.* 2010;50(2):281-9.
3. Yelima JD, Ufelle SA, Milgwe DM, Oyeleke KO, Denué BA, Chikwendu CK. Complications of blood transfusion and management: definitions and history of blood group systems. *Eur J Biol Med Sci Res.* 2019;7(3):25-34.
4. Mitra R, Mishra N, Rath GP. Blood groups systems. *Indian J Anaesth.* 2014;58(5):524-8.
5. Groot HE, Villegas Sierra LEV, Said MA, Lipsic E, Karper JC, van der Harst P. Genetically determined ABO blood group and its associations with health and disease. *Arterioscler Thromb Vasc Biol.* 2020;40(3):830-8.

6. Van Hylckama VA, Flinterman LE, Cannegiester SC, Reitsma PH, Arellano AR, Tong CH et al. Genetic variations associated with recurrent venous thrombosis. *Circ Cardiovasc Genet.* 2014; 7:806-13.
7. Ewald DR, Sumner SCJ. Blood type biochemistry and human disease. *Wiley Interdiscip Rev Syst Biol Med.* 2016; 8(6):517-35.
8. Gundrajukuppam DK, Vijaya SB, Rajendran A, Sarella JD. Prevalence of principal Rh blood group antigens in blood donors at the blood bank of a Tertiary Care Hospital in Southern India. *J Clin Diagn Res.* 2016;10(5):EC07-10.
9. Jahanpour O, Pyuza JJ, Ntiyakunze EO, Mremi A, Shao ER. ABO and rhesus blood group distribution and frequency among blood donors at Kilimanjaro Christian Medical Center, Moshi, Tanzania. *BioMed Central Research Notes.* BMC Res Notes. 2017;10(1):738.
10. Pratima V, Shradha S, Akhilesh K. Prevalence of haemoglobin variants, ABO and rhesus blood groups in northern Uttar Pradesh, India. *Biomed Res.* 2013;24:377-82.
11. Vege S, Westhoff CM. Rh and RhAG blood group systems. In: Shaz BH, Hillver CD, Gil MR, editors. *Transfusion medicine and hemostasis.* 3rd ed. Amsterdam: Elsevier. 2019;149-55.
12. Wafi M, Housse EL, Nourichafi N, Bauisk K, Benajiba MH, Abti N. Prevalence of weak Antigen D among Antigen D Negative C/E and donors in Morocco [international journal] of *Blood Transfusion and Immunohaematology.* 2016;6(1):3-6.
13. Weiss SR, Leibowitz JL. Coronavirus pathogenesis. *Adv Virus Res.* 2011; 81(1):85-164.
14. Wang W, Tang J, Wei F. Updated understanding of the outbreak of 2019 novel coronavirus (2019-nCoV) in Wuhan, China. *J Med Virol.* 2020; 92(4):441-7.
15. Croxen MA, Law RJ, Scholz R, Keeney KM, Wlodarska M, Finlay BB. Recent advances in understanding enteric pathogenic *Escherichia coli*. *Clin Microbiol Rev.* 2013;26(4):822-80.
16. Jaff MS. Relation between ABO blood groups and *Helicobacter pylori* infection in symptomatic patients. *Clin Exp Gastroenterol.* 2011;4:221-6.
17. Barbé L, Le Moullac-Vaidye B, Echasserieu K, Bernardeau K, Carton T, Bovin N et al. Histo-blood group antigen-binding specificities of human rotaviruses are associated with gastroenteritis but not with in vitro infection. *Sci Rep.* 2018;8(1):1.
18. Cooling L. Blood groups in infection and host immunity. *Clin Microbiol Rev.* 2015;28(3):801-70.
19. Moses O, Ugochi EE, Okeke HU, Francis OI. An assessment of the environmental impacts of land use dynamics in eleme, Rivers State, Nigeria. *IOSR J Environ Sci Toxicol Food Technol.* 2020;14(10):41-55.
20. Wenham C, Smith J, Morgan R, Gender and COVID-19 Working Group. COVID-19: the gendered impacts of the outbreak. *Lancet.* 2020;395(10227):846-8.
21. Trouillet-Assant S, Viel S, Gaymard A, Pons S, Richard JC, Perret M et al. Type 1 IFN immunoprofiling in COVID -19 patients. *J Allergy Clin Immunol.* 2020;146(1):206-208.e2.
22. Rehman S, Ravinayagam V, Nahvr I, Aldossary H, Al-shammari M, Al-Amiri MS Et al. Suhaimi, E.A. Front Immunol. Immunity, sex hormones and environmental factors as determinants of COVID-19 disparity in women.2021;12:1-12.
23. Konopińska J, Obuchowska I, Lisowski Ł, Dub N, Kozera M, Rękas M. Intention to get COVID-19 vaccinations among ophthalmology Residents in Poland: A cross sectional survey. *Vaccines.* 2021;9(4):371-80.
24. Cvetković VM, Nikolić N, Ocal A, Martinović J, Dragašević A. A predictive model of pandemic disaster fear caused by coronavirus (COVID-19): implications for decision makers. *Int J Environ Res Public Health.* 2022;19(2):1-27.
25. Adefisoye JO, Adefisoye MA, George F, Bisi-Johnson MA. Knowledge, attitude and perception of people towards the novel coronavirus disease (COVID-19) outbreak and its effects on their livelihood: perspective from diverse populations. *Health Econ Outcome Res.* 2021;7(20):033-42.
26. Olawuni P, Olowoporoku O, Daramola O. Determination of residents' participation in disaster, risk management in Lagos Metropolis, Nigeria. *Int J Disaster Risk Manag.* 2020;2(2):1-18.
27. Oleribe O, Ezechi O, Osita-Oleribe P, Olawepo o, Musa AZ, Omoluabii A et al.

- Public Perception of COVID-19 management and responses in Nigeria: A cross-sectional survey. *BMJ Open*. 2020;10:1-7.
28. Anifowoshe AT, Owolodun OA, Akinseye KM, Iyiola OA, Oyeyemi BF, B.F. (20217). Gene frequencies of ABO and Rh blood groups in Nigeria: A review. *Egypt J Med Hum Genet*. 2017;18(3):205-10.
  29. Arend P. Position of human blood group O (H) and phenotype determining enzyme in growth and infectious diseases. *Ann N Y Acad Sci*. 2018;1425(1):5-18.
  30. Gérard C, Maggipinto G, Minon JM. COVID-19 and ABO blood group: another viewpoint. *Br J Haematol*. 2020;190(2):e93-4.
  31. Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O et al. *Cry- E. Science*. 2020;367(6483):1260-3.
  32. Kwaghe VG, Habib ZG, Bawa A, Adebayo FO, Akor AA, Thairu Y et al. Blood group distribution among COVID-19 patients in a Nigerian treatment centre. *West Afr J Med*. 2021;38(5):494-7.
  33. Kotila TR, Alonge TO, Fowotade A, Famuyiwa OI, Akinbile AS. Association of the ABO blood group with SARS-Cov-2 infection in a community with low infection rate. *Vox Sang*. 2021;116(8):910-5.
  34. Udomah FP, Isaac IZ, Aliyu N, Erhabor O, Ahmed MH, Yakubu A et al. Haemoglobin electrophoretic patterns, ABO and rhesus D blood groups distribution among antenatal women in Sokoto, Nigeria. *Obstet Gynaecol Cases Rev*. 2015;2(3):2-4.
  35. Ifeanyi OE. An update on susceptibility of individuals to diseases based on ABO blood groups. *Int J Curr Res Med Sci*. 2019;5(3):1-8.
  36. Christian SG, Eze EM, Ezimah ACU, Buseri FI. Rhesus E, rhesus D and ABO blood groups distribution among indigenes of Ogoni ethnic group of Rivers State, Nigeria. *Int J Res Rep Hematol*. 2020; 3(1):1-7.
  37. Arak E, Solmaz, Akkoc H, Donmezdil S, Karahan Z, Kaya S. mertsoy, y., Yildirim. M.S., Ekin, N., Arac, S. & Demir, C. (2020). Association between the Rh blood group and the COVID-19 susceptibility. *International Journal of Haematology and Oncology*, 30 (2), 81-86.
  38. Zietz M, Zucker J, Tatonetti NP. Association between blood type and COVID-19 infection, intubation and death. *Nat Commun*. 2020;119(1):1-22.
  39. Taha SAH, Osman MEM. abdoelkarim, E.A.A., Holie, M.A. I. Elbasheir, M.M., Abuzeid, N.M.K., al-Thobaiti, S.A. Fabul, S.B. & Knozu, E.H.E. New Microbes New Infect. Individuals with a Rh-positive but not Rh-negative blood group are more vulnerable to SARS-COV-2 infection: demographics and trend study on COVID-19 cases in Sudan. 2020;38:1-7.
  40. Saify K, Alborz MS, Saadat M. Susceptibility to the novel coronavirus disease (COVID-19) is associated with ABO and Rh blood groups: A case-control study from Afghanistan. *Egypt J Med Hum Genet*. 2021;22(1):1-5.
  41. Anderson JL, May HT, Knight T, Bair TL, Horne, B. D. & knowtton, K. U. *Br J Haematol*. 2022. Association of Rhesus factor blood type with risk of SARS-COV-2 infection and COVID-19 severity;100:1-3.
  42. Erhabor O, Adias TC, Jeremiah ZA, Hart ML. Abnormal haemoglobin variants, ABO, and rhesus blood group distribution among students in the Niger Delta of Nigeria. *Pathol Lab Med Int*. 2010;2(1):31-46.

© 2022 Davies et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
 The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/90321>