



Patients with Parathyroid Disorders: Oral Manifestations and Dental Management

Georges Aoun ^{a*}, Houssam Abou Hamdan ^b and Pamela Aoun ^c

^a Department of Oral Medicine and Maxillofacial Radiology, Faculty of Dental Medicine, Lebanese University, Beirut, Lebanon.

^b Department of Periodontics, Faculty of Dental Medicine, Lebanese University, Beirut, Lebanon.

^c Department of Food Technology, American University of Beirut, Beirut, Lebanon.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMPS/2021/v23i1230274

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/83719>

Review Article

Received 13 October 2021
Accepted 26 December 2021
Published 28 December 2021

ABSTRACT

Parathyroid hormone secreted by the parathyroid glands has a key role in the regulation of calcium and phosphorus metabolism and plays an essential role in tooth and bone mineralization. Disorders of the parathyroid glands most frequently result in abnormalities of serum calcium and can induce various oral and extra-oral manifestations. Therefore, a sound understanding of these conditions is essential for the dental practitioner, with emphasis on alerting signs, clinical and radiological findings and mandatory communication with patient's physician.

Keywords: Dentistry; hyperparathyroidism; hypoparathyroidism; management; oral manifestations; parathyroid gland.

1. INTRODUCTION

The parathyroid glands, located behind the thyroid gland, are part of the endocrine system. They are generally four, though additional glands have been found in up to 13% of autopsies [1].

Microscopically, the parathyroid glands are composed of two different types of cells, the chief cells and the oxyphil cells. While the functions of the oxyphil cells are not completely understood, the chief cells, which are smaller and more numerous, control the secretion of

*Corresponding author: E-mail: dr.georges@aoun@gmail.com, aoungesorges@yahoo.com;

parathyroid hormone (PTH) [2]. PTH has a key role in the regulation of calcium and phosphorus metabolism [3,4]; it enhances the bone resorption, consequently releasing the calcium in the bloodstream, augments the formation of the active form of Vitamin D in the kidneys, thereby promoting the absorption of calcium in the intestine and lastly, it decreases renal phosphate reabsorption [2,4].

Parathyroid disorders include either hyperparathyroidism (increase of PTH secretion) or hypoparathyroidism (decrease of PTH secretion). They can induce various oral and/or extra-oral manifestations [5], accordingly, the dental practitioner must be aware of these findings to be able to correctly diagnose and treat the patient.

The aim of this article is to review the parathyroid disorders and their implications in dentistry.

2. PARATHYROID DISORDERS

2.1 Hyperparathyroidism

Hyperparathyroidism affects approximately 0.05 to 0.1% of the general population with an incidence of 1/400 in women and 1/1000 in men usually in their middle age [6]. It is classified into 3 types: a) primary, b) secondary, and c) tertiary.

Primary hyperparathyroidism is usually caused by an adenoma of the gland (85% of cases) leading to a PTH hypersecretion and consequently to hypercalcemia and hypophosphatemia [4,5]. This condition is normally asymptomatic but it can result in bone

disease, renal calculi, or neuromuscular symptoms [8].

Secondary hyperparathyroidism is generally due to increased quantities of hormones via stimulation of parathyroid glands in order to correct abnormally low levels of serum calcium in various conditions such as chronic kidney disease, intestinal malabsorption syndrome, decreased Vitamin D production [4,5].

Tertiary hyperparathyroidism results from chronic secondary hyperparathyroidism.

Table 1. Summarizes the typical signs and symptoms of hyperparathyroidism / hypercalcemia [9,10-12]

Diagnosis of primary hyperparathyroidism is based on presented signs and symptoms, an elevated PTH, and a high serum calcium level [7,13]. In the secondary type of the disease, the serum calcium level can be low or normal [14].

2.2 Hypoparathyroidism

Hypoparathyroidism commonly occurs after surgery in the neck region leading to the glands' damage and it can also be caused by an autoimmune destruction of the glands or developmental defects [7, 8].

Table 2 summarizes the typical signs and symptoms of hypoparathyroidism / hypocalcemia [7,9,15-17].

Diagnosis of hypoparathyroidism is based on existing signs and symptoms, low serum levels of calcium and PTH, and a high serum level of phosphorus.

Table 1. Signs and symptoms of hyperparathyroidism / hypercalcemia

Organ	Signs and symptoms
Kidney	Polydipsia, polyuria, kidney stones
Gastrointestinal tract	Anorexia, constipation, epigastric pain, nausea, vomiting
Cardiovascular system	Angina, dyspnea, palpitations, syncope
Neuromuscular system	Anxiety, confusion, depression, fatigue, forgetfulness, impaired vision, insomnia, lethargy, weakness
Skeleton	Bones pain, fractures

Table 2. Signs and symptoms of hypoparathyroidism / hypocalcemia

Organ	Symptoms
Cardiovascular system	Dyspnea, edema, palpitations, syncope
Neurologic system	Headache, impaired vision, neuropsychiatric symptoms
Neuromuscular system	Labial and perioral numbness, paresthesia of finger and toes, myalgia, muscular spasm, laryngospasm or bronchospasm
Ectodermal tissues	Alopecia, loss of axillaries and pubic hair, roughness of body hair, nails deformities

Table 3. Oral manifestations of hyperparathyroidism

Oral manifestations of patients with hyperparathyroidism	
-	Brown tumor
-	Decrease in bone density
-	Teeth mobility and drifting
-	Indistinct pain of the jaw
-	Sensitive teeth
-	Soft tissue calcifications
-	Dental abnormalities (development defects, alterations in dental eruption)
-	Obliteration of pulp chamber by pulp stone
-	Sialolithiasis

2.3 Oral manifestations of Parathyroid Disorders

The most common oral manifestation associated with hyperparathyroidism is the brown tumor. This lesion develops in any of the three types of hyperparathyroidism and is characterized by an extensive bone resorption, which is replaced by fibrovascular tissue and giant cells with abundant hemorrhage and hemosiderin deposition [18]. In the oral cavity, it usually occurs at the level of the mandible, especially in the premolars and molars region; it is rare in the maxilla [4,18]. Radiographically, it shows as well-defined uni- or multilocular radiolucencies with a widespread loss of the lamina dura and changes in the pattern of the trabecular bone of the jaw. Chronic lesions normally induce an important expansion of cortical bone, and roots resorption and displacement [4].

Table 3 summarized additional oral manifestations of hyperparathyroidism.

For hypoparathyroidism, the most common oral manifestations are enamel hypoplasia and delayed dental eruption. Additionally, oral chronic candidiasis, tongue or lips paresthesia and facial twitching can occur [8].

2.4 Dental Management of Patients with Parathyroid Disorders

Dental management of patients with parathyroid disorders does not necessitate any special consideration. However, some recommendations have been implemented in order to avoid any risk associated with these conditions.

As a matter of fact, it is essential to know that a high bone fracture risk exists in patients with hyperparathyroidism and consequently the practitioner must take into consideration specific precautions in surgical treatments. Moreover, it is mandatory to correctly diagnose a brown

tumor and adequately treat it; in some cases, spontaneous regression of the lesion occurs after surgical treatment of the glands; however, in other cases the tumor does not regress and must be removed surgically [18,19].

Furthermore, the dental practitioner must be aware of the susceptibility of the patients to caries because of dental abnormalities such as enamel hypoplasia [4].

3. CONCLUSION

Even though the majority of the oral manifestations occurring in patients with parathyroid disorders are not specific, the dental practitioner must be aware of these conditions, with emphasis on alerting signs along with clinical and radiological findings. In addition, a communication with the patient's physician for further medical investigation may be crucial for the success of the treatment.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Tattera D, Wong LM, Vikse J, Sanna B, Pękala P, Walocha J, et al. The prevalence and anatomy of parathyroid glands: a meta-analysis with implications for parathyroid surgery. *Langenbecks Arch Surg.* 2019;404(1):63-70.

- DOI: 10.1007/s00423-019-01751-8.
2. Ilahi A, Muco E, Ilahi TB. Anatomy, Head and Neck, Parathyroid. [Updated 2021 Aug 11]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022. Available: <https://www.ncbi.nlm.nih.gov/books/NBK537203/>
 3. Žofková I. Hypercalcemia. Pathophysiological aspects. *Physiol Res*. 2016;65(1): 1-10. DOI: 10.33549/physiolres.933059.
 4. Mittal S, Gupta D, Sekhri S, Goyal S. Oral manifestations of parathyroid disorders and its dental management. *J Dent Allied Sci* 2014;3:34-38. DOI: 10.4103/2277-4696.156527
 5. Khalekar Y, Zope A, Brahmanekar U, Chaudhari L. Hyperparathyroidism in dentistry: Issues and challenges!! *Indian J Endocrinol Metab*. 2016;20(4):581-582. DOI: 10.4103/2230-8210.183452.
 6. Wu JX, Yeh MW. Asymptomatic Primary Hyperparathyroidism: Diagnostic Pitfalls and Surgical Intervention. *Surg Oncol Clin N Am*. 2016;25(1):77-90. DOI: 10.1016/j.soc.2015.08.004.
 7. Michels TC, Kelly KM. Parathyroid disorders. *Am Fam Physician*. 2013;88(4): 249-257.
 8. Kelly A, Pomarico L, de Souza IP. Cessation of dental development in a child with idiopathic hypoparathyroidism: a 5-year follow-up. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2009;107(5):673-677. DOI: 10.1016/j.tripleo.2008.11.017.
 9. Sarkar S, Mondal M, Das K, Shrimal A. Mucocutaneous manifestations of acquired hypoparathyroidism: An observational study. *Indian J Endocrinol Metab*. 2012;16(5):819-820. DOI: 10.4103/2230-8210.100637.
 10. Stearns M, Cox J. Parathyroid disease: symptoms, differential diagnosis, and management In: Arora A, Tolley NS, Tuttle RM, eds. *A Practical Manual of Thyroid and Parathyroid Disease*. Chichester, UK: Wiley-Blackwell. 2010;145-163.
 11. Taniegra ED. Hyperparathyroidism. *Am Fam Physician*. 2004;69(2):333-339.
 12. Silverberg SJ, Bilezikian JP. The diagnosis and management of asymptomatic primary hyperparathyroidism. *Nat Clin Pract Endocrinol Metab*. 2006;2(9):494-503. DOI: 10.1038/ncpendmet0265.
 13. Pallan S, Rahman MO, Khan AA. Diagnosis and management of primary hyperparathyroidism. *BMJ*. 2012;344:e1013. DOI: 10.1136/bmj.e1013.
 14. Muppidi V, Meegada SR, Rehman A. Secondary Hyperparathyroidism. [Updated 2021 Sep 6]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. Available: <https://www.ncbi.nlm.nih.gov/books/NBK557822/>
 15. Shoback D. Clinical practice. Hypoparathyroidism. *N Engl J Med*. 2008; 359(4):391-403. DOI: 10.1056/NEJMcp0803050.
 16. Bringhurst FR, Demay MB, Kronenberg HM. Hormones and disorders of mineral metabolism. In: Larsen PR, Kronenberg HM, Melmed S, Polonsky KS, eds. *Williams Textbook of Endocrinology*. 10th ed. Philadelphia, Pa.: Saunders. 2003;1303-1348.
 17. Cooper MS, Gittoes NJ. Diagnosis and management of hypocalcaemia. *BMJ*. 2008;336(7656):1298-1302. DOI:10.1136/bmj.39582.589433.BE. Erratum in: *BMJ*. 2008;336(7659): DOI: 10.1136/bmj.a334.
 18. Shetty AD, Namitha J, James L. Brown tumor of mandible in association with primary hyperparathyroidism: a case report. *J Int Oral Health*. 2015;7(2):50-52.
 19. Triantafillidou K, Zouloumis L, Karakinaris G, Kalimeras E, Iordanidis F. Brown tumors of the jaws associated with primary or secondary hyperparathyroidism. A clinical study and review of the literature. *Am J Otolaryngol*. 2006;27(4):281-286. DOI: 10.1016/j.amjoto.2005.11.004.

© 2021 Aoun et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/83719>