



Evaluation of Patients' Responses to Oral Hypoglycemic Agents at a University Health Centre

Olugbenga Matthew Ajulo^{1*} and Oluwatoyin Hephzibah Ajulo²

¹Faculty of Pharmacy, University of Uyo, Uyo, Akwa-Ibom State, Nigeria.

²Faculty of Agriculture, University of Uyo, Uyo, Akwa-Ibom State, Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2018/v25i530113

Editor(s):

(1) Dr. Jongwha Chang, University of Texas, College of Pharmacy, USA.

Reviewers:

(1) Dennis, Amaechi, Veritas University, Abuja, Nigeria.

(2) Muruganandan Shanmugam, Wayne State University, USA.

(3) Rupali Sengupta, SNTD Women's University, India.

(4) Arthur N. Chuemere, University of Port Harcourt, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/46851>

Original Research Article

Received 25 October 2018

Accepted 18 January 2019

Published 11 March 2019

ABSTRACT

Background: There were 1.5 million deaths caused by diabetes in 2012, of which more than 80% of diabetes deaths occurred in developing countries. WHO estimated diabetes would be the 7th leading cause of death by 2030.

Aim: The study aimed at evaluating type 2 diabetes mellitus patients' clinical responses after use of oral hypoglycaemic agents.

Study Design: The study was a retrospective observational study.

Place and duration of Study: The study was undertaken at Primary healthcare facility, University Health Centre. The study monitored case notes of type 2 diabetes mellitus patients who attended the endocrinology clinic within the ten years of review.

Methods: After ethical approval was given, a retrospective evaluation of type 2 diabetes mellitus patients' folders was done for one hundred and nineteen patients who attended the endocrinology clinic. Relevant information obtained from patients' folders were collated and analysed.

Results: Out of one hundred and nineteen participants who received oral hypoglycaemic agents,

*Corresponding author: E-mail: ajugbeng@gmail.com, matthewajulo@uniuyo.edu.ng;

seventy-six (63.8%) participants were in the age range of 45-55 years, followed by twenty-four (20.2%) participants with age range of greater than 55 years. Sixty-eight (57.1%) participants were females while fifty-one (42.9%) were males. Forty two (35.3%) participants had a controlled plasma glucose level of <110 mg/dl while seventy-seven (64.7%) participants had plasma glucose level of >110 mg/dl. Efficacy index was highest for Daonil+Glycomet followed by Diabinese+Glucophage and Glucovance respectively.

Conclusion: The study indicated that fewer type 2 diabetes mellitus patients' plasma glucose levels were controlled by two drugs combination therapy involving metformin.

Keywords: Diabetes mellitus; oral hypoglycaemic agents; efficacy index; plasma glucose.

1. INTRODUCTION

The human, social and economic consequences of non-communicable diseases were felt by all countries but were particularly devastating in developing countries of the world. Reducing the global burden of non-communicable diseases was an overwhelming priority and an unavoidable condition for sustainable development. Non-communicable diseases was the leading cause of death globally responsible for 38 million (68%) of world's 56 million deaths in 2012. Sixteen million were premature deaths under age 70 years. 28 million deaths associated with non-communicable diseases occurred in developing countries and mostly (82%) premature deaths [1].

The leading causes of deaths associated with non-communicable diseases in 2012 were reported as cardiovascular diseases (17.5 million deaths), cancers (8.2 million deaths), respiratory diseases (4 million deaths) and diabetes (1.5 million deaths). These four major non-communicable diseases were responsible for 82% of death associated with non-communicable diseases [2].

Diabetes mellitus was a chronic disease that occurred when the pancreas did not produce enough insulin or when the body could not effectively use the insulin it produced. Insulin was an hormone that regulated blood glucose [3]. Hyperglycaemia was a common effect of uncontrolled diabetes and over time led to serious damage to many of the body's systems, especially the nerves and blood vessels. The global prevalence of diabetes in 2014 was estimated to be 9% among adults aged 18 years and above [4]. There were 1.5 million deaths caused by diabetes in 2012. More than 80% of diabetes deaths occurred in developing countries [3]. WHO estimated diabetes would be the 7th leading cause of death by 2030 [4]. Africa

experienced an increasing prevalence of diabetes mellitus [5]. In 2010, 12.1 million people were assumed to be living with diabetes mellitus in Africa and it was expected to increase to 23.9 million by 2030 [6]. Diabetes assumed to cause other diseases such as cardiovascular disease, renal disease, pneumonia, bacteraemia and tuberculosis [7-12]. Consequently, it increased morbidity and mortality in the region [13-18]. Therefore attention should be given to the management of diabetes mellitus.

The total economic cost of diabetes mellitus in the Africa region in 2000 was US\$67.03 billion, or US\$8836 per person with diabetes per year [19]. The prevalence of diabetes mellitus (T2DM) appeared to have increased considerably from that recorded in earlier surveys conducted in the region, which found the prevalence in Sub-Saharan Africa was typically below 1%, with the exception of studies in South Africa (3.6%) and the Ivory Coast (5.7%) [20-21].

The main goal of treatment of diabetes mellitus was to recreate normal or nearly normal blood glucose levels without causing low blood glucose while preventing tissue damage due to hyperglycemia. The main goal of treatment was to obtain an HbA1c of 6.5% or fasting glucose of less than 6.1mmol/L (less than 110mg/dL) [22]. There were many brands of oral hypoglycaemic agents used in Nigeria to treat diabetes mellitus. This study aimed at evaluating clinical responses to different oral hypoglycaemic agents used in the University Health Center, Uyo.

2. METHODS

2.1 Study Design

It was a retrospective observational study. A survey of records of patients on hypoglycaemic agents were observed, collated and compared.

2.2 Study Setting

This study was undertaken in a secondary healthcare facility located in Uyo. Records of patients attending endocrinology clinic were used for the study.

2.3 Study Location

The study took place in Endocrinology clinic, University Health Centre, University of Uyo, Akwa-Ibom state, Nigeria. The Health Centre was a primary healthcare facility with about 50 bed spaces. The Health Centre had nine medical practitioners, five Pharmacists, twenty nurses, five medical laboratory scientists and a radiographer.

2.4 Ethical Consideration

The Ethics and Research Committee of the University Health Centre approved the study to be carried out at the Centre.

2.5 Study Population

Folders of one hundred and nineteen type 2 diabetic mellitus patients who attended the Endocrinology Clinic at the University Health Centre for the management of their disease condition were used for the collation of data.

2.6 Sample Size

All the available folders of type 2 diabetes mellitus patients attending endocrinology clinic in the Health Centre were used for the survey. One hundred and nineteen folders were used.

2.7 Data Collection

After the study gained approval from the Health Centre Research Committee, the Health Centre record book was used to select

folders of patients that were currently attending the Centre for the management of type 2 diabetes mellitus. The medical information of the participants that were extracted from the record book included age, weight, gender, patients' complaints, diagnostic test report, physicians' diagnostics, prescribed medication and serum glucose level. The reported serum glucose levels were taken after participants had taken oral hypoglycaemic medications for three months. Data were collated from one hundred and nineteen folders of type 2 diabetes mellitus patients who attended the Centre from December 2013 to November 2014.

2.8 Inclusion Criteria

These included patients diagnosed of type 2 diabetes mellitus and patients were receiving oral hypoglycaemic agents.

2.9 Exclusion Criteria

The study excluded patients that were not having type 2 diabetes mellitus and patients that were receiving non-oral hypoglycaemic agents such as injectables and insulin.

2.10 Data Analysis Plan

Descriptive statistical tools were used to analyse serum glucose levels of participants. SPSS version 21 software package was used for the statistical analysis.

A format of data obtained from patient's folder was shown in Table 1.

2.11 Data Analysis

Data were stored in Microsoft word and analyse by using descriptive analysis and chi test. SPSS version 21 software package was used while significance was considered at $p=0.05$.

Table 1. A format of data obtained from patient's folder

S/n	Patient information			Diagnostic report	Physician diagnostic	Prescription on clinic visit				Pharmacist's intervention	Drug related problem	Diet control report	Exercise	Clinical response
	Complaint	Weight	Age			V1	V2	V3	V4					

3. RESULTS

One hundred and nineteen patients' folders were assessed for clinical response after use of oral hypoglycaemic agents. Among the study participants, diabetes mellitus was most prevalent (63.86%) in age 45-55 years old, followed by ages 55 years (20.17%). Among diabetic patients below 45 years of age, 13 (68.4%) patients had blood glucose level above 110 mg/dL. Among diabetic patients between 45-55 years old, 57 (75%) patients had blood glucose above 110 mg/dL. Among diabetic patients above 55 years old, 14 (58.3%) patients

had blood glucose above 110 mg/dL (Table 2). Oral hypoglycaemic agents could not reduce blood glucose to 110 mg /dL in 28 (54.9%) male diabetic patients and 29 (42.6%) female diabetic patients (Table 3).

Clinical responses of diabetic patients showed that combination of Daonil+Glucomet controlled plasma glucose below 110 mg/dl in 50% of users while Diabinese+Glucophage controlled plasma glucose below 110 mg/dl in 45.5% of users. Glucovance controlled plasma glucose below 110 mg/dl in 42.9% of users (Table 4).

Table 2. Comparison of age with clinical response to oral hypoglycaemic agents

Age	Clinical response			No of participants
	<90 mg/dL	90-110 mg/dL	>110 mg/dL	
<45 years	1 (5.26%)	5 (26.3%)	13 (68.4%)	19 (16.0%)
45-55 years	5 (6.6%)	14 (18.4%)	57 (75%)	76 (63.8%)
>55 years	1 (4.2%)	9 (37.5%)	14 (58.3%)	24 (20.2%)
Total	7	28	84	119

Table 3. Comparison of sex with clinical response to oral hypoglycaemic agents

Sex	Clinical response			No of participants
	<90 mg/dL	90-110 mg/dL	>110 mg/dL	
Male	10 (19.6%)	13 (25.5%)	28 (54.9%)	51 (42.9%)
Female	13 (19.1%)	26 (38.2%)	29 (42.6%)	68 (57.1%)
Total				119

Table 4. Clinical responses of oral hypoglycaemic agents

Drug therapy	Clinical response		Frequency
	<110 mg/dL	≥110 mg/dL	
Daonil + Glucophage	30 (37.0%)	51 (62.9%)	81 (68.1%)
Diabinese + Glucophage	5 (45.5%)	6 (54.5%)	11 (9.2%)
Daonil + Glycomet	1 (50.0%)	1 (50.0%)	2 (1.7%)
Daonil	3 (16.7%)	15 (83.3%)	18 (15.1%)
Glucovance	3 (42.9%)	4 (57.1%)	7 (5.9%)
Total	42	77	119

Glucophage: Metformin, Daonil: Glibenclamide, Diabinese: Chlorpropamide, Glycomet: Metformin, Glucovance: Metformin + Glibenclamide

Table 5. Efficacy index

Drug therapy	Clinical response		Efficacy index
	% Benefit	% No benefit	
Daonil + Glucophage	37.0%	63.0%	0.37
Diabinese + Glucophage	45.5%	54.5%	0.46
Daonil + Glycomet	50.0%	50.0%	0.5
Daonil	16.7%	83.3%	0.17
Glucovance	42.9%	57.1%	0.43

Glucophage: Metformin, Daonil: Glibenclamide, Diabinese: Chlorpropamide, Glycomet: Metformin, Glucovance: Metformin + Glibenclamide

Efficacy index was highest among users of oral hypoglycaemic agents who received Daonil + Glucomet combination followed by those who received Diabinese + Glucophage and Glucovance only (Table 5).

4. DISCUSSION

Africa profoundly increased in prevalence of non-communicable diseases such as diabetes mellitus from 12.1 million people living with diabetes in 2010 to 23.9 million people living with diabetes mellitus in 2030 [22]. Diabetes mellitus was most prevalent in the 45-55 years age group. Similar age group, 45-64 years was reported in a study in Asia as most prevalent [23]. Another study which determined the global prevalence of diabetes concluded that most prevalent age group in developing countries of the world was 45-64 years [24]. Their report supported this study outcome. Another study in Nigeria on prevalence of diabetes indicated an increasing prevalence of diabetes with increasing age [25]. The age range >55 years were few probably due to limited resources to maintain the health challenges of the elderly and to ameliorate the complications of uncontrolled type 2 diabetes mellitus [9].

In this study, more than half of the study participants in the three age groups could not have their plasma glucose lower to 110 mg/dl suggesting difficulty of managing diabetes mellitus in developing countries with limited resources and a predisposition to risks associated with uncontrolled diabetes mellitus such as cardiovascular risk. Previous study had documented that majority of diabetes mellitus patients on oral hypoglycaemic agents did not have controlled plasma glucose level. Previous study had indicated cardiovascular risk as one of the consequences of uncontrolled plasma glucose [26].

In this study more than half of the study's male participants could not have their plasma glucose lower to 110 mg/dl while less than half of the study female participants could not have their plasma glucose lower to 110mg/dl suggesting improved clinical responses in female participants. This report was in contrast to an earlier study which indicated that it was difficult to achieve glucose control in female diabetic participants [27]. The difference in our reports may be due to the fact that their study involved the use of both insulin and oral hypoglycaemic agents while this study involved the use of oral

hypoglycaemic agents only. Other study indicated that combination of hypoglycaemic agent and insulin glardine could control plasma glucose [28].

Nearly 20% of both male and female participants had lowered plasma glucose below 90 mg/dl suggesting. No study has suggested that there was a lowering of plasma glucose to 90 mg/dl by oral hypoglycaemic agents. However, drug adherence, lifestyle modification and diet control could make glycemic goal achievable [29].

Daonil+Glycomet, Diabinese+Glucophage and Glucovance produced nearly 50% users with lowered plasma glucose at <110mg/dl suggesting efficacy at reducing plasma glucose. This combination therapies involved metformin which justified the reason for its inclusion in first line therapy. Metformin had been recommended as first line therapy in the management of type 2 diabetes mellitus [30].

The two drugs combinations including different brands of metformin with Daonil did not indicate similar efficacy which probably suggested the effect of patients' factors such as drug adherence, body mass index or drug formulation effect. Previous study explained different pattern of adherence among type 2 diabetes mellitus participants [31].

Most participants did not have plasma glucose control with the use of two combination therapies. Thus three drugs combination therapy and life style modification might be adequate to control plasma glucose in those patients whose plasma glucose levels were uncontrolled.

5. CONCLUSION

This study indicated that fewer participants had plasma glucose controlled with two drugs combinations that included metformin.

6. RECOMMENDATION

Type 2 diabetes mellitus patients whose plasma glucose levels were uncontrolled with two drugs combination should be given education on lifestyle modification and encouraged to commence three drugs combination therapy.

CONSENT

It is not applicable.

ETHICAL APPROVAL

The ethical approval was granted by the Research Ethics Committee of the University Health Centre, University of Uyo.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organisation. Global status report on non-communicable diseases: Attaining the nine global noncommunicable diseases targets; a shared responsibility. www.who.int. Geneva; 2014.
2. World Health Organization. Global Health Estimates: Deaths by Cause, Age, Sex and Country, 2000-2012. Geneva, WHO; 2014.
3. World Health Organisation. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: Diagnosis and classification of diabetes mellitus. Geneva, World Health Organization; 1999. (WHO/NCD/NCS/99.2)
4. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med*. 2006;3(11): 3:e442.
5. World Health Organisation. The Global Burden of Disease: Update World Health Organisation: Geneva; 2004.
6. Sicree R, Shaw J, Zimmet L. The Global Burden: Diabetes and Impaired Glucose Tolerance. *Diabetes Atlas*, IDF. 4 Edition. International Diabetes Federation: Brussels; 2009.
7. Saydah SH, Eberhardt MS, Loria CM, Brancati FM. Age and the burden of death attributable to diabetes in the United States, *Am J Epidemiol*. 2002;156(8):714-19.
8. Brown WV. Microvascular complications of diabetes mellitus: Renal protection accompanies cardiovascular protection. *Am J Cardiol*. 2008;102(12A):10L-13L.
9. Kornum JB, Thomsen RW, Riis A, Lervang HH, Schønheyder HC, Sørensen HT. Diabetes, glycemic control, and risk of hospitalization with pneumonia: A population-based case-control study. *Diabetes Care*. 2008;31(8):1541-45.
10. Thomsen RW, Hundborg HH, Lervang HH, Johnsen SP, Schønheyder HC, Sørensen HT. Risk of community-acquired pneumococcal bacteremia in patients with diabetes: A population-based case-control study. *Diabetes Care*. 2004;27(4):1143-47.
11. Thomsen RW, Hundborg HH, Lervang HH, Johnsen SP, Schønheyder HC, Sørensen HT. Diabetes mellitus as a risk and prognostic factor for community-acquired bacteremia due to enterobacteria: A 10-year, population-based study among adults. *Clin Infect Dis*. 2005;40(4):628-31.
12. Jeon CY, Murray MB. Diabetes mellitus increases the risk of active tuberculosis: A systematic review of 13 observational studies. *PLoS Med*. 2008;5(7):e152.
13. Rudan I, Boschi-Pinto C, Biloglav Z, Mulholland K, Campbell H. Epidemiology and etiology of childhood pneumonia. *Bull World Health Organ*. 2008;86(5):408-16.
14. Seale AC, Mwaniki M, Newton CR, Berkley JA. Maternal and early onset neonatal bacterial sepsis: Burden and strategies for prevention in sub-Saharan Africa. *Lancet Infect Dis*. 2009;9(7):428-38.
15. Mayanja BN, Todd J, Hughes P, Van Der Paal L, Mugisha JO, Atuhumuza E, et al. Septicaemia in a population-based HIV clinical cohort in rural Uganda, 1996-2007: Incidence, aetiology, antimicrobial drug resistance and impact of antiretroviral therapy. *Trop Med Int Health*. 2010;15(6): 697-05.
16. Reddy EA, Shaw AV, Crump JA. Community-acquired bloodstream infections in Africa: A systematic review and meta-analysis. *Lancet Infect Dis*. 2010;10(6):417-32.
17. Steen TW, Aruwa JE, Hone NM. The epidemiology of adult lung disease in Botswana. *Int J Tuberc Lung Dis*. 2001; 5(8):775-82.
18. Lonnoth K, Kastro KG, Chakaya JM, Chanuhan LS, Floyd K, Glaziou P, Raviglione MC. Tuberculosis control and elimination 2010-50: Cure, care, and social development. *Lancet*. 2010;375(9728): 1814-29.
19. Kirigia JM, Sambo HB, Sambo LG, Barry SP. Economic burden of diabetes mellitus in the WHO African region. *BMC Int Health Hum Rights*. 2009;9:6. DOI: doi.org/10.1186/1472-698X-9-6
20. Marine N, Vinik AI, Edelstein I, Jackson WP. Diabetes, hyperglycemia and glycosuria among Indians, Malays and

- Africans (Bantu) in Cape Town, South Africa. *Diabetes*. 1969;18(12):840-57.
21. McLarty DG, Pollitt C, Swai AB. Diabetes in Africa. *Diabet Med*. 1990;7(8):670-84.
 22. Motala AA, Omar MA, Pirie FJ. Diabetes in Africa: Epidemiology of type 1 and type 2 diabetes in Africa. *J Cardiovasc Risk*. 2003;10(2):77-3.
 23. Hall V, Thomsen RW, Henriksen O, Lohse N. Diabetes in Sub-Saharan Africa 1999-2011: Epidemiology and public health implications, a systematic review. *BMC Public Health*. 2011;11:564. DOI: 10.1186/1471-2458-11-564
 24. Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: Estimates for the year 2000 and projections for 2030. *Diabetes care*. 2004; 27(5):1047-53.
 25. The DECODA study group. Age- and sex-specific prevalence of diabetes and impaired glucose regulation in 11 Asian cohorts. *Diabetes care*. 2003;26:1770-80.
 26. Adeloye D, Ige JO, Aderemi VA, Adeleye N, Amoo EO, Auta A, Oni G. Estimating the prevalence, hospitalisation and mortality from type 2 diabetes mellitus in Nigeria: A systematic review and meta-analysis. *BMJ Open*. 2017;7:e015424. DOI: 10.1136/bmjopen-2016-015424
 27. McGills JB, Vlajnic A, Knutsen PG, Recklein C, Rimler M, Fisher SJ. Effect of gender on treatment outcomes in type 2 diabetes mellitus. *Diabetes Res Clin Pract*. 2013;102(3):167-74.
 28. Ketema EB, Kibret KT. Correlation of fasting and postprandial plasma glucose with HbA1c in assessing glycemic control, systematic review and meta analysis. *Archives of Public Health*. 2015;73:43. DOI: 10.1186/s13690-015-0088-6
 29. Arnetz L, Ekberg ND, Alvarsson M. Sex differences in type 2 diabetes: Focus on diseases course and outcomes. *Diabetes Metab Syndr Obes*. 2014;7:409-20. DOI: 10.2147/DMSO.S51301
 30. Fujihara K, Igarashi R, Matsunaga S, Matsubayashi Y, Yamada T, Yokoyama H, et al. Comparison of baseline characteristics and clinical course in Japanese patients with type 2 diabetes among whom different types of oral hypoglycaemic agents were chosen by diabetes specialists as initial monotherapy (JDDM42). *Medicine*. 2017;96:7. Available:<http://dx.doi.org/10.1097/MD.0000000006122>
 31. De Vries McClintock HF, Morales KH, Small DS, Bogner HR. Pattern of adherence to oral hypoglycaemic agents and glucose control among primary care patients with type 2 diabetes. *Behav Med*. 2016;42(2):63-71.

© 2018 Ajulo and Ajulo; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/46851>*