



Growth Response, Carcass Yield and Meat Quality of West African Dwarf Sheep Fed Graded Level of Shea Cake Meal

**D. O. Okunlola^{1*}, A. J. Amuda², M. D. Shittu³, O. O. Olatunji⁴,
O. T. Ojoawo³, O. S. Olaniyan⁴, B. O. Olateju¹, A. A. Fasola¹ and A. J. Alao¹**

¹Department of Animal Nutrition and Biotechnology, Ladoké Akintola University of Technology, Ogbomoso, Oyo State, Nigeria.

²Department of Animal Production and Health, Federal University, Wukari, P.M.B. 1020, Taraba State, Nigeria.

³Department of Animal Production and Health, Ladoké Akintola University of Technology, Ogbomoso, Oyo State, Nigeria.

⁴Department of Animal Science, University of Ibadan, Ibadan, Oyo State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JABB/2021/v24i730237

Editor(s):

(1) Dr. Maria Serrano, University Miguel Hernández, Spain.

Reviewers:

(1) Farhad mirzaei, Animal Science Research Institute, Iran.

(2) Brishketu Kumar, Navsari Agricultural University, India.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here: <https://www.sdiarticle5.com/review-history/77197>

Original Research Article

Received 20 July 2021

Accepted 24 September 2021

Published 26 September 2021

ABSTRACT

Growth response, carcass characteristics and meat quality of West African Dwarf (WAD) sheep fed Shea Cake Meal (SCM) supplement was investigated. A total of 24 WAD sheep of average body weight ranges from 11-12.65 kg comprising of twelve (12) males and twelve (12) females were randomly allotted to four dietary treatments with six animals per treatment in a completely randomized design in a twelve (12) weeks experimental period. Treatment 1 was control and had no inclusion of SCM. Treatments 2, 3 and 4 had 10%, 15% and 20% inclusion of SCM. Results were significantly different ($P < 0.05$) across treatments. Average weight gain ranged from 10.0 kg (SCM-10) to 13.12 kg (SCM-15). T3 recorded highest values for average feed intake (AFI) at a record value of 63.50 kg. T2 had the least feed conversion ratio of 4.81. The carcass quality and sensory evaluation were significantly affected ($P < 0.05$) across the treatments. Hot carcass weight

*Corresponding author: E-mail: davidolanrewaju_2008@yahoo.co.uk, okunlola@gmail.com;

(HCW) ranged from 18.90 kg (SCM-10) to 20.85 kg (SCM-15). Cold carcass weight was recorded 18.05 kg (SCM-10) and 19.95 kg (SCM-20), respectively. Minimum chilling loss of 3.96% was recorded at SCM-15 inclusion level. Animals on SCM-15 inclusion recorded the highest value of 7.55, 7.25 and 7.25 respectively for appearance, tenderness and overall acceptability. SCM-20 had highest value of 7.50 for juiciness compared to SCM-10 and SCM-15 with recorded values of 6.80 and 7.20, respectively. The study showed that all the dietary inclusion was satisfactory as feed for small ruminants without any deleterious effect on their performance as well as on carcass yield and meat quality. Inclusion level of SCM-15 supplement had highest overall performance when fed to (WAD) sheep.

Keywords: WAD sheep; growth response; carcass characteristics; meat quality; Shea cake meal.

1. INTRODUCTION

The major problem of ruminants in developing countries is inadequate supply of quality feed on a year round basis [1]. This is as a result of poor quality of tropical grasses, during the dry season [2]. Consequently, livestock production becomes very low both in quantity and quality, effect of which has resulted to inadequate consumption of protein of animal source to humans. Then the need for supplementation of natural forage with agro industrial by products such as residues from local farm as well as unconventional feed resources to improve livestock production cannot be overemphasized, considering the increase in human population with their corresponding demand for animal protein. This becomes necessary due to the high cost of conventional concentrates required to boost the performance of these animals. Small ruminants play significant role in animal protein production via meat and milk production, as well as a source of income to peasant farmers in rural communities. Improvements in performance of small ruminants using nutritious but yet untapped unconventional feed resource such as Shea cake is an effort in right direction. This crop discards have been studied to have potential benefits when used as a feed ingredient in livestock production. Shea tree (*Vitellaria paradoxa*) which bears shea fruit, grows naturally in African savannah. The fruit has a thin nutritious pulp around a large, oil-rich kernel which is used to make shea butter. The fruit is tasty and rich in vitamins and antioxidants [3]. The effluent of its fruit in the process of shea butter production solidifies; forming cake when dried. This cake is highly rich in nutrients, which instead of allowing it to be wasted and constitute nuisance can be utilized by the animals. This study therefore, aimed at investigating the potential of shea nut extracts as feed resource in the production and management of West African Dwarf (WAD) sheep.

2. MATERIALS AND METHODS

2.1 Experimental Site

The experiment was carried out at the small ruminant unit of Ladoke Akintola University of Technology, Ogbomoso, Oyo State. The institution is located on longitude 4°5' East of the Greenwich Meridian and at latitude of 8°7' North of the Equator in the derived savannah zone of Nigeria with a mean annual rainfall of 1247 mm and mean annual temperature of about 27°C. Laboratory analysis was carried out at the Animal Science Department laboratory, University of Ibadan.

2.2 Preparation of Experimental Diets

Solidified effluent (Shea cake) was collected from Shea butter production factory in Saki, a popular community in Okeogun Area of Oyo State. The shea cake alongside cassava peels and palm kernel cake were milled separately and mixed with other feed components, namely, wheat ofal, Di-calcium phosphate, ruminant premix and salt. A diet tagged Shea Cake Meal (SCM) was then formulated at the inclusion level of 0, 10, 15 and 20% of shea cake, respectively, as contained in the table of gross composition of the experimental diets. Samples of the experimental diets were taken and stored in a covered plastic container for laboratory analysis; following the procedures of AOAC [4].

2.3 Sample Collection

Samples were taken from four locations to ensure evenness and creates record for variation in the proximate if exists. The samples were collected from Ofiki, Ipapo, Ogbooro and Igbeti and each sample location represent a treatment before the best was selected and fed to the animal at varied levels.

2.4 Experimental Animal Management

A total of twenty four (24) growing West African dwarf sheep comprising of 12 males and 12 females weighing between 11–12.65 kg were used for the study. They aged between 5-6 months as dentition was used to estimate the age. The experimental animals were purchased from Okeogun region of Oyo State of Nigeria. On arrival, animals were given prophylactic treatments, using standard procedure of health management but were introduced to the experimental diets two weeks before prior to the commencement of experiment and data recording after weight balance for even distribution. Four experimental diets having Shea cake meal (SCM) at inclusion levels of 0, 10, 15 and 20% were offered to the experimental animals in individual pen in a completely randomized design of 6 replicates per treatment. The experimental animals were allowed to graze on guinea grass paddock for 1 hour every day, followed by shea cake meal supplement at 5% of their body weight. The orts was weighed at 08:00 am every morning and deducted from the quantity offered for intake determination prior to serving new feed daily. Individual animal was tagged for the purpose of identification. The feeding and relevant data collection lasted for 12 weeks.

2.5 Measurement of Growth

Weight of individual animal was measured at the onset of the trial and subsequently on weekly basis. Feed was offered at 5% of body weight of individual experimental animals. The intake and orts were measured daily to determine weekly average intake. Weights were taken weekly to determine growth change of the animals. Data collection lasted for 84 days.

2.6 Slaughtering, Processing of Carcass and Sensory Evaluation

At 84th day, representatives from each treatment were randomly selected and weighed to obtain slaughter weight (SW), and thereafter slaughtered. After bleeding, the skin, head and limbs were removed before the carcass was eviscerated. The hot carcass weight (HCW) was obtained before the carcasses were taken to the cold chamber for 24 hours at 4°C for cooling to obtain cold carcass weight (CCW) and chilling losses (CL) was quantified using the formula of Maciel et al. [5].

$$CL (\%) = \frac{HCW - CCW}{HCW} \times 100$$

However, hot carcass yield (HCY) and cold carcass yield (CCY) were computed using the formulas of Urbano et al. [6].

$$HCY (\%) = \frac{HCW}{SW} \times \frac{100}{1}$$

$$CCY (\%) = \frac{CCW}{SW} \times \frac{100}{1}$$

2.7 Sensory Evaluation

Cooked meat from loin chops were served in plates to a 12 member taste panel drawn from the staff and students population to judge the test for appearance, flavor, tenderness, juiciness and overall acceptability of meat samples on a 9 - point hedonic scale where (1) corresponds to extremely dislike and (9) to extremely like. Panelists were trained prior to the evaluation test.

2.8 Statistical Analysis

Data obtained were subjected to analysis of variance (ANOVA) using the procedure of SAS [7] package to determine the effect of dietary treatments on the various parameters studied. Significant means were separated using Duncan multiple range test of the same software at 0.05 significant levels.

3. RESULTS AND DISCUSSION

Table 1 presents the proximate composition of shea cake taken from four locations in the Derived Savannah Area of Nigeria, since each location represent a treatment. The dry matter value ranges from 87.02 – 89.65%; with the sample T3 (Sample from Ogboro) recording the highest value (89.65%). T2 (Ipapo) recorded the least dry matter of 87.02%. The least value recorded by T4 was just in comparison among samples from other location as its value (87.02%) and not of less quality for use in ruminant feed composition or formulation. The dry matter value across the treatment is an indication of good shelf life and storage value of shea cake. Crude protein (CP) values on the other hand recorded a value range of 7.00 (T2) - 8.75 (T3). This was in agreement with the findings of Enaberue et al. [8] on shea fruit pulp. The CP values recorded justified shea cake as a good feed resource in ruminant nutrition. The ether extract (EE) recorded was high across

treatments. The values ranged from 19.20% (T4) – 22.00% (T2). This could be linked to the nature of shea tree (*Vitellaria paradoxa*); being an oil rich tree. Comparatively, Shea cake from Ogbooro (T3) had best results among other treatments, and was considered for use in this study at various inclusion levels.

Table 2 presents the gross composition of experimental diets. Shea cake at the inclusion level of 0, 10, 15 and 20% was mixed with other feed ingredients. These include wheat offal, Di-calcium phosphate, ruminant premix and salt to form experimental diets tagged shea cake meal (SCM). The change in body weight of the experimental animals were significantly different ($P < 0.05$) across treatments (Table 3). Animals

on all treatments recorded appreciable values for intake and weight gain with animals on 15% (SCM-15) having the highest value of average weight gain of 13.12 kg compared to T1, T2 and T4. The feed intake recorded across treatments reflects on the weight gained by the animals, as well as feed conversion ratio was an indication of good efficiency and availability of nutrients embedded in SCM which was maximally utilized by the animals. This result agreed with the findings of Okunlola [9] where Baobab fruit meal supplement was fed to Red Sokoto sheep with highest weight gain of 13.3kg and Aina [10], where the best efficiency in goat was traced to feed supplementation. Results obtained in this study shows that shea cake meal supplement improved the performance of WAD sheep.

Table 1. Proximate composition of Shea Cake (SK) in the Derived Savannah Area of Nigeria

Parameters (%)	Ofiki (T1)	Ipapo (T2)	Ogbooro (T3)	Igbeti (T4)
Dry matter	89.15 ^b	88.50 ^d	89.65 ^a	87.02 ^c
Crude protein	7.60 ^b	7.00 ^c	8.75 ^a	7.26 ^{ab}
Crude fibre	4.00 ^b	4.00 ^b	4.50 ^a	4.22 ^{ab}
Ether extract	20.00 ^b	22.00 ^a	21.50 ^a	19.20 ^c
Ash	4.55 ^b	4.60 ^b	5.05 ^a	4.30 ^c
Nitrogen free extract	53.00 ^a	50.90 ^b	49.85 ^c	52.04 ^a

^{abcd} Means within each row with different superscripts are significantly different ($P < 0.05$)

Table 2. Gross composition of experimental diets

Component/Treatment	T ₁ (SCM-0)	T ₂ (SCM-10)	T ₃ (SCM-15)	T ₄ (SCM-20)
Shea cake	0.00	10.00	15.00	20.00
Salt	1.00	1.00	1.00	1.00
Ruminant premix	0.50	0.50	0.50	0.50
Di-calcium phosphate (DCP)	0.50	0.50	0.50	0.50
Palm Kernel Cake (PKC)	5.00	5.00	5.00	5.00
Wheat offal	18.00	18.00	18.00	18.00
Cassava peel	75.00	65.00	60.00	55.00
Total	100.00	100.00	100.00	100.00

SCM-0: diet without Shea cake, SCM-10: diet with 10% Shea cake, SCM-15: diet with 15% Shea cake, SCM-20: diet with 20% Shea cake.

Table 3. Intake, growth and feed conversion ratio of West African Dwarf sheep Shea cake meal

Treatment/Parameter	Treatments				SEM
	T ₁ (SCM-0)	T ₂ (SCM-10)	T ₃ (SCM-15)	T ₄ (SCM-20)	
Initial body weight (kg)	12.50 ^a	11.00 ^b	12.65 ^a	11.80 ^b	1.50
Final body weight (kg)	22.50 ^c	23.90 ^b	25.77 ^a	23.85 ^b	1.26
Average feed intake (kg)	60.20 ^c	62.00 ^b	63.50 ^a	60.75 ^c	5.00
Average weight gain (kg)	10.00 ^c	12.90 ^b	13.12 ^a	12.05 ^{ab}	1.05
Feed conversion ratio (FCR)	6.02 ^a	4.81 ^b	4.84 ^b	5.04 ^b	0.75

^{abcd} Means within each row with different superscript are significantly different ($P < 0.05$)

SCM-0: diet without Shea cake, SCM-10: diet with 10% Shea cake, SCM-15: diet with 15% Shea cake, SCM-20: diet with 20% Shea cake.

Table 4. Carcass characteristics and sensory evaluation of West African Dwarf sheep fed Shea cake meal

Parameters	Treatment diets				SEM
	SCM-0	(SCM-10)	(SCM-15)	(SCM-20)	
Carcass characteristics					
Slaughter weight (kg)	22.50 ^c	23.90 ^b	25.77 ^a	23.85 ^b	1.05
Hot carcass weight (kg)	17.65	18.90 ^b	20.85 ^a	18.95 ^b	0.75
Cold carcass weight (kg)	16.48	18.05 ^b	19.95 ^a	18.20 ^b	0.05
Chilling losses (%)	6.63	4.50 ^a	4.32 ^a	3.96 ^b	0.55
Hot carcass yield (%)	78.44	79.08 ^b	80.91 ^a	79.45 ^a	1.00
Cold carcass yield (%)	73.24	75.52 ^c	77.42 ^a	76.41 ^b	0.75
Sensory evaluation					
Appearance	6.00 ^c	7.20 ^b	7.55 ^a	7.50 ^a	0.25
Flavour	5.00 ^b	6.95 ^a	7.00 ^a	7.00 ^a	0.35
Tenderness	5.50 ^d	6.50 ^c	7.25 ^a	7.00 ^b	0.45
Juiciness	5.25 ^d	6.80 ^c	7.20 ^b	7.50 ^a	0.20
Overall acceptability	5.00 ^c	6.00 ^b	7.25 ^a	7.00 ^a	0.45

^{abcd} Means with different superscripts along the same row are significantly different ($P < 0.05$). SEM = standard error of mean. SCM-0: diet without Shea cake, SCM-10: diet with 10% Shea cake, SCM-15: diet with 15% Shea cake, SCM-20: diet with 20% Shea cake.

Table 4 presents carcass characteristics and sensory evaluation of West African Dwarf sheep fed shea cake meal. There were no significant difference ($P > 0.05$) in the hot carcass and cold carcass weights of T2 (10%) and T4 (20%). Experimental animals on control diet (SCM-0) recorded lowest values in all parameters evaluated. The Hot Carcass Weight (HCW) was highest at T3 (SCM-15) with record value of 20.85 kg. This was higher than the value obtained by McGregor [11], 13.4 kg for angora sheep with a live weight of 30 kg, and carcass weight of 14 kg in a study which involved grazing Saanen (diary) sheep. Meat consumers generally prefer lean meat. Butchers need consistent products that always meet specification, while producers of sheep for meat are in the business of growing saleable carcasses. According to the result obtained in this study, tissue composition highly affects commercial quality of the carcass at various level of SCM inclusion. This makes the experiment suitable as feed supplement in goat production for meat purpose. Study have shown that goat have 69.50% muscles and 11.8% bones [12]. SCM-15 in this study recorded the highest value of tenderness (7.25), appearance (7.55), and overall acceptability (7.25). Appearance (colour) is an important attribute that affects consumer purchase and willingness to buy, Tenderness often affects consumer acceptability and satisfaction, the metabolic-biochemical reaction that happen after rigor mortis let a progressive tenderization [11] McGregor 1985). The sensory evaluation in this

study clearly justified shea cake meal as a good supplement to promote quality of WAD sheep mutton. Worthy of note of the meat quality in this study was appreciable reduction in odour associated with mutton with increase in level of SCM inclusion. This further led to high level of acceptability especially at T3 (SCM-15) inclusion level. This study has shown that an increase in shea cake meal supplement up to 20% of the total diet improves the carcass quality in terms of carcass yield and overall acceptability of West African Dwarf sheep.

4. CONCLUSION AND RECOMMENDATION

The results from this study showed Shea cake meal as a profitable supplement in WAD sheep diets as there was no deleterious effect on the performance of the experimental animals. Shea cake meal promotes rapid growth and as such makes it appropriate in fattening program. Its inclusion in ruminants feed will supply the nutrients required for healthy performance. Its uses would reduce cost of production because it is available free at shea butter production factory. Its use would reduce environmental pollution; thereby enhance promotion of health in humans. Supplementing WAD sheep feed with SCM increased the acceptability, table and market value of mutton. Its use in this experiment removed bias associated with mutton due to its odour which repels some consumers, thereby making them reject its consumption.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Makkar HPS. Recent advances in vitro gas method for evaluation of nutritional quality of feed resources. Animal Production and Health Section, Joint FAO/IAEA Division, International Atomic Energy Agency Publications. 2004;1-10.
2. Olorunnisomo OA. Sweet potato as a ruminant feed: performance of sheep fed mixtures of the forage and root. Nigerian Journal of Animal Production. 2008;35(2): 242-251.
3. Olorode BR, Onifade AA, Babatunde GM. Comparative Utilization of Shea butter Cake and Palm Kernel Cake by Broiler Chicken. Nigerian Journal of Animal Production. 1997;24:124-131.
4. AOAC. Official methods of analysis of the association of official's analytical chemists, 17th Edn. Association of official analytical chemists, Arlington, Virginia; 2003.
5. Maciel MV, Carvaliho FFR, Batista GA, Souza EJO, Maciel LPA, Pereira –Neto JD, Lima–Junior DM. Carcass and non – carcass characteristics of sheep fed cassava (*Manihot pseudoglaziovii* Pax and Hoffm, K.) Chilean Journal of Agricultural Research. 2015;75(3):307–312.
6. Urbano SA, Ferrera MA, Wilson MD, Andrade RP, Michelle CB, Sabrina CR. Carcass Characteristics of sheep fed with Castor bean hulls in replacement of Tifton Hay. Cienc Agrotech Lauras. 2013;36(1): 85-93.
7. SAS. Statistical Analysis Systems, User'Guide, Version 8 ed., SAS Institute Inc. SAS Campus Drive Cary, North Carolina, USA; 2003.
8. Enaberue LO, Obisesan IO, Okolo EC, Ojo A. Proximate and chemical composition of shea (*Vitellaria paradoxa* C.F. Gaertn) fruit pulp in the Guinea Savanna of Nigeria. World Journal of Agricultural Sciences. 2014;2(4):078-083.
9. Okunlola DO. Growth and Milk Production of Red Sokotosheep fed Baobab (*Adansonia digitata*) fruit meal based diets. Ph.D Thesis, Dept. of Animal Science, University of Ibadan, Nigeria; 2016.
10. Aina ABJ. Effect of dietary salt levels on the performance of West African Dwarf sheep. Nigerian Journal of Animal Production. 2002;29(1):47-54.
11. McGregor BA. Growth development and carcass composition of sheep. A review in goat production and research in the Tropic, Copeland, Australia Centre for International Agricultural Research, Canberra. 1985;82-90.
12. Ana G, Maria de Mar C, Jose LO, Carlos S. Carcass and Meat Quality in Goat, Goat Science, Sandor Kukovics, Intech Open; 2017. Available: <http://www.intechopen.com/book/goat-science/carcass-and-meat-quality-in-goat>

© 2021 Okunlola et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/77197>