



Efficacy and Phytotoxicity of Triafamone 18.52% SC in Direct Sown Rice and their Residual Effect on Succeeding Blackgram

B. Jyothi Basu ^{a*}, P. Swathi ^a, N. Sambasiva Rao ^a
and V. Saida Naik ^a

^a Agricultural Research Station, Jangamaheswarapuram, Acharya N.G Ranga Agricultural University, Guntur District, Andhra Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i203928

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106243>

Original Research Article

Received: 08/07/2023
Accepted: 14/09/2023
Published: 11/10/2023

ABSTRACT

Rice (*Oryza sativa*), the staple food of more than half of the population of the world, is an important target to provide food security and livelihoods for millions. Direct seeding of rice (DSR) refers to the process of establishing the crop from seeds sown in the field rather than by transplanting seedling from the nursery. Therefore, field experiment was conducted on clay loam soils at Agricultural Research Station, Jangamaheswarapuram, Guntur Dist. Andhra Pradesh, India, during 2017-18 and 2018-19 in direct sown rice (DSR) to test the efficacy of Triafamone 18.52 SC herbicide on production and to evaluate the lasting impact of herbicides on succeeding blackgram. Among the herbicides tested only Triafamone 18.52 SC @ 100 g a.i. ha⁻¹ had a slight phytotoxic effect on direct sown rice. Phytotoxicity of Triafamone 18.52 SC @ 100 g a.i. ha⁻¹ applied at 2-3 leaf stage of weed effect the rice crop, exhibiting slightly stunted growth and the leaves failed to fully expand and

*Corresponding author: E-mail: jyoba226@gmail.com;

became yellow at 7 days after application. However, those symptoms disappeared, and the Rice plants recovered within a week. Further, it was observed that none of the herbicides applied at tested rates had adverse effect on succeeding blackgram. In among the herbicides Triafamone 18.52 SC @ 100 g a.i. ha⁻¹(T₅) and Triafamone 18.52 SC @ 50 g a.i. ha⁻¹(T₄) applied at 2-3 leaf stage of weed were at par recorded significantly higher grain yield over weedy check (T₁).

Keywords: Rice; phytotoxicity; blackgram; DSR.

1. INTRODUCTION

"The primary crop grown in the tropics, particularly in India, is rice. The phrase "Rice is Life" perfectly captures the significance of rice in the food and nutritional security of Asian nations. India is the second-largest producer of rice in the world, with a production of 118.4 million tonnes and a productivity of 2.7 t ha⁻¹ on an area of 43.8 million hectares" [1]. "With a yield of 8.23 million tonnes and a productivity of 3.73 t ha⁻¹, it is grown on 2.21 million hectares in Andhra Pradesh [2]. Before the advent of Green revolution and adoption of irrigation, rainfed rice was often broadcasted into moist soil and yields were low, variable and highly prone to weed competition. Weed spectrum and degree of infestation in rice field are often determined by rice ecosystems and establishment methods. Research evidences at different places has shown around 20-100% losses due to weeds such as *Echinochloa spp.*, *Leptochloa spp.*, *Cyanotis spp.*, *Commelina sp.*, *Digitaria spp.* and *Alternanthera sp* in DSR. Integrated weed management approach based on the critical period of crop weed competition, involving different direct and indirect control measures, has been developed and widely adopted by farmers to overcome weed problem in DSR in a sustainable way. Weed infestation is the major biotic constraint for higher productivity especially in dry direct sown rice (DSR)" [3-5]. "Herbicides are frequently employed in high-input crop production systems to suppress weeds. Yet, the majority of the unused fractions of herbicides may stay within soils. Weed competition reduced multiple rice yield components, and weed biomass in wet-seeded rice was six-fold greater than in rice transplanted into puddled soil and twice as much again in dry-seeded rice sown either after dry tillage or without tillage" [6]. "Hence, information regarding persistence and residual effect of herbicides in soil is essential to use them safely and effectively. Because of this, bioassay continues to be a crucial technique for both qualitative and quantitative analysis of herbicide residue in soil" [7]. In light of the aforementioned facts, an effort has been made to

assess the Triafamone 18.52% SC efficacy and phytotoxicity in direct sown Rice and their residual effect on succeeding Blackgram.

2. MATERIALS AND METHODS

A field experiment was conducted on clay loam soils of at Agricultural Research Station, Jangamaheswarapuram, Guntur Dist. Andhra Pradesh, India for two consecutive years (2017 - 18 and 2018-19). The soils of the experimental locations were medium to deep black texture with PH 7.6, organic carbon 0.57, available Nitrogen 228 kg/ha, available Phosphorus 47.8 kg/ha, and available Potassium 310.4 kg/ha. The experiment was laid out in randomized block design with nine treatments.

Triafamone belongs to the keto sulfonamide herbicides. In plants, Triafamone is taken up by leaves and roots and is very quickly converted into an intermediate form by reduction of the keto group. Contrary to rice, in weeds a 2nd metabolite is formed by N-demethylation, and this metabolite strongly inhibits acetolactate synthase (ALS). Phytotoxic effect of herbicides on Rice crop, if any, was assessed at 7, 14, 21, 28 and 35 days after spraying by using simple rating scale of 0-10 (equal to 0 to 100%) [8] as given in Table 2.

After harvest and threshing of crop, grain yield was recorded in net plot wise and converted to grain yield per hectare. The data of each year was analyzed separately. Weed index indicates percent reduction in grain yield due to weed competition. The weed index was calculated by using the formula as suggested by Gill and Vijay Kumar (1969).

$$WI (\%) = \frac{X-Y}{X} \times 100$$

Where,

X = Grain yield from weed free plot
Y = Grain yield from plots for which WI is to be calculated.

Table 1. Treatments, dose and time of application

Treatment	Dose (g ha ⁻¹)	Time of Application
T ₁ :Untreated control	-	-
T ₂ :Triafamone 18.52 SC	30	2 to 3 leaf stage of weed
T ₃ :Triafamone 18.52 SC	40	2 to 3 leaf stage of weed
T ₄ :Triafamone18.52 SC	50	2 to 3 leaf stage of weed
T ₅ :Triafamone18.52 SC	100	2 to 3 leaf stage of weed
T ₆ :Pyrazosulfuron ethyl 10% WP	15	2 to 3 leaf stage of weed
T ₇ :Cyhalofop Butyl 10% EC	80	2 to 3 leaf stage of weed
T ₈ :Farmer practice (two hand weedings)	-	20 DAS and 40 DAS
T ₉ :Weed free	-	-

Table 2. Phytotoxicity Rating Scale (PRS)

Effect	Rating	Description on crop	Crop response/ Crop injury
None	0	No injury, normal	0-00
Slight	1	Slight stunting injury or discolouration	1-10%
	2	Some stand loss, stunting discolouration	11-20%
	3	Injury more pronounced but not persistent	21-30%
Moderate	4	Moderate injury, recovery possible	31-40%
	5	Injury more persistent, recovery doubtful	41-50%
	6	Near severe injury, no recovery possible	51-60%
Severe	7	Severe injury, stand loss	61-70%
	8	Almost destroyed, a few plants surviving	71-80%
	9	Very few plants alive	81-90%
Complete	10	Complete destruction	91-100%

The observations on growth attributes were taken at different intervals and yield attributing characters like no of panicles m⁻², panicle length, no of grains per panicle and test weight were recorded at harvest. The data was analyzed using ANOVA and the least significant difference (LSD) values at 5% level of significance were calculated and used the RBD test significant difference between treatment means.

3. RESULTS AND DISCUSSION

3.1 Crop Injury Score

Among the herbicides tested only T₅ (Triafamone 18.52 SC @ 100 g a.i. ha⁻¹ at 2-3 leaf stage of weed) had a slight phytotoxic effect on Rice. Phytotoxicity of Triafamone 200 SC @ 100 g a.i. ha⁻¹ was characterized by slightly stunted plant growth and leaves fail to expand fully and became yellowish as observed at 7 days after application. However, those symptoms disappeared, and the Rice plants recovered within a week. Similar trend was close conformity of Mohapatra et al. [9] and Murali Arthanari [10].

3.2 Grain Yield (kg ha⁻¹)

Grain yield of Rice was significantly influenced by different weed management practices in both the years of study (Table 4).

The highest grain yield (5076 and 5651 kg ha⁻¹ during 2017-18 and 2018-19, respectively) was recorded under weed free treatment (T₉), which was significantly superior to rest of the treatments except treatments T₈ (Two hand weedings), which was however, comparable to the treatments T₅ (Triafamone 18.52 SC @ 100 g a.i. ha⁻¹ at 2-3 leaf stage of weed) and T₄ (Triafamone 18.52 SC @ 50 g a.i. ha⁻¹ at 2-3 leaf stage of weed). The lowest grain yield (1922 and 2552 kg ha⁻¹) was obtained in untreated (T₁) plot, which was significantly lower than any herbicidal treatment. The results are conformity with the findings of Kumar et al., [11], Mohapatra et al. [9] and Murali Arthanari [10].

3.3 Phyto-toxicity Effect on Succeeding Crop

The data on succeeding crop (Blackgram) did not show any crop injury as well as there was no adverse effect on its germination, plant height, branches and yield of crop.

3.4 Seed Yield of Blackgram (kg ha⁻¹)

The seed yield of succeeding blackgram crop after rice was nonsignificant among the treatments during both the years of study [Table 6]. This indicates that there was no marked

Table 3. Phytotoxic effect of Triafamone 18.52 SC on direct seeded Rice as influenced by weed management practices during Rabi, 2017-18 and Kharif, 2018-19

Treatments	Dose (g a.i. ha ⁻¹)	Phyto-toxicity Days after herbicide application									
		2017-18					2018-19				
		7	14	21	28	35	7	14	21	28	35
T ₁ .Untreated (control)	-	-	-	-	-	-	-	-	-	-	-
T ₂ . Council Prime (Triafamone 18.52 SC)	30	0	0	0	0	0	0	0	0	0	0
T ₃ . Council Prime (Triafamone 18.52 SC)	40	0	0	0	0	0	0	0	0	0	0
T ₄ . Council Prime (Triafamone 18.52 SC)	50	0	0	0	0	0	0	0	0	0	0
T ₅ .Council Prime (Triafamone 18.52 SC)	100	1	0	0	0	0	1	0	0	0	0
T ₆ .Pyrazosulfuron ethyl 10% WP	15	0	0	0	0	0	0	0	0	0	0
T ₇ .Cyhalofop Butyl 10% EC	80	0	0	0	0	0	0	0	0	0	0
T ₈ .Farmer practice (two hand weedings)	-	-	-	-	-	-	-	-	-	-	-
T ₉ .Weed free	-	-	-	-	-	-	-	-	-	-	-

Table 4. Grain yield and weed index of direct seeded Rice as influenced by weed management practices during Rabi, 2017-18 and Kharif, 2018-19

Treatments	Dose (g a.i. ha ⁻¹)	Grain yield (kg ha ⁻¹)		*Weed index (%)	
		2017-18	2018-19	2017-18	2018-19
T ₁ .Untreated (control)	-	1922	2552	52.07 (62.1)	48.27 (55.6)
T ₂ . Council Prime (Triafamone 18.52 SC)	30	3440	3667	34.48 (32.1)	36.37 (35.5)
T ₃ . Council Prime (Triafamone 18.52 SC)	40	3751	4143	30.70 (26.7)	31.01 (26.8)
T ₄ . Council Prime (Triafamone 18.52 SC)	50	3993	4460	27.68 (22.0)	27.07 (21.3)
T ₅ .Council Prime (Triafamone 18.52 SC)	100	4376	4730	21.80 (14.4)	22.83 (16.4)
T ₆ .Pyrazosulfuron ethyl 10% WP	15	3226	3537	37.09 (36.5)	37.64 (37.4)
T ₇ .Cyhalofop Butyl 10% EC	80	3492	3713	33.83 (31.4)	35.93 (34.6)
T ₈ .Farmer practice (two hand weedings)	-	4649	5185	16.82 (8.8)	13.90 (7.7)
T ₉ .Weed free	-	5076	5651	0.00 (0.0)	0.00 (0.0)
SEm_±	-	168.27	177.16	2.06	2.68
CD (P = 0.05)	-	491.13	517.09	6.03	7.82

*Note: Data transformed to arc sine transformations. Figures in parenthesis are original values

Table 5. Phytotoxic symptoms and germination of succeeding Blackgram as influenced by Triafamone 18.52 SC during Rabi, 2017-18 and Kharif, 2018-19

Tr. No	Treatment	Dose (g a.i.ha ⁻¹)	Phyto-toxicity (Days after sowing)				Plant germination per 3 row at 20 DAS	
			2017-18		2018-19		2017-18	2018-19
			20	Harvest	20	Harvest		
T ₁ .	Untreated control	-	0	0	0	0	47.5	48.1
T ₄ .	Council Prime (Triafamone 18.52 SC)	50	0	0	0	0	47.3	47.1
T ₅ .	Council Prime (Triafamone 18.52 SC)	100	0	0	0	0	46.1	46.3

Table 6. Plant height (cm), braches (No. plant⁻¹) and seed yield (kg ha⁻¹) of Blackgramas influenced by weed management practices in Rice-Blackgram sequence during Rabi, 2017-18 and Kharif, 2018-19

Treatments	Dose (g a.i. ha ⁻¹)	Plant height		No of branches plant ⁻¹		Seed yield (kg ha ⁻¹)	
		2017-18	2018-19	2017-18	2018-19	2017-18	2018-19
T ₁ . Untreated control	-	31.98	34.83	7.17	7.73	506	549
T ₂ . Council Prime (Triafamone 200 SC)	30	33.41	34.76	7.18	7.95	520	547
T ₃ . Council Prime (Triafamone 200 SC)	40	34.90	35.33	7.06	7.39	525	547
T ₄ . Council Prime (Triafamone 200 SC)	50	34.86	35.89	7.26	8.00	557	561
T ₅ . Council Prime (Triafamone 200 SC)	100	36.12	36.59	8.02	8.17	566	574
T ₆ . Pyrazosulfuron ethyl 10% WP	15	34.33	36.35	6.97	7.80	545	560
T ₇ . Cyhalofop Butyl 10% EC	80	32.87	34.95	7.44	8.44	534	558
T ₈ . Farmer practice (two hand weedings)	-	34.96	34.99	8.27	8.81	560	569
T ₉ . Weed free	-	36.33	35.88	7.78	8.06	555	580
SEm±	-	0.95	0.52	0.35	0.33	14.07	8.24
CD (P = 0.05)	-	NS	NS	NS	NS	NS	NS

difference among the treatments and the impact of herbicides applied to rice. The applied herbicides left no lingering effects on the germination, plant height, number of branches per plant or seed yields of blackgram after they had adequately broken down in the soil. This extraordinary manifestation suggests that the various weed management techniques used on rice had no discernible impact on the development and productivity of the next blackgram crop. Jyothi Basu et al., [7] found similar findings, stating that herbicides applied to rice crops had no lasting effects on subsequent crops' growth and production. These findings were in agreement with the views expressed by Deivasigamani [12], Deivasigamani [13], Murali Arthanar [10].

4. CONCLUSIONS

On the basis of foregone studies, it is concluded that, Triafamone 18.52 SC was not shown any phytotoxicity on Rice crop and Triafamone 18.52 SC applied in Rice doesn't have any residual effect either positive or negative on growth and yield of succeeding Blackgram crop. Further, Triafamone 18.52 SC @ 100 g a.i. ha⁻¹ at 2-3 leaf stage of weed (T₅) followed by T₄ (Triafamone 18.52 SC @ 50 g a.i. ha⁻¹ at 2-3 leaf stage of weed) may be used for attaining effective weed management, maximum rice grain yield with no residual effect on succeeding blackgram.

ACKNOWLEDGEMENT

The facilities provided by the Acharya N. G. Ranga Agricultural University and the financial support provided by Ms. Bayer crop science pvt. Ltd., India are gratefully acknowledged by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. GOI (Government of India). Economic Survey 2020-21. Statistical Appendix. Ministry of Finance. Government of India, New Delhi. 2021;2. Available:<https://www.indiabudget.gov.in/economicsurvey/>
2. Reserve Bank of India. Handbook of Statistics on Indian states. Reserve Bank of India, New Delhi, India; 2020.
3. Jyothi Basu B, Prasad PVN, Murthy VRK, Ashoka Rani Y, Prasad PRK. Productivity of direct seeded rice in response to various weed management practices and their residual effect on greengram. *Journal of Rice Research*. 2020^a;13(1):66-74.
4. Jyothi Basu B, Prasad PVN, Murthy VRK, Ashoka Rani Y, Prasad PRK. Evaluation of sequential application of herbicides for controlling complex weed flora in direct sown rice. *Indian Journal of Plant Protection*. 2020^b;48(4):482-490.
5. Jyothi Basu B, Prasad PVN, Murthy VRK, Ashoka Rani Y, Prasad PRK. Efficacy of sequential application of herbicides on weed management, rice nutrient uptake and soil nutrient status in dry direct-seeded ricegreengram sequence. *Indian Journal of Weed Science*. 2021;53(4):398-404.
6. Singh VP, Singh SP, Tripathi N, Kumar A and Banga A. Studies on bioefficacy and phytotoxicity of penoxsulam for weed control in direct seeded rice. In: Biennial Conference of Indian Society of Weed Science on "Weed Threat to Agriculture, Biodiversity and environment. 19-20 April, 2012, Kerala Agriculture University, Thrissur, India. 2012;72.
7. Jyothi Basu B, Prasad PVN, Murthy VRK, Ashoka Rani Y, Prasad PRK. Bioefficacy and phytotoxicity of herbicides in Rice and their residual effect on Succeeding greengram. *International Journal of Agriculture Sciences*. 2020^c;12(11):9940-9944.
8. Rao VS. Principles of Weed Science, Second Edition Oxford & IBH Publishing Company Private Limited, New Delhi. 2000;127-142.
9. Mohapatra S, Tripathy SK, Tripathy S, Mohanty AK. Effect of sequential application of herbicides on productivity and profitability of transplanted rice. *Indian Journal of Weed Science*. 2021;53(2):129-134.
10. Murali Arthanari P. Weed management with Triafamone herbicide in transplanted rice ecosystem. *Emirates Journal of Food and Agriculture*. 2023;35(4):351-356.
11. Kumar S, Rana SS, Chander N, Ramesh. Mixed weed flora management by Bispyribac sodium in transplanted rice. *Indian Journal of Weed Science*. 2013; 45(3):151-155.
12. Deivasigamani S. Bio-efficacy and phytotoxicity of triafamone and ethoxysulfuron in transplanted Rice (*Oryza*

- sativa*). International Journal of Multidisciplinary Research and Modern Education. 2016^a;2(1):274-279.
13. Deivasigamani S. Study of bio-efficacy and Phytotoxicity of New Generation Herbicides on Triafamone and Ethoxysulfuron in Direct Seeded Rice (*Oryza sativa*). IRA-International Journal of Applied Sciences. 2016^b;3(2):106-112.

© 2023 Basu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/106243>