



# Diversity and Abundance of Butterflies (Lepidoptera: Rhopalocera) in Sethu Bhaskara Agricultural College and Research Foundation, Karaikudi, Tamil Nadu, South India

A. Narmadha <sup>a++</sup>, R. Ravinasree <sup>a++</sup>,  
S. R. Sakthi Kayalvizhi <sup>a++</sup>, M. Mathialagan <sup>a#\*</sup>,  
Kishan Tej Mitta <sup>bt</sup>, R. Meena <sup>a#</sup>, V. Sarathkumar <sup>a#</sup>,  
S. Kumaravel <sup>a#</sup>, S. Lekha Priyanka <sup>c#</sup>,  
S. Mangayarkarsi <sup>a#</sup> and U. Ganesh <sup>a++</sup>

<sup>a</sup> Sethu Bhaskara Agricultural College and Research Foundation, Karaikudi, Tamil Nadu, India.

<sup>b</sup> Department of Entomology, SMGR Agricultural College, ANGRAU, Udayagiri- 524 226,  
Andhra Pradesh, India.

<sup>c</sup> Pushkaram College of Agricultural Sciences, Pudukkottai-622 303, Tamil Nadu, India.

## Authors' contributions

This work was carried out in collaboration among all authors. Author AN designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Author RR and Author SRSK managed the analyses of the study. Author SRSK managed the literature searches. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2023/v13i102933

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:  
<https://www.sdiarticle5.com/review-history/106145>

Original Research Article

Received: 02/08/2023

Accepted: 05/09/2023

Published: 08/09/2023

<sup>++</sup>UG Student;

<sup>#</sup>Assistant Professor;

<sup>†</sup>Assistant Professor and Head;

<sup>\*</sup>Corresponding author: E-mail: mathialagan08637@gmail.com;

## ABSTRACT

Butterflies play a vital role in the ecosystem, acting as pollinators, pollution indicators, pests, predators, weed killers, a good source of food and being of aesthetic, economic and ecological importance. World-wide butterfly populations are on the decline due to habitat destruction and deforestation. This study was taken with the prime objective of preparing the checklist of butterflies at Sethu Bhaskara Agricultural College and Research Foundation, Karaikudi, Tamil Nadu, from April 2023 to July 2023. A total of 976 individuals of 76 species belonging to 5 families have been recorded in the survey. For this study, the college was mapped into four different habitats: botanical garden area, agricultural field area, horticultural field area and grassland area. Results revealed that the family Nymphalidae showed the maximum number of species (22.0 species) from 12 genera (29.0%), followed by Lycaenidae (20.0 species) from 17 genera (26.5%), Pieridae (18.0 species) from 9 genera (24.0%), Hesperidae (12.0 species) from 11 genera (16.0%) and the least being Papilionidae (4.0 species) from 2 genera (5.5%). Among the four sites, the botanical garden area ranks first with the highest number of butterflies (140 species, N = 280), followed by the horticultural field area (129 species, N = 258), the agricultural field area (115 species, N = 230) and the grassland area (104 species, N = 208). According to the monthly distribution of butterflies, the maximum number of species and abundance were recorded in the months of July (307 species, N = 332), followed by June (141 species, N = 282), April (112 species, N = 224) and least in May (69 species, N = 138). The overall Shannon-Weiner diversity index was  $H = 1.50$ , Pielou's evenness index was  $E = 0.60$  and Margalef's index was  $R = 13.30$ . This study illustrated useful information on butterfly diversity in the study region, which serves as a baseline for future monitoring programs.

**Keywords:** Butterfly; conservation; diversity; ecological indicator; nymphalidae.

## 1. INTRODUCTION

Butterflies (Lepidoptera: Rhopalocera) are fascinating, colorful and the most studied insects in the world [1]. Butterflies occupy a vital position in the ecosystem as keystone species that act as valuable pollinators and help more than 50 economically important crops in the pollination process and the food chain and food web [2,3,4], biological and pollution indicators of terrestrial ecosystem [5], nutrient recycling (N, P, K) and decomposition [6], biological pest control [7], good source of food and have aesthetic, economic and ecological importance [8,9,10]. Butterflies can be used as a good biological indicator for vegetative structure, habitat quality [11,12], climate change, environmental health and degradation [7], and the impact of different threats due to anthropogenic disturbances [13]. Approximately 28,000 species of butterflies are recorded in the world [14]. Of these, 1504 species of butterflies are recorded in the Indian subcontinent, which is about 8.74% of the total butterfly species in the world and constitutes 65.0% of the total Indian fauna [14,15]. The number of Indian butterflies amounts to one-fifth of the world's species [16,17]. Among which, Tamil Nadu hosts about 323 species of butterflies [18,19]. Butterflies are classified into two superfamilies, viz., Hesperioidea and Papilionoidea. Hesperioidea consists of a single

family of Hesperidae (Skippers), whereas Papilionoidea consists of the rest of the butterfly families, viz., Papilionidae (Swallowtails), Pieridae (Whites and Sulphurs), Nymphalidae (Brush-footed butterflies) and Lycaenidae (Blues) [20]. The recent worldwide trend is that anthropogenic disturbances like urbanization, deforestation, habitat destruction, industrialization, and climate change, along with agricultural intensification like the use of pesticides and herbicides, cause migration or local extinction of butterflies [21-24]. In the last 40 years, there has been a dramatic decline of about 35.0% in butterfly abundance across the globe [25] and due to anthropogenic drivers of defaunation, many species (about 40.0%) may face extinction in the next few decades [26]. Studies on the diversity of butterflies in any area help understand the status of an ecosystem [27]. Studies related to butterfly diversity are necessary to prevent extinction and further decline, particularly in urban regions and educational institutional campuses, where they are very rare [28] and extremely necessary for effective and proper conservation of butterflies [29]. However, no previous study has been carried out on the butterflies' diversity at Sethu Bhaskara Agricultural College and Research Foundation (SBAC & RF), Karaikudi, Sivagangai district. Hence, the present study was the first preliminary attempt to document the

species diversity and abundance of butterflies in SBAC and RF, Karaikudi. The result of the survey will be helpful in making a butterfly checklist and conservation plan for the study area.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

The study was conducted at Sethu Bhaskara Agricultural College and Research Foundation, Karaikudi, Sivagangai district, Tamil Nadu, India (Fig. 1). The college is situated between 9o51'13.3" N latitude and 78o44'44.6' N longitude, with average rainfall and temperature ranging from 603–800 mm and 29°C–41°C. The study area occupies an area of 234 acres. The study area is predominantly covered by agricultural, horticultural and agroforestry ecosystems. For the purpose of the butterfly survey, the entire campus was mapped into four different habitats: botanical garden area, agricultural field area, horticultural field area and grassland area.

**Habitat 1 (Botanical Garden area):** The botanical garden area comprises a lot of flowering plant species, *i.e.*, Ixora, Hibiscus, Jasmine, Plumeria, Yellow Elder, Paper Flower, Pin Wheel Flower, Water Lily, Cosmos, Table Rose, Rose, and Nerium and is beautifully maintained.

**Habitat 2 (Agricultural field area):** The agricultural field areas mainly produce Paddy, Ragi, Maize, Pulses (Black gram and green gram), Groundnut, Gingelly, Sorghum and Daincha. In addition, the area includes organic farming areas, traditional paddy cultivation fields and medicinal plant gardens.

**Habitat 3 (Horticultural field area):** The horticultural field area supports a variety of crops, *i.e.*, Chilli, Brinjal, Bhendi, Tomato, Mango, Guava, Sapota, Citrus, Marigold, Banana, Curry leaf, Grape, Watermelon, Castor and Water apple.

**Habitat 4 (Grassland area):** The grassland area is occupied by mixed vegetation, mainly grasses, climbers, creepers and shrubs, *i.e.*, Country mallow, Indian jointvetch, Water leaf, Milk weed, Giradol, Tick weed, Butterfly pea, Three-leaved caper, Wild poinsettia, Puncture vine, Morning glory, Rosy milk weed vine, Red pea eggplant, Prosopis, Coat buttons, Acacia, Desert horse purslane, Bamboo and Lantana. Some common trees that are grown on the campuses are Neem, Pungam, Rain tree, Jamun tree, Foxtail palm, Traveler's palm, Royal palm, Cutch tree, Trumphet tree, Gulmohar, Copper pod tree, Golden Shower tree, Giger tree, Singapore Cherry, Acacia, Marudham and Nuna tree. Artificial canals, fish ponds and standing water are additional specialties in the study area.



Fig. 1. Map showing the study location

## 2.2 Butterfly Survey

The survey and documentation of the butterfly were conducted for a period of four months, from April 2023 to July 2023. The standard modified Pollard walk method was used for field surveys with necessary modifications [30,31]. The butterflies were observed within 2.5 meters on the left and right sides and five meters in front of the observer. Field observations were taken twice a day, from early morning (8.0 am–10.0 am) to evening (3.0 pm–5.0 pm) during good weather conditions (no heavy rain or strong winds) to ensure maximum detection of butterflies on all working days. Photography was taken during the field visit using Canon DSLR cameras (77 and 80 D, 135-250 mm). Species identification was done with the help of field guides and books, viz., 'South Indian Butterfly: Field Guide [32], Butterflies of India [33], Butterflies of Tamil Nadu' [19] and other available literature [34]. Identification was also done with the help of consultation with experts and the online website I-Found-Butterflies (Indian Foundation for Butterflies) (<http://www.ifoundbutterflies.org/>). No specimens were collected or killed during the study period. On the basis of occurrence, the butterflies' species were classified as abundant (80–100%), common (60–80%), occasional (40–60%), rare (20–40%) and very rare (below 20%) [35]. Recorded species were categorized under the IUCN Red List [36] and their status in the Wildlife Protection Act (WPA) 1972 of India [37].

## 2.3 Data Analysis

Shannon-Wiener was calculated for butterfly diversity [38]. Species evenness was calculated by Pielou's evenness [39] and species richness was calculated by Margalef's index [40,41]. Also, a dominance index or relative diversity was calculated to compare the species abundance. Relative diversity is also known as percentage occurrence. The overall mean monthly sighting was calculated by the Seasonal Index [42]. All the data were pooled and analyzed using Microsoft Office Excel 2007.

### Shannon-Wiener Index (H')

$$H' = -\sum(P_i)(\ln P_i)$$

Where,

P<sub>i</sub> = Proportion of each species  
ln = Natural logarithm

### Pielou's Evenness Index (E):

$$E = H' / \ln(S)$$

Where,

S = Number of species

### Margalef's index (R):

$$R = (S-1) / \ln N$$

Where,

S = Total number of species

N = Total number of individuals in the sample

ln = Natural logarithm

### Dominance Index (Ds):

$$D_s = n_i \times 100 / N$$

Where,

n<sub>i</sub> = number of butterflies in the family

N = Total number of butterflies in all the family

### Seasonal Index (SI):

$$SI = \text{Month-wise mean} / \text{Overall mean} \times 100$$

Where, month-wise mean is the mean number of butterflies sighted during the study period and overall mean is the mean of all month-wise means.

## 3. RESULTS AND DISCUSSION

### 3.1 Family Wise Composition of Butterflies

The study revealed that a total of 976 individuals of 76 species belonging to 51 genera and 5 families of butterflies (Nymphalidae, Lycaenidae, Pieridae, Hesperidae, and Papilionidae) were recorded in this study (Tables 1–5). Photographs of each species categorized into their respective families were represented (Plates 1–5). The highest number of species and highest abundance have been observed in Nymphalidae (22 species from 12 genera, 29.0%, N=321), followed by Lycaenidae (20 species from 17 genera, 26.5%, N=285), Pieridae (18 species from 9 genera, 24.0%, N = 185), Hesperidae (12 species from 11 genera, 16.0%, N = 125) and the lowest number of species and abundance in Papilionidae (four species from 2 genera, 5.5%,

N = 60), as illustrated (Table 6).The study shows the highest species-to-genus ratio (S/G ratio) for Pieridae and Papilionidae compared to the other three families. The species-to-genus ratio (S/G) that determines species distribution among genera was 1.8, 1.2, 2.0, 1.1, and 2.0 for Nymphalidae, Lycaenidae, Pieridae,

Hesperiidae, and Papilionidae, respectively (Table 6). Among the total recorded species, their relative abundance indicates that 33.0% are common (25species), followed by abundant (32.0%, 24 species), occasional (18.0%, 14 species), rare (13%, 10 species) and Very rare (3.0%,2 species) (Fig. 2; Tables 1-5).

**Table 1. Check list of butterfly species belonging to the family Nymphalidae recorded in the study area**

S. No.	Common Name	Scientific Name	IUCN Status	Abundance
1	Lemon Pansy	<i>Junonia lemonias</i> (Linnaeus, 1758)	NE	A
2	Blue Pansy	<i>Junonia orithiya</i> (Linnaeus, 1758)	NE	R
3	Peacock Pansy	<i>Junonia almana</i> (Linnaeus, 1758)	LC	O
4	Grey Pansy	<i>Junonia atlites</i> (Linnaeus, 1763)	NE	C
5	Yellow Pansy	<i>Junonia hierta</i> (Fabricius, 1798)	LC	O
6	Chocolate Pansy	<i>Junonia iphita</i> (Cramer, 1779)	NE	O
7	Plain Tiger (Male)	<i>Danaus chrysippus</i> (Linnaeus, 1758)	LC	A
8	Striped Tiger (Male)	<i>Danaus genutia</i> (Cramer, 1779)	NE	A
9	Angled Castor	<i>Ariadne ariadne</i> (Moore, 1884)	NE	C
10	Common Castor	<i>Ariadne merione</i> (Cramer, 1777)	NE	C
11	Danaid Egg Fly (Female)#	<i>Hypolimnas misippus</i> (Linnaeus, 1764)	LC	R
12	Great Egg Fly	<i>Hypolimnus bolina</i> (Linnaeus, 1758)	NE	R
13	Common Four Ring	<i>Ypthima huebneri</i> (Kirby, 1871)	NE	C
14	White Four Ring	<i>Ypthima ceylonica</i> (Hewitson, 1865)	LC	R
15	Common Bush Brown	<i>Mycalesis perseus</i> (Fabricius, 1775)	NE	C
16	Glade Eye Bush Brown	<i>Mycalesis patina</i> (Moore, 1857)	NE	C
17	Joker	<i>Byblia ilithyia</i> (Drury, 1773)	NE	C
18	Common Indian Crow****	<i>Euploea core</i> (Cramer, 1780)	LC	C
19	Tawny Costor	<i>Acraea violae</i> (Linnaeus, 1758)	NE	C
20	Common Evening Brown	<i>Melanitis leda</i> (Linnaeus, 1758)	NE	A
21	Blue Tiger	<i>Tirumala limniace</i> (Cramer, 1775)	NE	A
22	Common Sailor	<i>Neptis hylas</i> (Linnaeus, 1758)	NE	C

IUCN Status: NE-Not evaluated; LC-Least concern Abundance: A-Abundant, C-Common, O-Occasional, R-Rare Wildlife Protection Act- 1972: \*\*\*\*Schedule IV, #Schedule I&II

**Table 2. Checklist of butterfly species belonging to the family Lycaenidae recorded in the study area**

S. No.	Common Name	Scientific Name	IUCN Status	Abundance
1	Small Grass Jewel	<i>Freyeria putli</i> (Kollar, 1844)	NE	A
2	Grass Jewel	<i>Freyeria trochylus</i> (Freyer, 1845)	NE	A
3	Plain Cupid	<i>Chilades panadava</i> (Horsefield, 1829)	NE	C
4	Lime Blue	<i>Chilades lajus</i> (Stoll, 1780)	NE	A
5	Gram Blue**	<i>Euchrysops cnejus</i> (Fabricius, 1798)	NE	A
6	Common Smoky Blue	<i>Euchrysops malathana</i> (Boisduval, 1833)	NE	C
7	Dull Babul Blue	<i>Azanius uranus</i> (Butler, 1886)	NE	C
8	Forest Pierrot	<i>Taraka hamada</i> (Druce, 1875)	NE	C
9	Common Pierrot##	<i>Castalius rosimon</i> (Fabricius, 1775)	NE	C
10	Indian Cupid	<i>Everes lacturnus</i> (Godart, 1824)	NE	C

S. No.	Common Name	Scientific Name	IUCN Status	Abundance
11	Rounded Pierrot	<i>Tarucus nara</i> (Kollar, 1848)	NE	VR
12	Zebra Blue	<i>Leptotes plinius</i> (Fabricius, 1793)	NE	A
13	Common Silverline	<i>Spindasis vulcanus</i> (Fabricius, 1775)	NE	A
14	Common Cerulean	<i>Jamides celeno</i> (Cramer, 1775)	NE	A
15	Lesser Grass Blue	<i>Zizina otis</i> (Fabricius, 1787)	NE	A
16	Tiny Grass Blue	<i>Zizula hylax</i> (Fabricius, 1775)	NE	A
17	Tamil Large Guava Blue	<i>Virachola Isocrates</i> (Fabricius, 1793)	NE	O
18	Tawny Silver Line	<i>Cigaritis acamas</i> (Klug, 1834)	NE	C
19	Forget-Me-Not	<i>Catochrysops strabo</i> (Fabricius, 1793)	NE	A
20	Ceraunus Blue	<i>Hemiargus ceraunus</i> (Fabricius, 1793)	NE	O

IUCN Status: NE-Not evaluated

Abundance: A-Abundant, C-Common, O-Occasional, VR-Very Rare Wildlife Protection Act-1972: \*\*Schedule II (Part II), ##Schedule I&IV

**Table 3. Checklist of butterfly species belonging to the family Pieridae recorded in the study area**

S. No.	Common Name	Scientific Name	IUCN Status	Abundance
1	Small Grass Yellow	<i>Eurema brigitta</i> (Stoll, 1780)	LC	C
2	One Spot Grass Yellow	<i>Eurema andersoni</i> (Moore, 1886)	LC	A
3	Common Grass Yellow	<i>Eurema hecabe</i> (Linnaeus, 1758)	NE	A
4	Small Orange Tip (Female)	<i>Colotis etrida</i> (Boisduval, 1836)	NE	C
5	Crimson Tip (Male)	<i>Colotis danae</i> (Fabricius, 1775)	NE	R
6	Plain Orange Tip	<i>Colotis aurora</i> (Cramer, 1780)	NE	O
7	Desert Orange Tip	<i>Colotis evagore</i> (Klug, 1829)	NE	O
8	Large Salmon Arab	<i>Colotis fausta</i> (Oliver, 1804)	NE	R
9	Common Albatross**	<i>Appias albino</i> (Boisduval, 1836)	NE	C
10	Striped Albatross***	<i>Appias libythea</i> (Fabricius, 1775)	NE	R
11	Mottled Emigrant	<i>Catopsilia pyranthe</i> (Linnaeus, 1758)	NE	A
12	Oriental Lemon/Common Emigrant	<i>Catopsilia pomona</i> (Fabricius, 1775)	NE	C
13	African Migrant	<i>Catopsilia florella</i> (Fabricius, 1775)	NE	C
14	Pioneer	<i>Belenois aurota</i> (Fabricius, 1793)	LC	A
15	Common Gull**	<i>Cepora nerissa</i> (Fabricius, 1775)	NE	A
16	Psyche	<i>Leptosia nina</i> (Fabricius, 1793)	NE	O
17	Andaman Great Orange Tip	<i>Hebomoia roepstorffii</i> (Wood-Mason, 1880)	NE	O
18	White Orange Tip	<i>Ixias marianne</i> (Cramer, 1779)	NE	C

IUCN Status: NE-Not evaluated; LC-Least concern Abundance: A-Abundant, C-Common, O-Occasional, R-Rare, Wildlife Protection Act-1972: \*\*Schedule II (Part II), \*\*\*Schedule III

Butterflies are important bioindicators of the ecosystem. Potent pollinators and ecological indicators like butterflies are important for estimating the general health of an ecosystem. Butterflies play a pivotal role in ecosystem restoration as pollinators. An increase in butterfly populations may indicate an increase in plant diversity in the area [43]. The present study, for the first time ever, aims at evaluating butterflies on our campus. The results accumulated so far

clearly indicate that the overall diversity of butterflies on our campus is quite good. Nymphalidae represented the highest number of species in the present study (22.0 species from 12 genera, 29.0%), while the Hesperidae (12.0 species from 11 genera, 16.0%) and Papilionidae (4.0 species from 2 genera, 5.5%) families represented the lowest number of species. Similarly, a high number of species from Nymphalidae were observed by [44-48] in their

studies. The highest species richness and abundance might be due to the presence of host plants suitable for Nymphalidae [49], high dispersal ability, ecological adaptation, being polyfag characteristic and strong or active flight, enabling them to search for resources in large geographical areas [6,50-53]. This is consistent with Nymphalidae being the largest butterfly family, accounting for one-third of all known Species worldwide [14]. According to [54] this phenomenon could be due to the large number of species in the family as well as their wide variety of food selection, including flowers, fruits, honey dew, tree sap, rotten materials and decomposing carcasses. The reason for the abundance of Lycaenidae and Pieridae butterflies in the study area can be attributed to the dominance of their food plants belonging to the genus *Cassia*, *Albizia* and *Bauhinia* in the study region [13,55]. However, in the present study, Hesperidae and Papilionidae were poorly

represented in the study area, which reported members of Papilionidae to be the least represented. One of the possible reasons for this difference could be due to the difficulties in observing Hesperidae butterflies because of the limited monitoring time (8.0–10.0 am and 3.0–5.0 pm), their dull color and their crepuscular habit, i.e., that they are active in the early morning and to a lesser extent in the evening. This probably indicates that the climate, vegetation type and availability of host plants in the present study area may not be suitable for members of the Hesperidae family. The availability of host plants determines butterfly abundance [56]. Butterflies solely depend on plants and their diversity highly depends on plant species diversity [23,57]. This, however, needs to be confirmed by further extensive surveys in this region. It is not unlikely that a few butterfly species may have escaped notice and will be added in the future.

**Table 4. Checklist of butterfly species belonging to the family Hesperidae recorded in the study area**

S. No.	Common Name	Scientific Name	IUCN Status	Abundance
1	Common Grass Dart	<i>Taractrocer a maeuius</i> (Fabricius, 1793)	NE	O
2	Tamil Grass Dart	<i>Taractrocer a ceramas</i> (Hewitson, 1868)	NE	VR
3	African Marbled Skipper	<i>Gomalia elma</i> (Trimen, 1862)	NE	R
4	Chinese Swift	<i>Pelopodas sinensis</i> (Mabille, 1877)	NE	R
5	Rice Swift	<i>Borbo cinnara</i> (Wallace, 1866)	NE	A
6	Indian Palm Bob	<i>Suastus gremius</i> (Fabricius, 1798)	NE	C
7	Brown Awl	<i>Badamia exclamationis</i> (Fabricius, 1775)	LC	A
8	Grass Demon	<i>Udaspes folus</i> (Cramer, 1775)	NE	O
9	Grizzled Skipper	<i>Pyrgus malvae</i> (Linnaeus, 1758)	NE	O
10	Acacia Skipper	<i>Cogia hippalus</i> (Edwards, 1882)	NE	C
11	Manfreda Giant Skipper	<i>Stallingsia maculosus</i> (Freeman, 1955)	NE	O
12	Common Bush Hopper	<i>Ampittia dioscorides</i> (Fabricius, 1793)	NE	R

IUCN Status: NE-Not evaluated; LC-Least concern

Abundance: A-Abundant, C-Common, O-Occasional, R-Rare, VR-Very Rare

**Table 5. Checklist of butterfly species belonging to the family Papilionidae recorded in the study area**

S. No.	Common Name	Scientific Name	IUCN Status	Abundance
1	Common Lime	<i>Papilio demoleus</i> (Linnaeus, 1758)	NE	A
2	Common Mormon	<i>Papili opolytes</i> (Linnaeus, 1758)	NE	A
3	Common Rose	<i>Pachliopta aristolochiae</i> (Fabricius, 1775)	LC	C
4	Crimson Rose*	<i>Pachliopta hector</i> (Linnaeus, 1758)	LC	O

IUCN Status: NE-Not evaluated; LC-Least concern Abundance: A-Abundant, C-Common, O-Occasional Wildlife Protection Act-1972: \*Schedule I (Part IV)





Lemon Pansy




Blue Pansy




Peacock Pansy



Striped Tiger



Yellow Pansy



Chocolate Pansy



Plain Tiger



Great Egg Fly



Angled Castor



Common Castor



Danaid Egg Fly



Common Evening Brown





Common Four Ring



White Four Ring



Common Bush Brown



Blue Tiger



Joker



Common Indian Crow



Tawny Costor



Common Sailor

**Plate 1. Photographs of butterfly species belonging to the family Nymphalidae recorded in the study area**



Small Grass Jewel



Grass Jewel



Plain Cupid



Lime Blue



Gram Blue



Common Smoky blue



Dull Babul Blue



Forest Pierrot



Common Pierrot



Indian Cupid



Rounded Pierrot



Zebra Blue



Common Silverline



Common Cerulean



Lesser Grass Blue



Tiny Grass Blue



Tamil Large Guava Blue

**Plate 2. Photographs of butterfly species belonging to the family Lycaenidae recorded in the study area**





Small Grass Yellow



One Spot Grass Yellow



Common Grass Yellow



Small Orange Tip



Crimson Tip



Plain Orange Tip



White Orange Tip



Large Salmon Arab



Common Albatross



Common Gull



Mottled Emigrant



Oriental Lemon/Common Emigrant



Psyche



Pioneer

**Plate 3. Photographs of butterfly species belonging to the family Pieridae recorded in the study area**



Common Grass Dart



Tamil Grass Dart



African Marbled Skipper



Chinese Swift



Rice Swift



Indian Palm Bob



Brown Awl



Grass Demon



Common Bush Hopper

**Plate 4. Photographs of butterfly species belonging to the family Hesperidae recorded in the study area**



Common Lime



Common Mormon



Common Rose



Crimson Rose


**Plate 5. Photographs of butterfly species belonging to the family Papilionidae recorded in the study area**



**Table 6. Family wise composition, species to genus ratio and diversity indices of butterflies in the study area**

S.No.	Family	Number of Species (S)	Number of Genus (G)	S/G	N	H	E	R	D (%)
1	Nymphalidae	22	12	1.8	321	0.36	0.12	3.64	29.0
2	Lycaenidae	20	17	1.2	285	0.35	0.12	3.36	26.5
3	Pieridae	18	9	2.0	185	0.34	0.12	3.26	24.0
4	Hesperiidae	12	11	1.1	125	0.29	0.12	2.28	16.0
5	Papilionidae	4	2	2.0	60	0.15	0.11	0.73	5.5
6	Total	76	51	-	976	1.50	0.60	13.30	100

S/G-Species to Genus Ratio; N-Number of individuals, H-Shannon-Wiener Index, E- Pielou's Evenness, R- Margalef's index, D- Dominance Index



**Fig.2. Local status of recorded butterflies in the study area**

**3.2 Butterfly Community, Composition in Different Habitats**

The maximum number of species and abundance were observed in the garden area (140 species, N = 280), followed by the horticultural field area (129 species, N = 258), the agricultural field area (115 species, N = 230) and the minimum in the grassland area (104 species, N = 208) (Fig. 3; Table 7).

The highest number of species with the highest abundance in the botanical garden area

compared to agricultural, horticultural and grassland areas may be due to the high diversity of host plants, rich nectar resources and variety of flowers, viz., Ixora, Hibiscus, Jasmine, Plumeria, Yellow Elder, Paper Flower, Pin Wheel Flower, Water Lily, Cosmos, Table Rose, Rose, Nerium, Lantana and Golden Shower Tree [58] and undisturbed habitats compared to other areas. The artificially created site, like a botanical garden, mainly contains cultivated plants, especially nectar-rich flowering plants, which host several butterfly species and hence report high diversity compared to the natural areas.

**Table 7. Butterflies' species composition in different months and habitats in the study area**

S. No.	Months	Garden area		Agricultural Field Area		Horticultural Field Area		Grassland Area		Total		Seasonal Index (SI) %
		S	N	S	N	S	N	S	N	S	N	
1	April	31	62	27	54	33	66	21	42	112	224	23.0
2	May	14	28	21	42	18	36	16	32	69	138	14.0
3	June	42	84	31	62	36	72	32	64	141	282	29.0
4	July	53	106	36	72	42	84	35	70	307	332	34.0
5	Total	140	280	115	230	129	258	104	208	629	976	100

S=Number of species; N=Number of individuals

Agricultural field areas have low plant diversity. Lower butterfly diversity in agricultural lands might be due to agricultural intensification, chemical fertilizers and the usage of pesticides and herbicides [59,60,61]. The lowest number of butterfly species recorded in the grassland area might be due to continuous anthropogenic activities and the limited availability of plants. Another reason for the low abundance of butterfly species in grassland areas could be the absence of shade in these areas. Anthropogenic disturbance around grassland areas affects butterfly communities [62,63]. The agricultural expansion and intensification, overgrazing by cattle in grassland areas, construction of buildings leading to habitat destruction and movement of heavy vehicles may cause a shift in the relative abundance and diversity of butterflies in the study area. These disturbances are degrading habitat quality and adversely affecting the food resources of butterflies, which in turn have the potential to accelerate declines in their abundance and diversity. Hence, conservation of butterflies is necessary to maintain biodiversity and prevent extinction. The primary goal of conservation is to identify the different areas or hot spots that support the butterfly population and their survival [64].

### 3.3 Butterfly Species Composition in Different Months

The maximum number of butterfly species and abundance was recorded in July (307 species,  $N = 332$ ), followed by June (141 species,  $N = 282$ ), April (112 species,  $N = 224$ ) and the minimum in May (69 species,  $N = 138$ ) (Fig. 4; Table 7). Similarly, the Seasonal Index (SI) also recorded the maximum number of butterfly species in July (34.1%), followed by June (29.5%), April (24.0%), and the minimum in May (14.2%).

The highest number of butterflies was seen in the month of July, followed by June and the least in the dry seasons (May and April). The butterflies tend to avoid dry habitats and prefer moist places [65]. The butterfly population rapidly declined during the period from April to May. Usually, in southern India, these months are very hot and dry. Moreover, factors such as scarcity of water, poor nectar butterfly pasture and dry vegetation result in lower butterfly abundance and lower survival abilities for most species. Seasonal fluctuations of butterflies are often influenced by climatic and ecological conditions such as temperature, photoperiod, rainfall, humidity, variation in the availability of food resources,

vegetation cover such as herbs and shrubs and the presence of natural enemies [66,67,68,69]. Low rainfall, high temperatures, and high wind speeds were recorded from April to May and the disappearance of green vegetation was witnessed during this period. When the summer heat is high, they tend to go into hiding inside the bushes and temperature highly affects the activity of butterflies, their distribution, growth and breeding. High air temperatures cause a decrease in the volume of nectar secretion in flowers [70]. Accordingly, the abundance of butterflies was low during this period. The distribution and diversity of butterflies vary depending on the season. They are abundant in some months and absent during other months [71]. When the distribution of families over four months was compared, we noted an increase in the number of butterflies towards the monsoon period (July). [72] reported that rainfall conditions greatly influence butterfly numbers and species distributions. In southern India, the monsoons promote an increase in foliage cover and density and have a profound effect on the seasonal occurrence of butterflies. The young, tender leaves in ample quantity stimulated by high rainfall provide better quality food for larvae and thus result in peak abundance of butterflies during monsoons [73].

### 3.4 Diversity Indices, WPA, 1972 and IUCN Status of Recorded Butterflies in the Study Area

Diversity indices of recorded species indicated that Nymphalidae was the most diverse and species-rich family ( $H = 0.36$ ,  $E = 0.12$ ,  $R = 3.64$ ), while the family Papilionidae was the least diverse and least species-rich ( $H = 0.15$ ,  $E = 0.11$ ,  $R = 0.73$ ) (Table 6). The overall Shannon-Weiner diversity index was  $H = 1.50$ , Pielou's evenness index was  $E = 0.60$  and Margalef's index was  $R = 13.30$ . Out of the total 76 species, twelve were categorized under the least concern (LC) status as per the IUCN RedList (IUCN, 2022). The remaining sixty-four species were marked as not evaluated (NE). Eight species were legally protected in India under the Wildlife Protection Act of 1972 under Schedules I, II, III, and IV of the acts (Tables 1–5). A total of eight species of butterflies from the study area are designated under the Wildlife Protection Act (WPA, 1972) while describing their status and justifying their inclusion in the scheduled list, suggesting the need for strict conservation measures.




Fig. 3. Species number and individual number of butterflies in different habitat in the study area




Fig. 4. Month-wise distribution of recorded butterflies in the study area

#### 4. CONCLUSION

The present study demonstrated that 976 individuals of butterflies of 76 species belonging to 51 genera and five families were recorded. More butterfly species and abundance were recorded in the study area during the month of July, as well as in the habitat of the garden area. The study area is rich in butterflies. Although the study area supports a good number of butterfly species, it requires detailed further study throughout the year. The present list is not conclusive and exploration is needed to update this checklist. This information will help in future research on butterflies and conserve them by

establishing a butterfly park and garden and by creating awareness among school and college students.

#### ACKNOWLEDGEMENTS

The authors are highly thankful to the Chairman, Dr. Sethu Kumanan, and Principal, Dr. K. Karunaniithi, SBAC & RF, Karaikudi, for granting permission and providing the necessary facilities to carry out the survey. The authors are grateful to Ms. K.S. Anu Priya Lashmi and Mr. M. Dhineshbabu for providing cameras to document butterflies during the survey. The authors are also thankful to Dr. P.M.M. David, retired

professor, for his constant encouragement and to Mr. M. Makizhan Mathi for his meticulous support while preparing the manuscript.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Tiple AD, Khurad AM. Butterfly species diversity, habitats and seasonal distribution in and around Nagpur City, central India. *World Journal of Zoology*. 2009;4(3):153-162.
2. Borges RM, Gowda V, Zacharias M. Butterfly pollination and high contrast visual signals in a low density distylous plant. *Oecologia*. 2003;136:571-573.
3. Tiple AD, Khurad AM, Dennis RL. Butterfly diversity in relation to a human-impact gradient on an Indian university campus. *Nota Lepidopterologica*. 2007;30(1):179.
4. Merlin C, Liedvogel M. The genetics and epigenetics of animal migration and orientation: birds, butterflies and beyond. *Journal of Experimental Biology*; 2019; 222(1):191890.
5. Alarape AA, Omifolaji JK, Mwansat GS. Butterfly Species Diversity and Abundance in University of Ibadan Botanical Garden, Nigeria. *Open Journal of Ecology*. 2015;5: 352-360.
6. Majumder J, Lodh R, Agarwala BK. Variation in butterfly diversity and unique species richness along different habitats in Trishna Wild-life Sanctuary, Tripura, Northeast India. *Check-List*. 2012;8(3): 432-436.
7. Ghazanfar M, Malik MF, Hussain M, Iqbal R, Younas M. Butterflies and their contribution in ecosystem: A review. *Journal of Entomology and Zoology Studies*. 2016;4(2):115-118.
8. Klein AM, Vaissiere BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C, Tscharntke T. Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society. Biological Sciences*. 2007;274(1608):303-313.
9. Syaripuddin K, Sing K-W, Wilson J-J. Comparison of Butterflies, Bats and Beetles as Bioindicators Based on Four Key Criteria and DNA Barcodes. *Tropical Conservation Science*. 2015;8(1):138-149.
10. Lestari DF, Putri RDA, Ridwan M, Purwaningsih AD. Butterfly diversity (Insecta: Lepidoptera) in Alas Bromo Forest Tourism, North Lawu BKP, Karanganyar, Central Tengah SS (in Indonesian). *Prosiding Seminar Nasional Masyarakat Biodiversitas Indonesia*. 2015;1(6):1284-1288.
11. Sawchik J, Dufrêne M, Lebrun P. Distribution patterns and indicator species of butterfly assemblages of wet meadows in southern Belgium. *Belgium Journal of Zoology*. 2005;135(1):43-52.
12. Castro A, Espinosa CI. Seasonal diversity of butterflies and its relationship with woody-plant resources availability in an Ecuadorian tropical dry forest. *Tropical Conservation Science*. 2015;8(2):333-351.
13. Kunte K. The Wildlife (Protection) Act and conservation prioritization of butterflies of the Western Ghats, southwestern India. *Current Science*. 2008;94:729-735.
14. Tiple AD. Butterflies of Vidarbha region, Maharashtra State, central India. *Journal of Threatened Taxa*. 2011;3(1):1469-1477.
15. Sidhu AK. Lepidoptera of India: A Review. *Journal of Entomology and Zoology Studies*. 2023;11(2):105-114.
16. Karmakar T, Nitin R, Sarkar V, Baidya S, Mazumder S, Chandrasekharan V, Veino J. Early stages and larval host plants of some northeastern Indian butterflies. *Journal of Threatened Taxa*. 2018;10(6): 11780-11799.
17. Nitin R, Balakrishnan V, Churi PV, Kalesh S, Prakash S, Kunte K. Larval host plants of the butterflies of the Western Ghats, India. *Journal of Threatened Taxa*. 2018; 10(4):11495-11550.
18. Pavendhan A. Butterfly Check list for the state of TamilNadu. In: *Pattampoochi - Wings 1 Veins 1*. Coimbatore, Tamil Nadu, India: Tamil Nadu Butterfly Society. 2017; 1:4-13.
19. Mohan Prasath P, Satheesh N. Butterfly of Tamil Nadu, Tamil Nadu Forest Department, Tiruchirappalli. 2018;223.
20. Kehmiker I. The book of the Indian Butterflies. *Bombay Natural History Society*. Mumbai. 2008;497p.
21. Mennechez G, Schtickzelle N, Baguette M. Metapopulation dynamics of the bog fritillary between a continuous and a highly fragmented butterfly: Comparison of demographic parameters and dispersal landscape. *Landscape Ecology*. 2003; 18:279-291.

22. Hoyle M, James M. Global Warming, Human Population Pressure, and Viability of the World's Smallest Butterfly. *Conservation Biology*. 2005;19(4):1113–1124.
23. Padhye AD, Dahanukar N, Paingankar M, Deshpande M, Deshpande D. Season and landscape wise distribution of butterflies in Tamhini, northern Western Ghats, India. *Zoos' Print Journal*. 2006;21(3):2175-2181.
24. Bates AJ, Sadler JP, Grundy D, Lowe N, Davis G, Baker D, Bridge M, Freestone R, Gardner D, Gibson C, Hemming R, Howarth S, Orridge S, Shaw M, Tams T, Young H. Garden and landscape-scale correlates of moths of differing conservation status: significant effects of urbanization and habitat diversity. *PlosOne*. 2014;9(1):e86925.
25. Dirzo R, Young HS, Galetti M, Ceballos G, Isaac NJB, Collen B. Defaunation in the Anthropocene. *Science*. 2014;345:401-406.
26. Sanchez-Bayo F, Wyckhuys KAG. Worldwide decline of the entomo fauna: A review of its drivers. *Biological Conservation*. 2019;232:8-27.
27. Meeta S, Noopur S. Suitability of Butterflies as Indicators of Ecosystem Condition: A Comparison of Butterfly Diversity across four habitats in Gir Wildlife Sanctuary. *International Journal of Advanced Research in Biological Sciences*. 2017;4(3):43-53.
28. Panicker, Varuna P, Anuraj R, Govind Shankar, Aswathy S, Anu Gopalakrishnan, and S. Biju. Studies on the diversity of butterfly (Lepidoptera: Rhopalocera) fauna in college of veterinary and animal sciences campus, Mannuthy, Thrissur, Kerala, India. *Journal of Entomology and Zoology Studies*. 2016;4(5):933-936.
29. Sharma K, Acharya BK, Sharma G, Valente D, Pasimeni M, Petrosillo I, Selvan T. Land use effect on butterfly alpha and beta diversity in the Eastern Himalaya, India. *Ecological Indicators*. 2020;110: 105605.
30. Pollard E. A method for assessing changes in the abundance of butterflies, *Biological Conservation*. 1977;12:115-134.
31. Pellet J, Bried JT, Parietti D, Gander A, Heer PO, Cherix D, Arlettaz R. Monitoring butterfly abundance: beyond Pollard walks. *PLoS One*. 2012;7(7):e41396-e41396.
32. Gunathilagaraj K, Perumal TNA, Jayaram K, Ganeshkumar, M. South Indian Butterfly: Field Guide. Krab media and Marketing. 2015;359.
33. Kehimkar I. Butterflies of Indian butterflies. Bombay Natural History Society, Mumbai. 2016;516pp.
34. Antram CB. A Descriptive Handbook of Indian Butterflies. Mittal publications, New Delhi. 2016;226 pp.
35. Tiple AD. Butterfly species diversity, relative abundance and status in Tropical Forest Research Institute, Jabalpur, Madhya Pradesh, central India, *Journal of Threatened Taxa*. 2012;4(7):2713-2717.
36. IUCN. The IUCN Red List of Threatened Species; 2022 April 25. Available: <https://www.iucnredlist.org>.
37. Anonymous. The Wildlife (Protection) Act, 1972. Natraj Publishers, Dehradun. 2006; 328.
38. Shannon CE, Weaver W. The mathematical theory of communication. Univ. Illinois Press, Urbana; 1949.
39. Pielou EC. An introduction to mathematical ecology. John Wiley: New York, USA. 1969;286.
40. Magurran AE. Measuring biological diversity. Blackwell Publishing Company: Hoboken, New Jersey. 2004;256.
41. Henderson PA. Practical Methods in Ecology, Blackwell Publishers, United Kingdom. 2005;1–148.
42. Mathew G, Anto M. In-situ conservation of butterflies through establishment of Butterfly Gardens: a case study at Peechi, Kerala, India. *Current Science*. 2007;93(3): 337-347.
43. Priya, Krishnaraj V, Janaranjini S, Lakeshmanaswamy. Studies on butterfly Diversity in adichanalloor Village, Kollam District, Kerala. *Journal of Entomology and Zoology Studies*. 2017;5(5):73-81.
44. Kumar P, Makesh Kumar B. Seasonal Diversity and Distribution of Butterflies in G.Venkataswamy Naidu College Campus, Kovilpatti, Tuticorin District, Tamil Nadu. *Advances in Zoology and Botany*. 2021; 9(5):91-99.
45. Miya MS, Gautam D, Neupane B, Chhetri A. Species diversity and abundance of Odonata in Sishaghat of Tanahun district, Nepal. *Journal of Animal Diversity*. 2021;3(3):45- 55.
46. Samal SK, Satapathy A, Pattanaik N. Diversity of butterflies (Lepidoptera: Rhopalocera) in Bhubaneswar, Odisha, India. *Notulae Scientia Biologicae*. 2021; 13(4):11074.

47. Nagarajan VM, Theivaprakasham H. Butterfly diversity of Chennai, Kancheepuram, Chengalpet and Thiruvallur districts in Tamil Nadu, India. *Journal of Entomology and Zoology Studies*. 2020;8(6):743-751.
48. Vinithashri G, Kennedy JS. Butterfly diversity in relation to host and nectar food plants in TNAU Botanical Garden, Coimbatore. *Journal of Environmental Biolpgy*. 2021;42:1141- 1151.
49. MalabikaSK. Impact of tropical forest degradation on nymphalid butterflies: A case study in Chandubi tropical forest, Assam, India. *International Journal of Biodiversity and Conservation*. 2011; 3(12):650-669.
50. Eswaran R, Pramod P. Structure of butterfly community of Anaikatty hills, Western Ghats. *Zoos' Print Journal*. 2005; 20(8):1939-1942.
51. Krishnakumar N, Kumaraguru A, Thiyagesan K, Asokan S. Diversity of papilionid butterflies in the Indira Gandhi wildlife sanctuary, Western Ghats, Southern India. *Tiger paper*. 2008;35(1):1-8.
52. Raut NB, Pendharkar A. Butterfly (Rhopalocera) fauna of Maharashtra Nature Park, Mumbai, Maharashtra, India. *Check List*. 2009;6(1):022-025.
53. Subedi B, Stewart A, Neupane B, Ghimire S, Adhikari H. Butterfly species diversity and their floral preferences in the Rupa wetland of Nepal. *Ecology and Evolution*. 2020;11:2086- 2099.
54. Rosmidi FH, Zahidin MA, Adanan A, Azizah A, Pesiu E, Abdullah MT. Checklist of butterflies in Pulau Perhentian and Pulau Bidong, Terengganu. *Journal of Sustainability Science and Management*. 2017;12(1):40-48.
55. Balasubramanian P, Mahendramani P, Padmapriya K. Comparison of plant biodiversity pattern of variously disturbed habitat of Moongilpallam area in the western ghats. *Salim Ali Centre for ornithology and Natural History, Coimbatore*. 2001;56-70.
56. Thomas JA. The ecology and conservation of Maculineaarion and other European species of large blue butterfly. *Ecology and Conservation of Butterflies*. 1995;pp180-197.
57. Ekroos J, Rundlöf M, Smith HG. Trait-dependent responses of flower-visiting insects to distance to semi-natural grasslands and landscape heterogeneity. *Landscape Ecology*. 2013;28(7):1283-1292.
58. Majumder J, Lodh R, Agarwala BK. Butterfly species richness and diversity in the Trishna Wildlife Sanctuary in South Asia. *Journal of Insect Science*. 2013; 13:79.
59. Geiger F, Bengtsson J, Berendse F, Weisse WW, Emmerson M, Morales MB, Ceryngier P, Liira J, Tscharntke T, Winqvist C, Eggers S, Bommarco R, Part T, Bretagnolle V, Plantegenest M, Clement LW, Dennis C, Palmer C, Onate JJ, Guerrerol, Hawro V, Aavik T, Thies C, Flohre A, Hanke S, Fischer C, Goedhart PW, Inchausti P. Persistent negative effects of pesticides on biodiversity and biological control potential on European farmland. *Basic and Applied Ecology*. 2010;11(2):97-105.
60. Batary P, Baldi A, Kleijn D, Tscharntke T. Landscape-moderated biodiversity effects of agri-environmental management: a meta-analysis. *Proceedings of the Royal Society B: Biological Sciences*. 2011; 278(1713):1894-1902.
61. Henry M, Beguin M, Requier F, Rollin O, Odoux JF, Aupine IP, Apte IJ, Tschamchian S, Decourtye A. Common pesticide decreases foraging success and survival in honey bees. *Science*. 2012; 336(6079):348-350.
62. Cabette RSH, Souza RJ, Shimano Y, Juen L. Effects of changes in the riparian forest on the butterfly community (Insecta: Lepidoptera) in Cerrado areas. *Revista Brasileira de Entomologia*. 2017;61:43-50.
63. An J, Choi S. Butterflies as an indicator group of riparian ecosystem assessment. *Journal of Asia-Pacific Entomology*. 2021; 24:195-200.
64. Myers N, Russell A, Mittermelert C, Mittermelert G, Gustavo AB, Fonseca KJ. Biodiversity hotspots for conservation priorities. *Nature*. 2000;403:853-858.
65. Rajagopal T, Sekar M, Archunan G. Diversity and community structure of butterfly of Arignar Anna Zoological Park, Chennai, TamilNadu. *Journal of Environmental Biology*. 2011; 32:201-207.
66. Grotan V, Lande R, Engen S, Sæther BE, DeVries PJ. Seasonal cycles of species diversity and similarity in a tropical butterfly community. *Journal of Animal Ecology*. 2012;81:714-723.



67. Valtonen A, Molleman F, Chapman CA, Carey JR, Ayres MP, Roininen H. Tropical phenology: biannual rhythms and interannual variation in an Afrotropical butterfly assemblage. *Ecosphere*. 2013; 4(3):36.
68. Sharmila EJ, Thatheyus AJ, Susaritha S, Snegapriya M. Seasonality of butterflies in Alagar Hills reserve forest, India. *Entomon*. 2020;45(1):53-60.
69. Tiple AD, Khurad AM. Butterfly species diversity, habitats and seasonal distribution in and around Nagpur City, central India. *World Journal of Zoology*. 2009;4(3):153-162.
70. Efendi MA. Diversity of butterflies (Lepidoptera: Ditrysia) in Corridor Forest, Gunung Halimun-Salak National Park, West Java. Thesis, School of Post Graduate Studies of Bogor Agricultural University, Bogor. 2009;500.
71. Kunte K. India, a Lifescape Butterflies of Peninsular India, Indian Academy of Sciences, University Press. 2000;243.
72. Reddi CS, Atluri JB, Venkata Ramana SP, Meera Bai G. The butterfly fauna of Vishakapatnam in South India. *Tiger Paper*. 2003;30:29-32.
73. Sengupta P, Banerjee KK, Ghorai, N. Seasonal diversity of butterflies and their larval food plants in the surroundings of upper Neora Valley National Park, a sub-tropical broad leaved hill forest in the eastern Himalayan landscape, West Bengal, India. *Journal of Threatened Taxa*. 2014;6(1):5327-5342.

© 2023 Narmadha et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/106145>