



Studies on Growth Characteristics and Quality Traits of Different Variety/ Genotypes of Brinjal (*Solanum melongena* L.)

Ram Ashish ^{a*}, Seema Sonkar ^b, C. L. Maurya ^c,
Rakesh Babu Gautam ^a, Nand Kumar ^a and Raj Kumar ^a

^a Department of Agricultural Biochemistry, CSAUA&T, Kanpur, India.

^b Department of Food Science and Nutrition, CSAUA&T, Kanpur, India.

^c Department of Seed Science and Technology, CSAUA&T, Kanpur, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i213947

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107892>

Original Research Article

Received: 02/08/2023

Accepted: 07/10/2023

Published: 13/10/2023

ABSTRACT

The present study was conducted to evaluate 20 variety/ genotypes of brinjal for quality parameters. The sample was collected from vegetable trial field of Chandra Shekhar Azad University of Agriculture and Technology, Kanpur as well as after grading of the sample and the quality analysis in the laboratory was conducted as per standard procedures in the laboratory of the department of agricultural biochemistry. A significant variation was detected in all traits studies. There was considerable variability among varieties. Among the growth characteristics data on the days to 50% flowering significantly varied from 31.78-40.42, plant height range from 64.05-82.65 cm, no. of branches per plant ranges from 9.0-14.5. Similarly among the quality traits such as ash content ranged from 9.71-11.39%, fat content varied from 0.23- 0.37%, fibre content ranged from 1.29-1.73%.

*Corresponding author: E-mail: ramashish271@gmail.com;

Keywords: Days to 50% flowering; plant height; No. of branches per plant; ash; fat; fibre.

1. INTRODUCTION

Eggplant or brinjal (*Solanum melongena* L.; $2n = 2x = 24$) is important vegetable crop and is considered a rich member of the family *Solanaceae*, which contains approximately 1300 species it can be grown in diversified climatic conditions of various ecological regions. It possesses high species richness with considerable flexibility of phenotypic adaptability that made the species the most important vegetable economically [1-3]. Eggplant is a general term for various *Solanum* species cultivated for their fruits, including the East Asian aubergine (*S. melongena* L.) and the two African native eggplants, Scarlet (*S. aethiopicum* L.) and Gboma (*S. macrocarpon* L.). Brinjal or eggplant (*Solanum melongena* L.) is one of the important solanaceous vegetable crop having diploid chromosome number $2n=2x=24$ [4-6,7,8]. One hundred gram edible portion of brinjal fruit contains 92.7% moisture, 24.0% calories, 4.0% carbohydrates, 1.4 g protein, 0.3 g fats, 1.3 g fibres, 124.0 (I.U.) vitamin A and 12.0 mg vitamin C [9]. It also contains 52.0 mg chlorine, 47.0 mg phosphorus, 44.0 mg sulphur and other minerals [10]. Plant height (cm), days to 50% flowering, primary branches per plant, fruit weight (g), fruit circumference (cm), fruits per plant, fruit length (cm,) were estimated according to Searle [11]. Brinjal has high nutritive value when compared with tomato [12-15]. It contains high amount of carbohydrates (6.4%), protein (1.3%), fat (0.3%), calcium (0.02%), phosphorus (0.02%), iron (0.0013 %) and other mineral matters [16].

2. MATERIALS AND METHODS

Plant height (cm): The height of the five randomly selected individual plants from ground to tip were measured in centimetre at final harvest stage, when the plant attained maximum growth and replication wise the average plant height of each genotype was calculated.

Days to 50% flowering: The number of days taken from the date of transplanting to the opening of first flower on 50% of the plants of each genotype in each replication was recorded as date and calculated in days.

Number of fruits per plant: Number of edible fruits from five tagged plants were counted at each picking and summed up for all the pickings and divided by five to get the number of fruits per plants.

Ash content: Ash content was determined by the method as described by Hart and Fisher, [17].

Fat content: Fat extractable lipid content in brinjal sample was determined by Soxhlet Extraction procedure using petroleum ether of boiling point 40-60 0C for six hours in a flash [18]. Fat content was calculated using the following formula:

$$\begin{aligned} & \text{Ether extract percentage in brinjal sample} \\ & = \frac{\text{weight of extracty (g)}}{\text{weight of sample}} \times 100 \end{aligned}$$

Fibre content: The fibre content in brinjal was analysed by the method as described by Hart and Fisher [17].

Statistical Analysis: All sample extract were prepared an analysis done using a Randomized Block Design (RBD) at 5% level of critical difference. Analysis of variance (ANOVA) for the design was carried out to determine the significance of differences among difference treatments.

3. RESULTS AND DISCUSSION

The data pertaining to days to 50% flowering of genotype cultivars for brinjal genotypes are presented in the Table 1. It was observed that the days to 50% flowering. The genotype C-9013 showed the maximum days to 50% flowering (40.42) followed by genotypes KS-453 (39.31), KS-556 (38.86) and C-8805 (38.37). The minimum days to 50% flowering was noted in genotype-KS- 331(31.78). The days to 50% flowering C-9013 were significantly superior than KS-331. The results are supported by Madhavi et al., [19]. Data regarding on Plant height of various cultivars of some brinjal genotypes are presented in the Table 1. It was observed that the Plant height. The genotype C-9012 showed the maximum plant height (82.65 cm) followed by genotype KS-554 (79.95 cm), KS-224 (78.10 cm) and C-9015 (77.75 cm). The minimum plant height was noted in genotype KS-453 (64.05 cm). The plant height C-9011were significantly superior than KS-453. These results are in close agreement with the reports Tripathy et al., [20]. Data regarding on No. of branches per plant of various cultivars of some brinjal genotypes are presented in the Table 2. It was observed that the no. of branches per plant. The genotype KS-

556 showed the maximum no. of branches per plant (14.99) followed by genotype C-7864-1 (14.5), C-9006 (14.16) and C- 9011 (13.83). The minimum No. of branches per plant was noted in genotype C- 8502-1 (07.50). The No. of branches per plant KS-556 were significantly superior than C-8502-1. These results are in close agreement with the reports Rahman et al., [21].

Data on ash content of various cultivars of some genotypes of brinjal are presented in the Table 2. It is shown that the data on the ash content varied from 9.71-11.39%. The genotype C-8805 showed the maximum ash content (11.395%) followed by genotype- Azad Kranti (11.33%), C-8841(11.28%) and C-9012 (11.17%). The minimum ash content was noted in genotype KS-235 (9.715%). The genotype of C-8805 superior than genotype of KS-235. The results are supported by Khan et al., [22]. Data on fat content of various cultivars of some genotypes of

brinjal are presented in the Table 2. It is shown that the data on the fat content varied significantly from 0.23-0.37%. The genotype of Azad Kranti showed the maximum fat content (0.37%) followed by genotype- C-9006 (0.36%), KS-555 (0.35%) and C-9012 (0.34%). The minimum fat content was noted in genotype of C-5623 (0.24). The genotype of Azad Kranti superior than the genotype of C-5623. The results are supported by Tetteh et al., [23]. Data on fibre content of various cultivars of some genotypes of brinjal are presented in the Table 2. It is shown that the data on the fibre content varied significantly from 1.29-1.73%. The genotype of KS-456 showed the maximum fibre content (1.73%) followed by genotype- C-9013 (1.63%), C-8841 (1.58%), and Azad Kranti (1.57%). The minimum fibre content was noted in genotype of C-9012(1.29%). The genotype of KS-456 superior than the genotype of C-9012. Similar results were reported by Rahman et al., [24].

Table 1. The data pertaining to days to 50 % flowering of genotype cultivars for brinjal

S.N.	Varieties	Days to 50 % flowering		Pooled mean	Plant height (cm)		Pooled mean	No. of branches per plant		Pooled mean
		2021	2022		2021	2022		2021	2022	
1	KS-235	34.32	32.23	33.275	65.4	67.6	66.5	08	10	9
2	KS-454	36.54	38.97	37.755	67.5	65.0	66.25	09	11	10
3	KS-456	31.22	35.43	33.325	70.7	67.9	69.3	11	14	12.5
4	KS-453	39.87	38.76	39.315	62.3	65.8	64.05	13	11	12
5	KS-224	36.54	38.78	37.66	78.9	77.3	78.1	09	12	10.5
6	KS-554	34.56	35.67	35.115	80.5	79.4	79.95	10	12	11
7	KS-555	37.65	36.76	37.205	78.2	76.5	77.35	13	11	12
8	KS-556	38.97	38.75	38.86	72.0	70.9	71.45	15	13	14
9	KS-331	31.23	32.34	31.785	68.2	67.6	67.9	08	10	9
10	C-9011	33.76	34.55	34.155	67.9	69.8	68.85	13	11	12
11	C-9012	33.08	35.46	34.27	83.4	81.9	82.65	10	12	11
12	C-9013	40.98	39.87	40.425	76.0	74.3	75.15	09	11	10
13	C-9015	37.65	36.72	37.185	78.9	76.6	77.75	08	10	9
14	C-9006	33.21	34.27	33.74	73.8	71.2	72.5	13	12	12.5
15	C-8502-1	36.75	34.56	35.655	67.5	65.7	66.6	15	14	14.5
16	C-8841	38.89	36.78	37.835	75.9	74.2	75.05	09	10	9.5
17	C-5623	35.65	37.62	36.635	69.8	70.9	70.35	10	11	10.5
18	C-7864-1	32.16	33.45	32.805	75.6	73.5	74.55	15	14	14.5
19	C-8805	39.09	37.65	38.37	72.3	70.9	71.6	11	13	12
20	Azad Kranti	37.65	38.97	38.31	77.7	76.5	77.1	10	12	11
	Mean	35.9885	36.3795	36.184	73.125	72.175	72.65	10.95	11.7	11.325
	S.E. m±	4.68	4.56	4.72	9.21	9.3	9.23	1.25	1.44	1.39
	C.D. (5%)	13.41	13.07	13.52	26.4	26.65	26.45	3.58	4.13	3.99

Table 2. Data on quality traits of various cultivars of some genotypes of brinjal

S.N.	Varieties	Ash Content (%)		Pooled mean	Fat (%)		Pooled mean	Feber (%)		Pooled mean
		2021	2022		2021	2022		2021	2022	
1	KS-235	09.56	09.87	9.715	0.26	0.35	0.305	1.5	1.3	1.4
2	KS-454	10.32	10.43	10.375	0.32	0.28	0.30	1.4	1.4	1.4
3	KS-456	10.87	10.76	10.815	0.38	0.32	0.35	1.8	1.6	1.7
4	KS-453	11.23	11.09	11.161	0.27	0.31	0.29	1.6	1.4	1.5
5	KS-224	10.87	10.65	10.76	0.29	0.35	0.32	1.3	1.3	1.3
6	KS-554	09.98	10.21	10.095	0.34	0.31	0.325	1.7	1.2	1.45
7	KS-555	11.37	10.95	11.16	0.31	0.37	0.331	1.2	1.6	1.4
8	KS-556	10.76	09.83	10.295	0.28	0.29	0.285	1.8	1.3	1.55
9	KS-331	10.54	10.93	10.735	0.29	0.26	0.275	1.3	1.4	1.35
10	C-9011	09.90	09.87	9.885	0.25	0.29	0.27	1.4	1.5	1.45
11	C-9012	11.21	11.13	11.17	0.36	0.31	0.335	1.2	1.3	1.25
12	C-9013	10.65	09.85	10.25	0.31	0.25	0.28	1.6	1.6	1.6
13	C-9015	10.87	10.21	10.54	0.29	0.31	0.30	1.3	1.4	1.35
14	C-9006	10.29	10.76	10.525	0.35	0.31	0.33	1.7	1.3	1.5
15	C-8502-1	09.87	09.56	9.716	0.26	0.32	0.29	1.4	1.6	1.5
16	C-8841	11.65	10.91	11.28	0.21	0.34	0.27	1.8	1.3	1.55
17	C-5623	10.90	09.85	10.375	0.27	0.24	0.25	1.2	1.4	1.3
18	C-7864-1	09.88	09.94	9.91	0.32	0.27	0.29	1.5	1.5	1.5
19	C-8805	11.56	11.23	11.395	0.34	0.28	0.31	1.7	1.4	1.55
20	Azad Kranti	11.68	10.98	11.33	0.37	0.31	0.34	1.4	1.7	1.55
	Mean	10.698	10.451	10.57425	0.3035	0.3035	0.3035	1.49	1.425	1.4575
	S.E. m±	1.35	1.35	1.36	0.04	0.04	0.07	0.2	0.18	0.37
	C.D. (5%)	3.86	3.87	3.88	0.11	0.12	0.21	0.57	0.52	1.11

4. CONCLUSION

The genotype C-9013, KS-453, KS-556 and C-8805 showed the maximum Days to 50% flowering, (40.42), (39.31), (38.86) and (38.37) Days to 50% flowering respectively. The minimum Days to 50% flowering was noted in genotype KS-331(31.78). The genotype C-9012, KS-544, KS-224 and C-9015 showed the maximum plant height (82.65 cm), (79.95 cm), (78.10 cm) and (77.75 cm) respectively. The minimum plant height was noted in genotype KS-453 (64.05 cm). The genotype KS-556, C-7864-1, C9006 and C-9011 showed the maximum no. of branches per plant (14.99), (14.50), (14.16) and (13.83) respectively. The minimum no. of branches per plant was noted in genotype C-8502-1 (07.50). The ash content ranged from in brinjal genotype C-8805, Azad Kranti, C-8841 and C-9012, 11.95%, 11.33%, 11.28% and 11.17%. While minimum ash content was noted in genotype KS-235 (9.715%). The genotype of Azad Kranti showed the maximum fat content (0.37%) by genotype- C-9006 (0.36 %), KS-555 (0.35%) and C-9012 (0.34%). The minimum fat content was noted in genotype of C-5623 (0.23%). Highest fibre content in various genotype of brinjal KS-456, C-9013, C-8841, and Azad Kranti (1.73%), (1.63%), (1.58%) and (1.57%). The minimum fibre content was noted in genotype of C-9012 (1.29%).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. AOAC. Official methods of analysis of the association of official Analytical Chemists, Washington D.C.
2. AOAC. Official methods of analysis: Association of Official Analytical Chemists, Washington, DC, USA; 1973.
3. Akpan N, Ogbonna P, Onyia V, Okechukwu E, Atugwu A. Dominic IO. Studies on the variability and combining ability for improved growth and yield of local eggplant genotypes (*Solanum melongena* L.) Not Sci. Biol. 2016;8(2):226-231.
4. Aykroyd WR, Gopalan C, Balasubramanian SC. The nutritive value of Indian foods and the planning of satisfactory diets (No. 23). New Delhi: Indian Council of Medical Research; 1963.
5. Chen NC, Li HM. Cultivation and seed production of eggplant. In Training workshop on vegetable cultivation and seed production technology (No. AVRDC Staff Publication). AVRDC; 1997.
6. Chinthagunti H, Sarnaik DA, Sharma D. Evaluation of brinjal (*Solanum melongena* L.) genotypes for flowering and yield parameters. Int. J. Curr. Microbiol. App. Sci. 2018;7(12):3101-3105.
7. Tirkey M, Saravana S, Lata P. Studies on variability, heritability and genetic advance for yield and its attributes in brinjal (*Solanum melongena* L.). Journal of Pharmacognosy and Phytochemistry, SP. 2018;(1):1181-1183.
8. Yadav S, Singh VB, Maurya R, Thapliyal V. Correlation and path coefficient analysis in brinjal (*Solanum melongena* L.). Int. J. Curr. Microbiol. and Appl. Sci. 2018;7: 2319-7706.
9. Chen LM, Hobbie S, Galan JE. Requirement of CDC42 for Salmonella-induced cytoskeletal and nuclear responses. Science, 274(5295). 1996;2115-2118.
10. Aykroyd UR. Indian Council of Medical Research, Special Report. Vegetable, National Book Trust India, New Delhi. 1963;42:188-191.
11. Searle SR. Phenotypic, genotypic and environmental correlations. Biometrics. 1961;17:474-780.
12. Dharmendra patidar1 MS, Shitap, Patel NA. Heterosis studies for fruit yield and its component in long type brinjal (*Solanum melongena* L.) Electronic Journal of Plant Breeding. 2017;8(4):1169-1176 (Dec 2017) ISSN 0975-928X
13. Ravali B, Reddy RK, Saidaiah P, Shivraj N. Genetic diversity in brinjal (*Solanum melongena* L.) International Journal of Current Microbiology and Applied Sciences. 2017;6(6):42-47.
14. Rodriguez-Jimenez JR, Amaya-Guerra CA, Baez-Gonzalez JG. Aguilera-Gonzalez C, Urias-Orona V, Nino-Medina G. Physicochemical, functional, and nutraceutical properties of eggplant flours obtained by different drying methods. Molecules. 2018;23(12): 3210.
15. Sharma TK, Pant SC, Kumar S, Paliwal A, Bahuguna P, Badhani HC. Combining ability studies in brinjal (*Solanum melongena* L.). Int. J. Bio-Resour. 2016;7 (6):1225-1231.
16. Kandoliya UK, Bajaniya VK, Bhadja NK, Bodar NP, Golakiya BA. Antioxidant and

- nutritional components of eggplant (*Solanum melongena* L.) fruit grown in Saurashtra region. Int. J. Curr. Microbiol. Appl. Sci. 2015;4(2):806-813.
17. Hart F, Fisher HJ. Modern food analysis. Springer-verlag; 1971.
 18. AOAC. Official methods of analysis of association of official agricultural chemist. 10th Ed. 1965;744.
 19. Madhavi N, Mishra AC, Prasad OJ, Bahuguna N. Studies on variability, heritability and genetic advance in brinjal (*Solanum melongena* L.). Plant Archives. 2017;15(1):277-281.
 20. Tripathy B, Sharma D, Jangde BP, Bairwa PL. Genetic variability and heritability studies in brinjal (*Solanum melongena* L.). The Bioscane, Special Issue. 2017; (10):109-116.
 21. Rahman MZ, Kabir H, Khan M. A study on brinjal production in Jamalpur district through profitability analysis and factors affecting the production. Journal of the Bangladesh Agricultural University. 2016; 14(1):113-118.
 22. Khan, Imtiaz Ali, H, Komal, A, Rasheed, K, Ashraf S, Muhammad, F, Abid, Ali Ijaz, A, Mukhtar. Proximate chemical composition of brinjal, *Solanum melongena* L. (*Solanales: Solanaceae*), genotypes and its correlation with the natural enemies in Peshawar Journal of Entomology and Zoology Studies 2015; 3(5): 07-11 in brinjal germplasm Journal of Crop and Weed. 2015;14(2): 51-60 (2018) ISSN: 2161-1009 Volume 5 • Issue 3 • 1000292 ISSN: 2320-7035
 23. Tetteh R, Aboagye LM, Boateng SK, Darko R, Obirih-Opareh J, Ibrahim AA. Variation in physiological seed quality of eggplant cultivars in relation to seed extraction time. Vegetos. 2022;1-7.
 24. Rahman MS, Rahman MH, Chowdhary MFN, Sultana MS, Ahmed KU. Effect of Spent Mushroom Substrate and Cowdung on Growth, Yield and Proximate Composition of Brinjal International Journal of Scientific and Research Publications, Volume 6, Issue 10, October 2016 468 ISSN 2250-3153; 2016.

© 2023 Ashish et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107892>