



# Effect of Application of Zn and Mn on Growth, Yield and Economics of Garden Pea (*Pisum sativum* var. *hortense* L.)

Heena <sup>a\*</sup>, Deependra Yadav <sup>a</sup>, Sandeep Kumar <sup>a</sup>,  
Ravinder <sup>b</sup> and Alok Kumar <sup>c</sup>

<sup>a</sup> Department of Horticulture, School of Agriculture, Abhilashi University, Mandi H.P. 175028, India.

<sup>b</sup> Department of Soil Science, School of Agriculture, Abhilashi University, Mandi, H.P. 175028, India.

<sup>c</sup> Department of Genetics and Plant Breeding, School of Agriculture, Abhilashi University, Mandi, H.P. 175028, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i193632

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105198>

Original Research Article

Received: 23/06/2023

Accepted: 29/08/2023

Published: 29/08/2023

## ABSTRACT

The present investigation entitled "Effect of application of Zn and Mn on growth and yield and economics of garden pea (*Pisum sativum* var. *hortense* L.) was carried out at the Research farm of the School of Agriculture, Abhilashi University, Mandi (H.P.) during Rabi season 2021. The experiment was laid out in a randomized block design with seven treatments in three replications. The result revealed that the soil chemical properties *i.e.*, Soil pH (5.98 pH) was maximum recorded in T<sub>6</sub> (basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5%) which showed non-significant effect on soil pH, available Zn and Mn was maximum recorded in T<sub>3</sub> basal application of Zn @ 100% of RDF + Mn @ 100% of RDF (1.67 ppm)

\*Corresponding author: E-mail: [hinathakurrajput44@gmail.com](mailto:hinathakurrajput44@gmail.com);

and (2.01 ppm), maximum organic carbon (0.74 %) was maximum observed in T<sub>6</sub> basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5%. Minimum days to 50% emergence (25.33), maximum plant height (31.66 cm, 53.33 cm at 30, 45 DAS respectively), number of leaves per plant (23.02 and 41.23 at 30 and 60 DAS respectively), fresh weight (62.06 g), dry weight (6.84 g), days to 50% flowering (37.33), number of pods per plant (15.79), shelling percentage (51.40 %), number of seeds/pod (8.46) individual pod weight (6.33 g), yield/plant (113.69 g), yield/plot (14.55 kg) and yield per hectare (194.00 q/ha) , highest net return (3,26,535) along with maximum benefit: net return per rupee invested (1:4.25) were recorded with the use basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5% in treatment T<sub>6</sub>. Basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5% was found highly beneficial for plant growth, yield and economics of garden pea.

**Keywords:** Zinc; manganese; growth; yield; garden pea.

## 1. INTRODUCTION

Garden Pea (*Pisum sativum* var. *hortense* L.) is a commercial legume vegetable crop of Himachal Pradesh as well as India. In Himachal Pradesh, it is grown during both main and offseason. It is a member of the Fabaceae family with a chromosome number  $2n = 14$ . Pea is a native of Europe and Northern Asia. Ethiopia is the likely the main centre of origin of garden pea. In India, Garden peas are grown over an area of 590 thousand hectares and are produced on an annual basis in the amount of 6182 million tonnes [1]. Garden Pea (*Pisum sativum* var. *hortense* L.) is a commercial legume vegetable crop of Himachal Pradesh as well as India. In Himachal Pradesh, area under pea cultivation is 26.00 thousand ha. production 328.80 million tonnes/ha [1].

Garden pea major producing states like Madhya Pradesh, Jharkhand, Himachal Pradesh, Punjab, West Bengal, Haryana, Bihar, Uttarakhand, Jammu & Kashmir, Odisha and portions of Rajasthan and Maharashtra the major pea growing states are Uttar Pradesh, Madhya Pradesh, Punjab, Himachal Pradesh, and Bihar. Major producing district in himachal Pradesh are Lahaul and Spiti, Kinnaur, Shimla, Kullu and Mandi districts in Himachal Pradesh are the principal pea-producing regions. The pea is a rich source of nutritional proteins. Being a legume crop it fixes atmospheric nitrogen into soil which helps in soil fertility status.

Himachal Pradesh is fifth and major leading pea producing state during main season, whereas major off season pea producing state of India. Micronutrients like zinc and manganese play an important role in increasing crop yield through their effect of plant itself and help in nitrogen

fixing symbiotic process. Garden pea is delicious and healthy for human consumption also good source of protein (7.2%) and other nutrients including slowly absorbed starch (50%) and sugars (12%), amino acids, vitamins A and vitamins C, calcium and phosphorus.

Micronutrients like zinc, manganese play an important role in increasing crop yield through their effect of plant itself and also help in nitrogen fixing symbiotic process. Zinc is an essential component of numerous physiological processes, including the production of chlorophyll, pollen, fertilisation, protein synthesis, cell elongation, and nodule formation. It also controls growth and development. Manganese (Mn) plays an important role in Nitrogen metabolism and forms several compounds needed or plant metabolism. Mn also acts as co-factor for several plant enzymes, where it activates about 35 different enzymes [2]. Keeping this in view, the present experiment is proposed to be undertaken to find out the effect of Zn and Mn on the growth, yield of garden pea.

## 2. MATERIALS AND METHODS

The present investigation was conducted during the Rabi season 2021-22 at the Agriculture Research Farm School of Agriculture, Abhilashi University, Mandi (H.P.), farm situated at 31°33'34" N latitude and 77°00'31" E longitude with an elevation of 1,426 m above mean sea level. The experiment was laid out in Randomized Block Design (RBD) Factorial with three replications comprising of seven treatment combination of Zn and Mn. The treatments were T<sub>1</sub>: Control (without Zn and Mn), T<sub>2</sub>: Basal application of Zn @ 50% of RDF + Mn @ 50% of RDF, T<sub>3</sub>: Basal application of Zn @ 100% of RDF + Mn @ 100% of RDF, T<sub>4</sub>: Basal application of

Zn @ 50% of RDF + foliar application of Zn @ 1%, T<sub>5</sub>: Basal application of Mn @ 50% of RDF + foliar application of Mn @ 0.5%, T<sub>6</sub>: Basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5% and T<sub>7</sub>: Foliar application of Zn @ 1% and Mn @ 0.5%. The plot size was 3 × 2.5 m<sup>2</sup> and a spacing of 30 cm × 15 cm was followed. Seeds of garden pea cv. 'Goldie' were sown in the bed on 12<sup>st</sup> November, 2021 and harvesting was done on 14<sup>th</sup> March, 2022. NPK, Zn and Mn were applied before garden pea was sown. Before the commencement of the experiment, soil samples were collected randomly from different plots of the experimental field from depth of 0-15 cm and the composite sample was prepared by mixing all these samples together. The soil sample after drying was passed through 2.0 mm sieve and was analyzed for soil pH, organic carbon and for available Zn and Mn. The results of the analysis and methods used are presented in Table 1. The recommended dose of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O were applied through urea, single super phosphate (SSP), muriate of potash (MOP) or Zinc (Zn) and manganese (Mn) respectively according to the treatment plan. The recommended dose of fertilizers like N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O were applied at the time of sowing in each plot as per the distributed to different treatments. Different growth and yield parameters like plant height, number of leaves per plant, number of pods per plant, average individual pod weight, yield per hectare. The statistical analysis was carried out by using the statistical package OPSTAT.

### 3. RESULTS AND DISCUSSION

#### 3.1 Soil Properties of Experimental Field (Post-Harvest)

Data pertaining to soil pH, OC (%), available Zn and Mn are present in (Table 1) showed that application of different level of zinc and manganese did not affected pH in the post-harvest soil. The treatments were found significant effect for organic carbon, available zinc and manganese. The result indicates that the combination of Zn and Mn significantly increase the organic carbon content in soil over control and the application of different level of Zn and Mn successively and significantly increased the available Zn and Mn content in soil.

The minimum pH value was recorded in T<sub>1</sub> control (without Zn and Mn) (5.94 pH) and

maximum pH was recorded in T<sub>6</sub> (5.98 pH) basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5%. Similar result for soil pH was observed by Alam (2020). In case of organic carbon, T<sub>6</sub> (basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5%) observed the highest organic carbon (0.74%) and T<sub>1</sub> [control (without Zn and Mn)] had the lowest organic carbon (0.53). This may be due to the fall in old leaves of the plant will leads to the amelioration of the crop residue, that will further accumulate the level of organic carbon [3].

Maximum available zinc (1.67 ppm) was recorded in T<sub>3</sub> (basal application of Zn @ 100% of RDF + Mn @ 100% of RDF) and minimum available zinc (0.67 ppm) recorded in T<sub>1</sub> [control (without Zn and Mn)]. Zinc concentration was significantly increased in soil when applied as soil and foliar application because of considerable exchange of Zn on the clay complexes with other cation on its addition. The results are in accordance with the findings of Kulandaivel et al., [4]. As a result of zinc foliar application, the soil zinc content clearly showed an increasing trend [5]. Maximum available manganese content (2.01 ppm) in soil was found in T<sub>3</sub> (basal application of Zn @ 100% of RDF + Mn @ 100% of RDF). Patel [6] reported that the soil application of manganese sulphate causes significant increase in Mn content of soil after harvest of crop.

#### 3.2 Plant Growth Attributes

Improvement of growth characters is pre-requisite to increased yield (Table 2). Treatment T<sub>6</sub> (basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5%) recorded a minimum days to 50% emergence (25.33), maximum plant height (31.66 cm and 53.33 cm at 30, 45 DAS respectively) and maximum number of leaves per plant 23.02 and 41.23 at 30 and 60 DAS respectively. The increase in plant height may be due accumulation of zinc nutrient in plant tissues and altered critical growth processes including photosynthesis and chlorophyll biosynthesis Stoyanova and Doncheva [5]. The reason for the maximum number of leaves can be the plant's cells dividing and lengthening more quickly. Numerous enzymatic processes involve zinc (Zn). It serves as a structural, functional, or regulatory co-factor.

**Table 1. Effect of Zn and Mn on soil properties**

Treatments	Soil pH	Organic carbon (%)	Available Zinc (ppm)	Available Mn (ppm)
T <sub>1</sub> : Control (without Zn and Mn)	5.94	0.53	0.67	0.78
T <sub>2</sub> : basal application of Zn @ 50% of RDF + Mn @ 50% of RDF	5.97	0.66	1.27	1.39
T <sub>3</sub> : basal application of Zn @ 100% of RDF + Mn @ 100% of RDF	5.96	0.70	1.67	2.01
T <sub>4</sub> : basal application of Zn @ 50% of RDF + foliar application of Zn @ 1%	5.97	0.60	1.13	1.17
T <sub>5</sub> : basal application of Mn @ 50% of RDF + foliar application of Mn @ 0.5%	5.96	0.64	1.61	1.77
T <sub>6</sub> : basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5%	5.98	0.74	1.50	1.53
T <sub>7</sub> : foliar application of Zn @ 1% and Mn @ 0.5%.	6.00	0.68	1.19	1.23
SE(m) ±	0.013	0.002	0.004	0.013
CD at 5%	NS	0.003	0.011	0.040

**Table 2. Effect of Zn and Mn practices in growth parameters of garden pea**

Treatments	Days to 50% emergence	Plant height (cm)		Number of leaves		Fresh weight of plant after harvesting (g)	Dry weight of plant after harvesting (g)
		30 DAS	45 DAS	30 DAS	60DAS		
T <sub>1</sub> : Control (without Zn and Mn)	34.00	18.66	38.66	15.33	24.08	44.55	4.78
T <sub>2</sub> : basal application of Zn @ 50% of RDF + Mn @ 50% of RDF	30.33	23.66	46.66	19.66	33.18	53.16	5.99
T <sub>3</sub> : basal application of Zn @ 100% of RDF + Mn @ 100% of RDF	27.33	29.33	51.05	22.33	38.48	58.10	6.54
T <sub>4</sub> : basal application of Zn @ 50% of RDF + foliar application of Zn @ 1%	32.66	20.00	41.33	18.00	27.12	47.50	4.90
T <sub>5</sub> : basal application of Mn @ 50% of RDF + foliar application of Mn @ 0.5%	31.00	22.03	43.99	18.66	29.01	50.72	5.32
T <sub>6</sub> : basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5%	25.33	31.66	53.33	23.02	41.23	62.06	6.84
T <sub>7</sub> : foliar application of Zn @ 1% and Mn @ 0.5%.	29.33	26.00	49.33	21.03	35.92	56.11	6.29
SE(m) ±	1.10	1.46	0.77	1.24	1.21	0.74	0.10
CD at 5%	3.42	4.56	02.41	3.92	3.77	2.32	0.31

**Table 3. Effect of Zn and Mn practices in yield parameters of garden pea**

Treatments	Days to 50% flowering	Number of pods per plant	Shelling percentage (%)	Number of seeds per pod	Average pod weight (g)	Yield/plant (g)	Yield/plot (kg)	Yield per hectare (q)
T <sub>1</sub> : Control (without Zn and Mn)	45.00	11.50	38.35	5.16	4.95	65.27	8.52	113.61
T <sub>2</sub> : basal application of Zn @ 50% of RDF + Mn @ 50% of RDF	41.33	13.49	44.68	6.60	5.99	100.83	12.46	166.13
T <sub>3</sub> : basal application of Zn @ 100% of RDF + Mn @ 100% of RDF	39.66	14.24	48.74	7.97	6.23	108.69	13.65	182.00
T <sub>4</sub> : basal application of Zn @ 50% of RDF + foliar application of Zn @ 1%	44.33	12.59	42.34	5.43	5.71	80.91	8.75	116.74
T <sub>5</sub> : basal application of Mn @ 50% of RDF + foliar application of Mn @ 0.5%	42.33	13.22	43.76	5.63	5.92	88.31	11.96	159.51
T <sub>6</sub> : basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5%	37.33	15.79	51.40	8.46	6.33	113.69	14.55	194.00
T <sub>7</sub> : foliar application of Zn @ 1% and Mn @ 0.5%.	40.00	14.06	46.86	6.88	6.13	103.26	13.01	173.47
SE(m) ±	1.49	0.37	1.29	0.25	0.10	1.36	0.32	2.93
CD at 5%	4.65	1.17	4.03	0.78	0.31	4.24	1.02	8.90

**Table 4. Effect of Zn and Mn on cost benefit ratio (CBR)**

Treatments	Cost of cultivation (₹/ha)	Gross Return (₹/ha)	Net Return (₹/ha)	Net return per rupee invested
T <sub>1</sub> : Control (without Zn and Mn)	94,258	249,942	1,55,684	1:2.65
T <sub>2</sub> : Basal application of Zn @ 50% of RDF + Mn @ 50% of RDF	97,223	365,486	2,67,063	1:3.75
T <sub>3</sub> : Basal application of Zn @ 100% of RDF + Mn @ 100% of RDF	1,00,189	400,400	2,99,011	1:3.99
T <sub>4</sub> : Basal application of Zn @ 50% of RDF + foliar application of Zn @ 1%	99,056	268,488	1,69,432	1:2.71
T <sub>5</sub> : Basal application of Mn @ 50% of RDF + foliar application of Mn @ 0.5%	97,474	350,922	2,53,448	1:3.60
T <sub>6</sub> : Basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5%	1,00,265	426,800	3,26,535	1:4.25
T <sub>7</sub> : Foliar application of Zn @ 1% and Mn @ 0.5%	97,301	381,634	2,84,333	1:3.92

The maximum fresh weight was observed (62.06 g) in T<sub>6</sub> (basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5%) Zinc stimulates the action of various enzymes and photosynthetic pigments, which promotes plant vegetative development [7]. Additionally, it affects protein synthesis, energy transfer, and nitrogen metabolism by altering the necessary enzymes [8] and maximum dry weight (6.84 g) observed in T<sub>6</sub> (basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5%).

### 3.3 Yield Attributes

Minimum days to 50% flowering (37.33), maximum number of pods/plant (15.79), shelling percentage (51.40) were observed in T<sub>6</sub> (basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5%). The increase in number of pods due to the application of micronutrients especially Zinc sulphate has a positive effect on formation of stamens and pollens and we can attribute the increase of number of pods per plant to this property of micronutrients. The large number of seed (8.46) and average pod weight were observed in T<sub>6</sub> (basal application of Zn @50% of RDF and Mn @50% of RDF + foliar application of Zn @1% Zinc was applied as a foliar spray, which enhanced seed weight. Seed weight will increase when photosynthesis assimilates is transferred from vegetative organs to the other portions.

The highest yield/plant (113.69 g), yield /plot (14.55 kg) and yield /hectare (194.00 q) observed in T<sub>6</sub> (basal application of Zn @50% of RDF and Mn @50% of RDF + foliar application of Zn @1% this is might due to the Zinc promotes male and female gametogenesis, which increases the number of flowers per plant, zinc is essentially highly advantageous for the reproductive yield of the crop. The formation of sporogenous tissue is additionally stimulated by zinc treatment, which raises the quantity of pollen grains per anther. These effects result in proper pollen grain germination, normal development and an increase in yield parameters such as the number, size, and weight of pods and seeds [9,10].

**Cost Benefit Ratio (CBR):** The Cost Benefit Ratio (CBR) of the various treatment combinations have been presented in Table 4. The data revealed that highest cost of production

₹/hectare (₹ 1,00,265), maximum gross income ₹/hectare amounting to (₹ 426,800), highest net return ₹/hectare (₹ 326,535), maximum net return/rupee invested (1:4:25) was incurred in treatment T<sub>6</sub> (basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5%), whereas lowest cost of cultivation ₹/hectare (₹ 94,258), minimum gross income ₹/hectare (₹ 249,942), lowest net return ₹/hectare (₹ 155,684), net return per rupee invested (1:2.65) was observed in treatment T<sub>1</sub> control (without Zn and Mn).

### 4. CONCLUSION

Based on present investigation, it can be concluded that the basal application of Zn @ 50% of RDF and Mn @ 50% of RDF + foliar application of Zn @ 1% and Mn @ 0.5% is more beneficial to the growth and yield of garden pea than the control or basal application only. It improves availability of nutrients in term with higher yield.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Anonymous. Department of Agriculture and Farmers Wekfare, horticulture statistics; 2022. Available:<https://agricoop.nic.in/en/StatHortEst>
2. Mengel K, Kirkby EA. Principles of plant nutrition 5<sup>th</sup>ed. Dordrecht: Kluwer Academic Publishers. 2001; 849.
3. Chethan KV, David AA, Thomas T, Swaroop N, Rao S and Hassan A. Effect of different levels of NPK and Zn on physico-chemical properties of soil growth parameters and yield by pea (*Pisum sativum* L.) Cv. Rachana Journal of Pharmacognosy and Phytochemistry. 2018;7:2212-2215.
4. Kulandaivel S, Mishra BN, Gangiah B, Mishra PK. Effect of levels of zinc and iron and their chelation on yield and soil micronutrient status in hybrid rice (*Oryza sativa*)- wheat (*Triticum aestivum*) cropping system. Indian J. Agron. 2004;49(2):80-83.
5. Stoyanova Z, Doncheva S. The effect of zinc supply and succinate treatment on plant growth and mineral uptake in pea plant Brazilian Journal of Plant Physiology. 2002;14:111-116.

6. Patel Rk. Effect of manganese application on growth, yield and its agronomic biofortification in Wheat (*Triticum aestivum* L.) in typic ustochrept of Ananad. M.Sc. (Ag) thesis submitted to Anand Agriculture University, Anand, Gujrat; 2013.
7. Sadeghzadeh B. A review of zinc nutrition and plant breeding. Journal of soil Science and Plant Nutrition. 2013;13:905-927.
8. Hafeez BM, Khanif YM, Saleem M. Apr. Role of zinc in plant nutrition-a review American Journal of Experimental Agriculture. 2013;1:3:374.
9. Pandey N, Gupta B. Improving seed yield of black gram (*Vigna mungo* L. var. DPU-88-31) through foliar fertilization of zinc during the reproductive phase. Journal of Plant Nutrition. 2012;1:35:1683-1692.
10. Alam I, Paul AK, Sultana S and Bithy PA. Effect of zinc and molybdenum on the growth and yield of garden pea (*Pisum sativum* L.). International Journal of Bio-resource and Stress Management. 2020; 11:425-431.

© 2023 Heena et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/105198>