



# Formation of Male Sex Tilapia Larvae (*Oreochromis niloticus*) Using Natural Methyltestosterone Hormone

Atriani <sup>a</sup>, Ardiansyah <sup>a\*</sup>, Hasniar <sup>a</sup>, Indrayani <sup>b</sup>,  
Adnan Baiduri <sup>c</sup> and Suryati <sup>c</sup>

<sup>a</sup> Fish Hatchery Study Program, Aquaculture Department, Pangkep State Polytechnic of Agriculture, Indonesia.

<sup>b</sup> Study Program of Agricultural Technology Education, Faculty of Engineering Makassar State University, Indonesia.

<sup>c</sup> Aquaculture Technology Study Program, Aquaculture Department, Pangkep State Polytechnic of Agriculture, Indonesia.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/AJBGMB/2024/v16i2359

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113156>

Original Research Article

Received: 07/12/2023

Accepted: 10/02/2024

Published: 13/02/2024

## ABSTRACT

**Aims:** Obtaining a productive tilapia, by changing the sex of the tilapia larvae into male monosex. This was achieved by determining the effect of the duration of administering the natural methyltestosterone (MT) hormone on the male sex of tilapia using the Immersion method.

**Methodology:** The study was conducted at the Northern Region Marine and Fisheries Service Branch, Wanayasa Freshwater Fish Seed Development Center, Purwakarta Regency, West Java Province, Indonesia. The experimental design used a completely randomized design (CRD) with 5 treatments and 3 replications. The study used the Immersion method where untreated tilapia and

\*Corresponding author: E-mail: ardi\_kimsan@yahoo.com;

the four hormones tilapia. The treatments are as follows:

A = untreated tilapia

B = tilapia treated in 17 $\alpha$ -MT (5h-100mg/L)

C = tilapia treated in MT (5h-350mg/L)

D = tilapia treated in MT (10h-350mg/L)

E = tilapia treated in MT (15h-350mg/L)

**Results:** Tilapia larvae used were aged 6 days after hatching and reared for 40 days, then the male and SR sex were observed. The results of this study showed that the duration of immersion of tilapia larvae using natural MT significantly affected the male tilapia which resulted in the highest male percentage at 15 h of immersion of 80.2%, then followed by 10-h immersion of 75.6% with the same SR value of about 90%.

**Conclusion:** Administration of natural MT using cow testis meal by immersion method at different durations significantly affects male sexual development in 6-day-old tilapia larvae. Based on the obtained data analyses, the percentage of male tilapia in the 40-day experimental study shows the most effective duration in treatment E (15 h) with a measure of 350 mg/l, which is 80.2%. Further studies regarding the relationship between sex change and the growth of sex-changed fish as an effect of immersion duration.

*Keywords: Bovine testicle; cow testicle flour; monosex culture; natural methyltestosterone; percentage of males; tilapia fish larvae.*

## 1. INTRODUCTION

Tilapia is ranked as one of the top ten major products from the freshwaters with significant economic value according to the Department of Marine Affairs and Fisheries of the Republic of Indonesia [1]. Growing demand and widespread tilapia aquaculture globally have brought tilapia to the world's attention as an affordable supplier of healthy animal protein, with total U.S. tilapia imports during the first half of 2022 reaching 89,104 tons, up 9.7 percent annually [2]. Hence, numerous studies are still being conducted in Indonesia, in an attempt to produce productive tilapia [3].

Tilapia is ranked as one of the top ten major products from the freshwaters with significant economic value and remains one of the most reasonably priced sources of animal protein for people of all social classes. Nevertheless, the high reproductive potential of tilapia fish makes it difficult to regulate and frequently results in inbreeding, which is most prevalent in tilapia aquaculture (Mangaro et al., 2018). This causes fish to grow slowly, taking a long time to reach the size needed for consumption, and sometimes even experiencing stalled growth (Tatalede et al., 2019; Maulana et al., 2023). To boost tilapia farming output, the most effective way to do so is to produce male tilapia larvae for mono-sex production (Malik et al., 2019; Susilawati et al., 2022). This is because, during the tilapia growing process, male tilapia grow more quickly than female tilapia.

One procedure to get male mono-sex tilapia is through sex change procedures by administering male hormones (testosterone). The commonly utilized hormone is the manufactured hormone 17 $\alpha$ -MT with a success rate of male sex alter that can reach 96-100% [4], (Mlalila et al., 2015). In addition, masculinization can also be done by controlling the environmental temperature in the form of increased temperature [5]. The higher the temperature, the higher the male-sex ratio (Tessema et al., 2006). Abou-El Fotoh et al. [6] found that a temperature of 28°C produced male tilapia by 52.33% while a temperature of 36°C produced male tilapia by 81%, thus it can be concluded that temperature and sex change are directly proportional.

This study aimed to determine the effect of the duration of the natural Methyltestosterone (MT) hormone treatment on the formation of the male sex tilapia. The expected hypothesis is that immersion of larvae in natural MT hormone with different immersion durations affects the increased percentage of male sex formation in tilapia.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

The research was conducted between October and December 2021 at the Northern Region Marine and Fisheries Service Branch, Wanayasa Freshwater Fish Seed Development Center, Purwakarta Regency, West Java Province, Indonesia.

## 2.2 Experimental Design

The study used a completely randomized design with 5 treatments and 3 replications. The treatments are as follows:

- A = untreated tilapia
- B = tilapia treated in 17 $\alpha$ -MT (5h-100mg/L)
- C = tilapia treated in MT (5h-350mg/L)
- D = tilapia treated in MT (10h-350mg/L)
- E = tilapia treated in MT (15h-350mg/L)

## 2.3 Experimental Set-Up

The test fish used in this study were tilapia larvae aged 6 days after hatching (weight between 0.02 - 0.03 g/larvae), the total number of larvae was 750 with a density of 50 larvae/container, and the larvae were obtained from natural spawning. The natural MT hormone used was derived from fresh cow testicles processed into cow testicle flour that had been previously prepared. The immersion method was made by dissolving the hormone in 70% alcohol in the amount of 3 ml and then put into an immersion jar containing 1 L of water then aerated and homogenized for  $\pm$  5 min to remove the alcohol content. Tilapia larvae were immersed according to each treatment and then transferred to a rearing container in the form of a pond that was blocked using nets and fed with pellet (Hi-Pro-Vite brand, type PSP) with a frequency of 3 to 4 times/day ad libitum. After 40 days of rearing [7], gonad observations were made by dissecting the abdomen of the fish then the gonads were taken and given 2-3 drops of Eosin solution on a glass object and then observed using a binocular microscope.

## 2.4 Larval Rearing

After the immersion, the larvae were transferred into the rearing container at a density of 1 ind/L (Adel et al., 2006). The larvae were fed with pellet starter in flour form for the freshwater fish fry, Hi-Pro-Vite brand, PSP type. During rearing, feeding was applied ad-satiation method with a frequency of 3 to 4 times a day. The rearing was carried out until the fish reached 40 days of age.

## 2.5 Parameters Observed

### 2.5.1 Percentage of male tilapia fish

The percentage of male tilapia is the percentage of the number of male tilapia fish from the total population of the tested tilapia fish. Damayanti et

al. [8], stated that to calculate the percentage of male sex using the following formula.

$$(\%) \text{ Male} = \frac{\text{Number of males}}{\text{Total number of fish}} \times 100\%$$

### 2.5.2 Survival rate

Survival is the percentage of the number of fish that live at the end of the experiment from the total fish population at the beginning of the experiment. The percentage of fish survival rate can be calculated with the formula proposed by Huisman [9]:

$$\text{SR} (\%) = \frac{N_t}{N_o} \times 100\%$$

Where SR is the survival rate of fish during rearing (%),  $N_t$  is the number of fish at the end of rearing (fish), and  $N_o$  is the number of fish at the beginning of rearing (fish).

### 2.5.3 Measurement of water quality

Water quality measurements during maintenance include measurements of water temperature, dissolved oxygen (DO), and acidity (pH). Measurements were taken twice a day at 8 am and 4 pm for 40 days. Water temperature was measured using a mercury thermometer, pH was measured using a digital pH meter and DO was measured using a digital DO meter.

## 2.6 Data Analysis

The data obtained were analyzed by the one-way Analysis of Variance (ANOVA) method. If the results show a real effect, it is continued with the smallest real difference test (BNT) to determine whether there are significant differences between treatment groups.

## 3. RESULTS AND DISCUSSION

### 3.1 Percentage of Male Fish

Based on the observation, the percentage of male tilapia treated with natural MT through immersion (72.8 - 80.2%) was higher than untreated tilapia (47.8 $\pm$ 0.02%). Data on the percentage of male tilapia in each treatment at the end of the experiment can be seen in Table 1.

**Table 1. Percentage (100%) of sex of male tilapia at the end of the experiment**

Replications	Treatments				
	A C-	B C+ (5-h)	C (5-h)	D (10-h)	E (15-h)
I	47.7	73	72.7	74.5	77.1
II	46.8	78	72.3	76.7	77.8
III	48.9	70.5	73.3	75.6	85.7
Mean	47.8±0.02 <sup>a</sup>	73.8±0.03 <sup>bc</sup>	72.8±0.02 <sup>bc</sup>	75.6±0.04 <sup>cd</sup>	80.2±0.04 <sup>d</sup>

*Different letters a, b, c, and d indicate significant differences between treatments*

where A = untreated tilapia, B = tilapia treated in 17a-MT (5h-100mg/L), C = tilapia treated in MT (5h-350mg/L), D = tilapia treated in MT (10h-350mg/L), E = tilapia treated in MT (15h-350mg/L)

The table above shows a significant difference ( $P < 0.05$ ) in the percentage value of male tilapia treated with natural MT hormone at different immersion durations. The 15-h treatment produced significantly different results than the untreated tilapia (C-), treatment B (C+), and treatment C (MT-5-h). However, the results were not significantly different from the 10-h treatment. It can be seen that the sex change of tilapia fish into males happens after immersion in natural MT and is regulated by the immersion duration of tilapia larvae, with a tendency to increase the percentage of male fish produced as the immersion time increases.

### 3.2 Survival of Tilapia

Survival of tilapia larvae is crucial for determining the effectiveness of treatment methods. The tilapia survival data collected at the end of this study is calculated by subtracting the number of fish at the start of the study from the number of fish that survive at the end of the study. Table 2 shows tilapia survival results for each treatment.

From the table above, the survival rate of tilapia in each treatment using cow testicle flour at different immersion durations did not bring a significant effect ( $P > 0.05$ ) throughout the treatment period until the end of the experiment. However, when the SR of fish treated with the natural hormone MT showed no statistically significant effect on untreated tilapia (C-) treatment, it was significantly different compared to C+ (treatment by immersing the hormone 17a-MT).

### 3.3 Water Quality

Water quality is one of the factors that support the survival of fish, but it also affects the formation of male sex when tilapia is still in the period of sexual differentiation. Water quality parameters in this study measured during the experiment included DO (4 - 6.8 mg/L), temperature (24 - 27), and pH (6.3 - 7.9). According to the results of water quality measurements during the experiment, they are still within the normal range that tilapia can tolerate, thus possibly supporting the survival rate of tilapia. Measurement data are shown in Table 3.

**Table 2. Percentage of experimental fish survival at the end of the experiment**

Replications	Treatments				
	A C-	B C+ (5-h)	C (5-h)	D (10-h)	E (15-h)
I	88	74	88	94	96
II	94	82	94	86	90
III	90	88	90	90	84
Mean	90,7 <sup>bc</sup>	81,3 <sup>a</sup>	90,7 <sup>bc</sup>	90 <sup>ab</sup>	90 <sup>ab</sup>

*Different letters a, b, and c indicate significant differences between treatments*

where A = untreated tilapia, B = tilapia treated in 17a-MT (5h-100mg/L), C = tilapia treated in MT (5h-350mg/L), D = tilapia treated in MT (10h-350mg/L), E = tilapia treated in MT (15h-350mg/L)

**Table 3. Water quality of fish-rearing media during the study**

Parameter Measured	Unit	Measurement Results	
		(8 am)	(4 pm)
Dissolved Oxygen	mg/L	4,0 – 6,8	4,0 – 6,8
Temperature	°C	24 – 27	24 – 28
pH	-log[H <sup>+</sup> ]	6,3 – 7,9	6,4 – 7,9

#### 4. DISCUSSION

In this study, the percentage of male tilapia that received natural MT hormone ranged from 72.8 to 80.2%. The administration of natural MT has been shown to influence the hormonal system in the body of tilapia [10], with each treatment increasing the percentage of different male tilapia. Further explained by Zairin [4] and Mlalila et al. (2015), MT hormone administration in the sex direction by the immersion method (dipping) is more efficient because the dose given is relatively small and the contact time is shorter. Cow testicle flour contains the testosterone hormone which is thought to increase the concentration of androgen hormones in tilapia larvae thus they become male. This is supported by the statement of Iskandar [11] that the administration of cow testicle flour containing the hormone testosterone through immersion applications, can affect the hormonal system in the body of fish thus it affects the formation of male tilapia.

Furthermore, Iskandar [11] and Wahidah et al. [12] explained that the administration of cow testicle flour extract containing the hormone testosterone through immersion can affect the hormonal system in the body of fish thus affecting the formation of male tilapia. The immersion method is usually applied to the sex-change of fish larvae [11,12], in which, the fish larvae used are still in the differentiation phase because other methods such as feeding and injection are not effectively applied the small size of the larvae.

In this study, the total percentage of males in treatment C (MT with 5-h immersion) was 72%, which was smaller than treatment B (17 $\alpha$ -MT) of 73.8%. It is presumed that the cause of the low percentage of male fish produced in treatment C is the presence of other factors such as the material used is still in the form of coarse flour which is not soluble in water therefore, it cannot be absorbed optimally by tilapia larvae through the skin, gills, and digestive system. However, the percentage of male fish produced is still higher than in previous studies, where Ridwan

[13] reported 72.14% male fish, previously Phelps and Popma [14] reported 65% male fish. The result of the present study indicates the use of cow testicle flour is more effective than the use of fresh cow testicles as used in the previous studies.

The success of the tilapia male sex direction through the immersion method is influenced by several factors including size, age, length of treatment, environment, fish species, genetics, hormone type, hormone dose, and treatment time [15]. The application of sex direction techniques is generally performed on fish that are still in the process of differentiation [16], in this labile period, fish sex can be directed through hormone induction [17]. The period of gonad differentiation in tilapia occurs up to 30 min after hatching. Still, the most sensitive period for sex direction in tilapia is 7-14 days after hatching [18] and according to Yuniarti et al. [19], sex differentiation in tilapia occurs when the larvae are 6-7 days after hatching until about 27-28 days after hatching.

In this study, the results of SR analysis of experimental fish were still relatively high, ranging from 81.3% to 90.7% compared to the report of Meyer et al. [20], about 40.2% using fresh cow testicles, Yustiati et al. [21] reported 86% using oven-dried cow testicles, and the report of Shalaby et al. [22], around 60-70% using the synthetic hormone 17 $\alpha$ -MT orally, and by Yustina et al. [23], about 14.66% in bonylip barb (*Osteochilus vittatus*) using the synthetic hormone 17 $\alpha$ -MT by immersion method. Furthermore, Ridha and Lone [24] emphasized that the use of synthetic hormones including 17 $\alpha$ -MT causes higher mortality. In addition, the use of chemicals in sex change can affect the survival rate of fish because it is believed that treated fish are subjected to physiological stress. The experimental fish are still in the larval stage and thus their physiological state is still weak, which also means that these synthetic materials that enter the body can have a two-step effect on the antibody system, which can initially stimulate the production of antibodies, but the next effect is to suppress immune responses. Long-term

administration of synthetic substances, even in low doses, can impair the ability of immune cells to reproduce (spread) [25]. Cow testicle flour, on the other hand, is a natural ingredient that does not cause physiological stress to the tilapia larvae, so the survival rate is relatively high.

In addition, cow testicle flour also contains a variety of essential and non-essential amino acids and contains saturated and unsaturated fatty acids [26]. The amino acids include arginine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, and valine, while non-essentials include aspartate, alanine, glutamate, glycine, Florine, serine, cysteine, and tyrosine. These amino acids are very useful in the formation of androgenic hormones, testosterone, which plays a role in increasing libido and the formation of spermatozoa [27,28].

Based on Table 3, the water quality during the study varied from the normal tolerance range of tilapia to very low water temperature. This means that one of the factors responsible for the low percentage of male tilapia fish obtained is the water quality of the rearing environment, especially the temperature. Rohmaniah et al. [29], the higher the water temperature, the faster the fish's metabolism accelerates, resulting in faster oxygen uptake and respiration and less dissolved oxygen in the water. Without oxygen, fish easily become stressed and even die. Temperature changes can affect the activity of the testosterone hormone; For example, low temperatures reduce the effect of the testosterone hormone, resulting in more female fish than male fish [30,31].

According to Chen et al. [32], water temperature affects larval development stages as well as the duration of gonadal sensitivity to hormone stimulation. Water temperature has a direct impact on the body's metabolic processes, which in turn alter the function of hormones. Meanwhile, Afpriyaningrum [33] found that tilapia larvae absorbed testosterone levels at 36°C during a 4-h immersion, resulting in 92.50% of males. The optimal water pH for fish survival ranges from 6.5 to 9, and DO levels vary from 5.5 to 6.7 mg/l. Maimunah dan Kilawati [34] recommend an optimal environment for tilapia rearing with DO 5.3-8.8 mg/l, pH 7.3-8.3, and water temperature 26-30°C. Nivellet et al. [18] recommend a water temperature of 27-30°C. However, in this study, the water temperature fluctuated between 24 and 28°C.

## 5. CONCLUSION

It can be concluded that the administration of natural MT by utilizing cow testicle flour through the immersion method at different durations has a significant effect on the formation of male sex in tilapia larvae aged 6 days. Based on the data analysis that has been obtained, the percentage of male tilapia sex during 40 days of the experimental study shows the most effective duration is in treatment E (15 h) with a dose of 350 mg/L, which is 80.2%.

## ACKNOWLEDGMENTS

The author would like to thank the head of the Wanayasa freshwater fish seed development center for the laboratory facilities and equipment used by the author during the research. To all center staff who helped the authors from planning, and experiment setup to data analysis.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Department of Marine Affairs and Fisheries of the Republic of Indonesia. Prohibited Drugs. Directorate General of Aquaculture, Center for the Development of Freshwater Fish Cultivation. Sukabumi; 2008.
2. FAO. Demand for tilapia resumes growth despite higher prices. Food and Agriculture Organisation for the UN; 2023. Available: <https://www.fao.org/in-action/globefish/market-reports/resource-detail/en/c/1632900/>
3. Hertanto MA. Male red tilapia (*Oreochromis niloticus*) production using forest bee honey. Thesis. Fakultas Teknologi Universitas Atma Jaya, Yogyakarta; 2013.
4. Zairin MJ. Endocrinology and its role for the future of Indonesian fisheries. Scientific oration of inauguration of permanent professor of reproductive physiology and endocrinology of aquatic animals. Institut Pertanian Bogor, Bogor; 2003.
5. Bowman MP, Bowker J, Carty DG, Straus DL, Farmer BD, Mitchell AJ, Ledbetter CK. The safety of 17 $\alpha$ -methyltestosterone administered in feed to larval Nile tilapia. AADAP. Drug Research Information Bulletin. Drib no. 2012;28.
6. Abou El-Fotoh EM, Ayat MS, Abd El Rahman GA, Farag ME. Mono-sex male

- production in Nile tilapia (*Oreochromis niloticus*) using different water temperatures. Zagazig Journal of Agricultural Research. 2014;41:1-8.
7. Laheng S, Widyastuti A. Effect of soaking time using coconut water on masculinization of masamo catfish (*Clarias sp*). Acta Aquatic: Aquatic Sciences Journal. 2019;6:2.
  8. Damayanti AA, Sutresna W, Wildan. Honey application for tilapia sex direction (*Oreochromis niloticus*). Depik. 2013;2(2):82-86. ISSN:2089-7790.
  9. Huisman EA. Principle of fish production. Department of fish culture and fisheries. Wageningen Agricultural University; 1987.
  10. Sarker B, Das B, Chakraborty S, Hossain MA, Alam MMM, Mian S, Iqbal MM. Optimization of 17 $\alpha$ -methyltestosterone dose to produce quality mono-sex nile tilapia *Oreochromis niloticus*. Heliyon. 2022, Dec 9;8(12):e12252. DOI: 10.1016/j.heliyon.2022.e12252
  11. Iskandar A. Effectiveness of bovine testicle flour extract in the sex change of tilapia *oreochromis niloticus* through immersion technique. Thesis. Faculty of Fisheries and Marine Science, Bogor Agricultural University, Bogor; 2010.
  12. Wahidah, Arham Rusli, Muh Alias LR, Amrullah. The morphological characters of Nile tilapia (*Oreochromis niloticus*) produced with bull testicle powder extract application AACL Bioflux. 2021;14(3):1811-1817. Available:http://www.bioflux.com.ro/aacl.
  13. Ridwan. Effect of immersion time of cow testicle flour on masculinization of tilapia (*Oreochromis niloticus*) larvae. Thesis. Universitas Muhammadiyah Makassar; 2022.
  14. Phelps RP, Popma TJ. Sex reversal of tilapia. tilapia. Pages 34-59 in B.A. costapierce and J.E. Rakocy, eds. Tilapia aquaculture in The Americas, The World Aquaculture Society, Baton Rounge. Lousiana, United States. 2000;2.
  15. Muslim. Masculinization of tilapia (*Oreochromis niloticus*) by feeding bovine testicle flour. Thesis Graduate School. IPB; 2011.
  16. Fernandino JI, Hattori RS. Sex determination in neotropical fish: Implications ranging from aquaculture technology to ecological assessment. General and Comparative Endocrinology. 2019;273:172-183. Available:https://doi.org/10.1016/j.ygcen.2018.07.002.
  17. Baroiller JF, D'Cotta H. The reversible sex of gonochoristic fish: Insights and consequences. Sex Dev. 2016;10(5-6):242–266. Available:https://doi.org/10.1159/000452362
  18. Nivelle R, Gennotte V, Kalala EJK, Ngoc NB, Muller M, Mélard C, Rougeot C. Temperature preference of Nile tilapia (*Oreochromis niloticus*) juveniles induces spontaneous sex reversal. PLoS One. 2019;14(2):e0212504. DOI: 10.1371/journal.pone.0212504.
  19. Yuniarti T, Sofi Hanif Teguh Prayoga, dan Suroso. Tilapia broodstock production technique. Jurnal Budidaya Air Tawar. 2007;4:32-36.
  20. Meyer D, Marco GW, Chan dan Castillo C. Use of fresh bull and hog testis in sex reversal of nile tilapia fry. Aquaculture Collaborative Research Support Program United State Agency for International Development (USAID). Honduras; 2008.
  21. Yustiati A, Bangkit I, Zidni I. Masculinization of nile tilapia (*Oreochromis niloticus*) using extract of bull testes. Conference Series Earth and Environmental Science. 2018;139(1):012008. DOI: 10.1088/1755-1315/139/1/012008
  22. Shalaby AME, Ashraf AR dan Yassir AEK. Sex reversal of nile tilapia fry using different doses of 17 $\alpha$ -methyltestosterone at different dietary protein levels. Central Laboratory for Aquaculture Research, Abbassa, Abo-Hammad, Sharkia Governorate, Egypt; 2007.
  23. Yustina, Arnetis D, Ariani. Effectiveness of sand sea cucumber flour (*Holothuria scabra*) on masculinization of cupang fish (*Betta splendens*). Jurnal Biogenesis. 2012;9(1):6773.
  24. Ridha M, Lone K. Effect of oral administration of different levels of 17 $\alpha$ -methyltestosterone on the sex reversal, growth and food conversion efficiency of the tilapia *Oreochromis spilurus* (Günther) in brackish water. Aquaculture Research. 2008;21(4):391–397. DOI: 10.1111/j.1365-2109.1990.tb00477.x
  25. Mariani E, Lisignoli G, Borzi RM, Pulsatelli L. Biomaterials: Foreign bodies or tuners for the immune response?. Int J Mol Sci. 2019;20(3):636.

- DOI: 10.3390/ijms20030636.
26. Odin RY, Bolivar RB, Liping L, Fitzsimmons K. Masculinization of Nile tilapia (*Oreochromis niloticus* L.), using lyophilized testes from carabao (*Bubalus bubalis* carabanesis L.), bull (*Bos indicus* L.) and boar (*Sus domesticus* L.). In Better science, better fish, better life. Proceedings of the Ninth International Symposium on Tilapia in Aquaculture, Shanghai, China. Aquafish Collaborative Research Support Program. 2011, April 22-24;105-120.
  27. Hutagalung RA. 2020. Differences in Sex Reversal Methods Using Cow Testicle Flour on Masculinization of Red Tilapia (*Oreochromis niloticus*). *Manfish Journal of Marine, Environment and Fishries*. 2020;1(1):9-14.
  28. Pearson K. Tilapia Fish: Benefits and Dangers. Nutrition. Healthline; 2023. <https://www.healthline.com/nutrition/tilapia-fish>
  29. Rohmaniah H, Syaputra Dn dan Syarif AH. Tilapia (*Oreochromis niloticus*) masculinization using Java chili extract (*Piper retrofractum*) through larval immersion. *Journal of Aquaculture Asia*. 2019;4(1):29-34
  30. Huda NR, Susilowati T, Yuniarti T. Application of Cow Testicle Flour Containing rGH in Artificial Feed on Sex Ratio, Growth and Survival of Tilapia (*Oreochromis niloticus*). *Pena Akuatika* 2018;17(2):59-69.
  31. Suyanto R. Nile Tilapia. PT. Penebar Swadaya. Jakarta; 2004.
  32. Chen H, Hu Q, Kong L, Rong H, Bi B. Effects of temperature on the growth performance, biochemical indexes and growth and development-related genes expression of juvenile hybrid sturgeon (*Acipenser baerii*♀ × *Acipenser schrenckii*♂). *Water*. 2022;14:2368. Available:<https://doi.org/10.3390/w14152368>
  33. Afpriyaningrum MD, Soelistyowati DT, Alimuddin, Zairin Jr M, Setiawati M, Hardiantho D. Tilapia masculinization by larval immersion at 36°C and residual 17 $\alpha$ -Methyltestosterone levels in fish body. *Omni-Akuatika*. 2016;12(3):106-113.
  34. Maimunah Y, Kilawati Y. Performance of growth in tilapia fish in policulture system. *J Food Life Sci*. 2020;4(1):42-49.

© 2024 Atriani et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/113156>