



# Management of a Bilateral Ectopia Lentis in a Patient with Marfan Syndrome: A Case Report

EL Filali El Mehdi <sup>a\*</sup>, Zineb Algouti <sup>a</sup>, Farah Benelkadri <sup>a</sup>,  
Houssaine Aitlhaj <sup>a</sup>, Mohamed Kriet <sup>a</sup> and Fouad El Asri <sup>a</sup>

<sup>a</sup> Department of Ophthalmology, Cadi Ayyad University, MED VI University Hospital, Avicenne Military Hospital, Marrakech, Morocco.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/OR/2024/v19i3424

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115465>

Case Report

Received: 03/02/2024

Accepted: 07/04/2024

Published: 29/04/2024

## ABSTRACT

**AIMS:** to highlight the management procedure of a bilateral subluxated lens in a patient with marfan syndrome.

**Case Report:** We report a case of a 51-year-old woman, with a family history of Marfan syndrome; who presented for a bilateral subluxated lens, iridodonesis and bilateral macular degeneration. Given the family history, the general physical features and the ocular findings, diagnostic of Marfan syndrome was established. The management of the ectopic lentis was very challenging. An intracapsular lens extraction with posterior chamber intraocular lens fixed to the sclera was performed and the patient regained 2/10 visual acuity on the right eye and remained at counting fingers in the left.

**Conclusion:** Marfan syndrome is a rare condition, with various ocular manifestations. The only guarantee for a better visual outcome is a well-designed therapeutic approach.

\*Corresponding author: E-mail: [dr.filali.mehdi@gmail.com](mailto:dr.filali.mehdi@gmail.com);

**Keywords:** *Bilateral ectopia lentis; marfan syndrome; tissue disorder.*

## 1. INTRODUCTION

Marfan syndrome is an autosomal dominant genetic connective tissue disorder resulting from FBN1 (15q21.1) gene mutation. It typically affects the cardiovascular, skeletal and ocular systems but can also involve other systems such as the central nervous system, skin and the respiratory system [1].

Ocular manifestations of Marfan syndrome are numerous, crystalline lens dislocation or ectopia lentis constitutes the most frequent one, and has been reported to happen in up to the 70% cases [2]. It can cause symptoms like blurred vision, or monocular diplopia. Its management is still, to this day very challenging and depends on the severity of the symptoms, patients' age and the degree of dislocation, among other parameters.

## 2. CASE REPORT

We describe the case of a 51-year-old female patient, with a family history of Marfan syndrome; who presented to our department with progressive decreased visual acuity for more

than twenty years. General examination revealed a patient of tall stature (177 cm), tall narrow face shaped, and long slender fingers (Fig.1).


Ophthalmological examination revealed a distance visual acuity limited to counting fingers in both eyes, the automated refraction was not obtainable. Ocular pressure was within normal range. Examination of the anterior segment showed clear corneas, iridodonesis, and ectopic opaque crystalline lenses bilaterally. Fundus revealed multiple drusens scattered throughout the retina, reduced macular reflex, and chorioretinal atrophy (Fig. 2).

Optical coherence tomography (OCT) showed bilateral diffuse drusens and remodelling of the pigment epithelial layer associated to choroidal thickening (Fig. 3).


Given the family history, the general physical features of the patient and ocular findings, diagnostic of Marfan syndrome was established. The patient was referred to the internal medicine department for further assessment of other possible systemic abnormalities.



**Fig. 1. Image showing the the main clinical signs that may point to Marfan syndrome : A suggestive morphotype (large size with a lanky morphotype) and arachnodactyly**



**Fig. 2.** Image resuming the biomicroscopique exam. 2a/ image showing fundus : multiple diffuse drusens scattered throughout the retina. 2b/ image showing bilateral iris atrophy and subluxated ectopic lens



**Fig. 3.** OCT image showing remodeling of the pigment epithelial layer and choroidal thickening. 3a/ right eye. 3b/ left eye.

On the other hand, the management of the ectopic lentis was very challenging due to the zonular weakness, the lens instability, and the possibility of a mediocre visual acuity recovery because of undiagnosed amblyopia. After considering the various therapeutic options, we opted for intracapsular extraction of the lens. The operation was performed under local anesthetic, using a wide keratotomy followed by lensectomy after endothelial protection with viscoelastic. Next, a 25-gauge anterior vitrectomy was performed and all vitreous prolapses were

excised. Implantation was performed after viscoelastic protection of the corneal endothelium and injection of pilocarpine for the myotic effect. The intraocular implant clipped to the iris (ICIOL) was inserted in a concave-convex position, with the IOL turned towards a horizontal plane and centred on the pupil. The inverted ICIOL optic was held in place with forceps. A haptic was gently slid behind the iris; the optic was lifted slightly forward over the pupil. The optic was lifted slightly forward over the posterior surface of the iris so that the haptic could be recognised on

the anterior surface of the iris. A long microspatula was used to insert the iris tissue into the claw. The second haptic was fixed in the same way. Only a small amount of iris tissue was embedded in the claw haptic to avoid pupil formation. Post-operative visual acuity at 3 months improved to 2/10 on the right eye and remained at counting fingers in the left. We followed up closely the patient, after 1 year no complication has occurred.

### 3. DISCUSSION

Ectopia lentis is observed in 50% to 80% of Marfan syndrome cases, frequently presenting as the primary clinical manifestation [3].

The primary approach in managing lens subluxation involves correcting refractive errors with eyeglasses. However, if the subluxation is significant, characterized by the lens intersecting the pupil, posterior dislocation into the vitreous cavity, or anterior dislocation of the lens with or without secondary glaucoma, solely relying on eyeglasses is unlikely to restore visual function. In such cases, extraction of the crystalline lens becomes necessary [4].

Historically, due to the elevated occurrence of intra- and postoperative complications and unsatisfactory visual results, surgery was not widely favored. However, with advancements in microsurgical instruments and surgical techniques, there has been a resurgence of interest in surgical intervention. Several surgical strategies for managing ectopia lentis in Marfan syndrome have been described including standard lensectomy with or without anterior vitrectomy, or lens extraction using capsular tension rings when mild zonular desinsertion is present [5]. Post-operative refractive rehabilitation may consist of the utilization of aphakic glasses, contact lenses, or intra-ocular lenses scleral-fixated, iris-fixated in the anterior or posterior chamber [6].

Following surgery, patients should undergo regular monitoring for IOL alignment, retinal degeneration and detachment that are reported to be present in 18% of eyes in Marfan syndrome; this incidence is higher up to 70% in patients with a subluxated lens [7] Moreover, patients should be encouraged to promptly seek ophthalmological assessment if they notice flashes, floaters, or encounter any partial or complete vision loss [8]. For our patient, none of

these symptoms occurred after one year of follow-up.

### 4. CONCLUSION

Marfan syndrome is a rare condition, with various ocular manifestations and complications. Therefore, Ophthalmologists play a substantial role in making a prompt clinical diagnosis and a well thought out therapeutic approach, sole guarantee of a better visual outcome and a significantly improved life quality.

### CONSENT

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

### ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Akram H, Aragon-Martin JA, Chandra A. Marfan syndrome and the eye clinic: From diagnosis to management. *Ther Adv Rare Dis.* 2021;2:26330040211055738. DOI:10.1177/26330040211055738
2. Faria MY, Ferreira N, Neto E. Retropupillary iris-claw intraocular lens in ectopia lentis in Marfan syndrome. *Int Med Case Rep J.* 2016;9:149–53.
3. Shah S, Shah M, Chandane P, Makhloga S, Thorat D, Sanghani M. Clinical profile and outcome of ocular manifestation in Marfan syndrome in India. *Indian J Ophthalmol.* 2022;70:626–629. DOI:10.4103/ijo.IJO\_1651\_21
4. Esfandiari H, Ansari S, Mohammad-Rabei H, Mets MB. Management Strategies of Ocular Abnormalities in Patients with Marfan Syndrome: Current Perspective. *J Ophthalmic Vis Res.* 2019; 14(1):71–77. DOI:10.4103/jovr.jovr\_29\_18
5. Al-Dwairi RA, Shihadeh WA, Alqudah AA, Alqudah NM, Msallam M, Alrazem F, et al. Outcomes of Iris-Claw IOL Implantation in

- Patients with Marfan's Syndrome in Jordan. Clin Ophthalmol. 2022;16:3811–9.
6. V Miraldi Utz, Coussa RG, Traboulsi EI. Surgical management of lens subluxation in Marfan syndrome. Journal of American Association for Pediatric Ophthalmology and Strabismus. 2014;18(2):140–146. DOI: 10.1016/j.jaapos.2013.12.007.
  7. Dogăroiu AC, Dogăroiu C, Tătaru CP. Ectopia lentis surgery in Marfan's syndrome: From “couching” to the use of intracapsular tension rings. Rom J Ophthalmol. 2022;66(1):2–7.
  8. Loeys BL, Dietz HC, Braverman AC, Callewaert BL, De Backer J, Devereux RB, et al. The revised Ghent nosology for the Marfan syndrome. Journal of Medical Genetics. 2010;47(7):476–85.

---

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/115465>