



Effect of Nitrogen and Silica on Growth and Yield of Paddy (*Oryza sativa* L.) Variety CO 51

G. Poovizhi Sindhu^{a*}, D. Jawahar^a, T. Chitdeshwari^a, P. Jeyakumar^b and D. Jeya Sundara Sharmila^c

^a Department of Soil Science and Agricultural Chemistry, TNAU, Coimbatore, India.

^b Department of Crop Physiology, TNAU, Coimbatore, India.

^c Department of Nano Science and Technology, TNAU, Coimbatore, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i130821

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/82318>

Original Research Article

Received 12 January 2022

Accepted 25 January 2022

Published 25 January 2022

ABSTRACT

World agriculture is facing increasing demand due to multiple challenges. The biggest challenge is to attain food security in the ever increasing growing population. Rice (*Oryza sativa* L.) is a staple food crop for half of the world's population and one of the world's most significant food crops. Farmers must produce more rice with enhanced quality in order to ensure food security in rice-consuming countries across the world. Rice (*Oryza sativa* L.) is considered to be a Si accumulator plant and tends to actively accumulate Si into tissue up to concentrations of 5% or even higher. The present study aimed to use different forms of N and Si sources at different levels to improve the growth traits contributing to the increase yield of rice. A pot experiment was conducted in glasshouse, TNAU, Coimbatore to assess the response of nitrogen and silica on paddy during October 2020 – January, 2021. The treatments comprised of the two factors, nitrogen and silica with three replications. The growth parameters like plant height, number of tillers per plant and total dry matter production were recorded at all stages and yield traits at harvest stage. The treatment, T₁₁ - N₂S₂ (53.4 and 110.2 g pot⁻¹ respectively) significantly enhanced the grain yield (34%) and straw yield (40%) than control in the pot experiment. Highest growth and yield parameters were recorded with application of combined fertilization of nitrogen and silica (T₁₁-N₂S₂) which comprises 150 kg ha⁻¹ of urea and 200 kg ha⁻¹ of calcium silicate compared to control.

Keywords: Urea; calcium silicate; growth parameters and yield attributes.

1. INTRODUCTION

Rice is the major staple food for human population of southeastern Asia and occupies a predominant position among the cereal crops. Rice grain production in India is about 118.87 and 121.46 million tons during the year 2019-2020 and 2020-2021 respectively [1]. Increased population in turn increased the global rice demand and the only way to meet the growing demand is through increased the production of cereals. Nitrogen (N) and silicon (Si) are two important nutritional elements required for plant growth and development in rice. Nitrogen (N) is the important and primary element. It is one of the key inputs to paddy fields along with phosphorus to improve initial vigor of rice plants. Healthy plants contain 3 to 4 percent nitrogen in the aerial biomass. It is a major component of photosynthesis; enhance the chlorophyll content, amino acids and serves as building blocks of protein. Silicon (Si) is a beneficial element for paddy growth and development [2,3]. Rice is a typical silicon accumulator that benefited through exogenous silicon nutrition. Silicon supply is essential for healthy growth of plant which results in improved economic yield of rice crop [4]. Hence, the application of silicon to soil or crop is practically applicable and also it increases the crop yield and alleviates the Fe toxicity [5]. Lavinsky et al. [6] reported that silicon application at reproductive growth stage (panicle initiation to heading) rather than the vegetative and ripening stages had a positive effect on photosynthesis with increased stomatal conductance. Along with improvement in photosynthetic traits, exogeneous application of Si provides biotic and abiotic tolerance and has improved plant metabolism [7].

Experimental variety is Rice variety CO51. It was released by Tamil Nadu Agricultural University in the year 2013. The parentage of this variety is ADT43 / RR272-1745. It is a high yielding semi dwarf rice variety with 105-110 days duration. It has been recommended for cultivation as transplanted rice throughout Tamil Nadu except Nilgiris district. Average yield of the variety is 6641 kg ha⁻¹. Si concentration is much higher than other nutrients such as nitrogen (N), phosphorus (P), and potassium (K) in rice plants when plants are supplied with exogenous silica [8]. Since, Si is a beneficial element, rice crop deficient in silica considerably reduces the growth and fertility which in turn declines yield

[3]. Application of N and Si fertilizers significantly increases the plant height and number of panicles than control [9]. When nitrogen and silica is used in combination, the ideal level of both nitrogen and silica has been determined to be raised from its current level which is responsible for improved crop growth and high productivity [10]. With this background, the present study was formulated to study the interactive effect of nitrogen and silica on growth parameters and yield traits in paddy variety CO 51.

2. MATERIALS AND METHODS

A pot experiment was conducted in glasshouse, TNAU, Coimbatore with treatment were N₀ – without urea, N₁– 100 kg ha⁻¹ urea, N₂–150 kg ha⁻¹ urea, N₃ – 175 kg ha⁻¹ urea, S₀–0 kg of Calcium silicate, S₁– 150 kg of Calcium silicate, S₂–200 kg of Calcium silicate, S₃–250 kg of Calcium silicate. The experiment was conducted in factorial completely randomized block design and replicated thrice. The experimental soil belongs to Noyyal soil series taxonomically grouped as Clay loamy, mixed iso-hyperthermic, *Typic haplustalf*. The composite soil samples were collected initially from the experimental field and had been subjected to analysis initial characteristics. The soil was clay loam in texture, pH - 7.4, non-saline (EC - 0.60 dSm⁻¹), bulk density - 1.26 Mg m⁻³, particle density - 2.53 Mg m⁻³ and total porosity - 50%. The initial soil fertility status revealed that the experimental soil has low available N (265 kg ha⁻¹), medium available P (20 kg ha⁻¹) and high available K (575 kg ha⁻¹) status. Fertilizers were applied basal, 20th and 45th DAT. Rice variety CO51 was released by Tamil Nadu Agricultural University in the year 2013. The parentage of this variety is ADT43 / RR272-1745. It is a high yielding semi dwarf rice variety with 105-110 days duration. Variety CO51 is moderately resistant to blast, brown plant hopper. It is white medium slender rice with high milling (69%) and head rice recovery (63%) percentage. The analytical methods of soil analysis were furnished below the Table 1.

2.1 Data Collection

The growth parameters were (plant height, number of tillers per plant and total dry matter production) measured at 30, 60 and harvest

stage and yield attributes (grain yield, straw yield and harvest index at harvest stage.

The plant height was measured from the base of the leaf to tip of the longest leaf stretched and the mean value was expressed in centimeter.

Tiller population was counted from plant and is denoted as number of tillers m^{-2} at 30, 60 and harvest stage.

Samples were collected from pot and shade dried and then oven dried at $60^{\circ}C$ for 72 hrs. The dry weight of the samples was used for the estimation of dry matter production and it is expressed in $g\ pot^{-1}$ at 30, 60 and harvest stage.

The harvested grains from the pot was threshed, cleaned, sun dried and weighed and are symbolized in terms of $g\ pot^{-1}$.

The paddy straw collected from the pot was sun dried, weighed and is represented as $g\ pot^{-1}$.

Harvest index is the ratio of harvested grain yield to total shoot dry matter and this can be used as a measure of reproductive efficiency.

Harvest index (%) = Economic yield /Biological yield $\times 100$

The results of growth and yield parameters were statistically analyzed as suggested by [11].

3. RESULTS AND DISCUSSION

Plant height is an important morphological trait which determines the overall plant frame work.

The treatment $T_{11} - N_2S_2$ significantly recorded higher plant height (35.4, 79.8, and 84.9 cm respectively), followed by the treatment $T_8 - N_1S_3$ (31.8, 75.6 and 80.1 cm respectively) and the treatments $T_1 - N_0S_0$ and $T_{16} - N_3S_3$ were recorded significantly lower plant height (11.6, 42 and 46.8 cm) and (10.3, 41.5 and 45.2 cm) respectively (Table 2) at 30, 60 and harvest stage. Application of nutrients helps in cell elongation and increased plant height. Similar findings were reported by Mazid et al., [11] and Arf [12] that the application of nitrogen at increasing level boosts the plant metabolism and significantly increased plant height. Present study were supported by Chu et al. [12] and Yoshida et al. [13] that the silica fertilization improves the plant rigidity and leaf erectness in high density population.

The application of nitrogen and silicon fertilizer greatly enhanced the number of tillers per hill. Increased number of tillers per plant were recorded in the treatment, $T_{11} - N_2S_2$ (12.4, 22.1 and 28.9 respectively) followed by the treatment, $T_8 - N_1S_3$ (11.8, 20.8 and 25.4 respectively). Least number of tillers per plant were recorded in the treatments, $T_1 - N_0S_0$ (6.2, 7.1 and 8.8 respectively) and the treatment $T_{16} - N_3S_3$ (6.1, 7.0 and 8.1 respectively) at 30, 60 and harvest stage (Table 3). Fertilization of rice crop with silicon and nitrogen positively increases plant vigor and increases tiller number per hill. Similar findings were reported by Mazid et al. [14]; Pati et al. [15]; Dorairaj et al. [16]; and Cuong et al. 17] that the number of tillers per hill were increased due Si silica fertilization at top dressing and reproductive stage.

Table 1. Analytical methods of soil analysis

Sl. No	Parameters	Methods	References
1.	pH	pH meter (soil water ratio 1:2.5)	Jackson (1973)
2.	Electrical conductivity	Conductivity meter (soil water ratio 1:2.5)	Jackson (1973)
3.	Bulk density	Measuring Cylinder Method	Sree ramulu (2003)
4.	Particle density	Measuring Cylinder Method	Sree ramulu (2003)
5.	Porosity	Measuring Cylinder Method	Sree ramulu (2003)
6.	Available nitrogen	Alkaline potassium permanganate method	Subbiah and Asija (1956)
7.	Available phosphorous	Bray No 1 extraction and photoelectric colorimetry	Jackson (1973)
8.	Available potassium	Neutral normal ammonium acetate extraction and flame photometry	Stanford and English (1949)

Table 2. Effect of Nitrogen and silica on plant height in paddy at 30, 60 DAT and harvest stage of CO 51 (cm)

Treatment	30 DAT					60 DAT					Harvest				
	S ₀	S ₁	S ₂	S ₃	Mean	S ₀	S ₁	S ₂	S ₃	Mean	S ₀	S ₁	S ₂	S ₃	Mean
N ₀	11.6	15.4	16.9	17.3	15.3	42.1	47.4	48.5	49.7	46.9	46.8	52.8	53.1	54.6	51.8
N ₁	18.2	19.8	25.8	31.8	23.9	50.3	51.2	65.9	75.6	60.7	55.2	56.9	67.9	80.1	65.0
N ₂	23.4	24.5	35.4	26.1	27.3	55.4	59.8	79.8	71.4	66.6	60.1	62.5	84.9	74.8	70.5
N ₃	20.1	21.9	22.4	10.3	18.6	52.1	53.7	54.8	41.5	50.5	57.2	58.4	59.8	45.2	55.1
Mean	18.3	20.4	25.1	21.3		49.9	53.0	62.2	59.5		54.8	57.6	66.4	63.6	
Factor	S	N	S*N			S	N	S*N			S	N	S*N		
SEd	0.6	0.8	1.1			1.5	2.1	2.9			1.6	2.2	3.2		
CD	1.2	1.7	2.4			3.1	4.4	6.3			3.3	4.8	6.9		

N₀ –without urea
N₁– 100 kg ha⁻¹ urea

N₂–150 kg ha⁻¹ urea
N₃– 175 kg ha⁻¹ urea

S₀–0 kg ha⁻¹ of Calcium silicate
S₁– 150 kg ha⁻¹ of Calcium silicate

S₂–200 kg ha⁻¹ of Calcium silicate
S₃–250 kg ha⁻¹ of Calcium silicate

Table 3. Effect of Nitrogen and silica on number of tillers in paddy at 30, 60 DAT and harvest stage of CO 51

Treatment	30 DAT					60 DAT					Harvest				
	S ₀	S ₁	S ₂	S ₃	Mean	S ₀	S ₁	S ₂	S ₃	Mean	S ₀	S ₁	S ₂	S ₃	Mean
N ₀	6.2	6.9	7.5	7.9	7.1	7.1	7.8	8.2	8.7	7.9	8.8	9.4	10.5	11.3	10
N ₁	8.1	8.4	10.6	11.8	9.7	9.5	10.8	16.5	20.8	14.4	12.4	13.9	19.8	25.4	17.8
N ₂	9.8	10.1	12.4	11.2	10.8	14.3	15.8	22.1	18.2	17.6	17.9	18.5	28.9	22.1	21.8
N ₃	9	9.3	9.6	6.1	8.5	11.2	12.7	13.8	7.0	11.1	14.2	15.7	16.5	8.1	13.6
Mean	8.2	8.6	10.0	9.2		10.5	11.7	15.1	13.6		13.3	14.3	18.9	16.7	
Factor	S	N	S*N			S	N	S*N			S	N	S*N		
SEd	0.2	0.3	0.5			0.3	0.5	0.7			0.4	0.6	0.8		
CD	0.5	0.7	1.1			0.7	1.0	1.4			0.9	1.2	1.7		

N₀ –without urea
N₁– 100 kg ha⁻¹ urea

N₂–150 kg ha⁻¹ urea
N₃– 175 kg ha⁻¹ urea

S₀–0 kg ha⁻¹ of Calcium silicate
S₁– 150 kg ha⁻¹ of Calcium silicate

S₂–200 kg ha⁻¹ of Calcium silicate
S₃–250 kg ha⁻¹ of Calcium silicate

Plant height, number of tillers was positively correlated with total dry matter production and the data was furnished in the Table 4. The treatment T_{11} - N_2S_2 (35.4, 43.6 and 55.8 g pot⁻¹ respectively) was recorded highest total dry matter production followed by the treatment T_8 - N_1S_3 (31.2, 39.4 and 51.2 g pot⁻¹ respectively). Lowest total dry matter production were recorded in the treatment, T_1 - N_0S_0 (10.3, 18.1 and 27.9 g pot⁻¹ respectively) and the treatment, T_{16} - N_3S_3 (9.8, 17.4 and 24.6 g pot⁻¹ respectively) at 30, 60 DAT and harvest. The present findings were supported by Yogendra et al. [18] that increase in number of tillers increases the total dry matter production with the application of calcium silicate at 2 t ha⁻¹ and Mazid et al. [14] observed that increased plant height and number of tillers per plant substantially increased the total dry matter production. The study suggested by Agarie et al., [19] found that the application of Si significantly enhanced the dry matter production which is mainly could be maintenance of photosynthetic activity in the leaves.

Number of productive tillers and total dry matter production positively enhanced the rice grain yield and straw yield. The results revealed that the grain yield and straw yield were recorded higher in the nutrient treatments. The treatment

T_{11} - N_2S_2 (53.4 and 110.2 g pot⁻¹ respectively) recorded maximum grain and straw yield followed by the treatment, T_8 - N_1S_3 (48.9 and 101.4 g pot⁻¹ respectively). The treatment, T_1 - N_0S_0 (19.5 and 46.8 g pot⁻¹ respectively) and the treatment T_{16} - N_3S_3 (18.2 and 44.2 g pot⁻¹ respectively) recorded the lowest grain yield and straw yield (Fig. 1 & 2). The present findings was in accordance with Chen et al., [20] that the silica transferred preferentially into the flag leaves and grains during the reproductive stage which increased the grain yield. The research findings found by Sudhakara et al., [21]; Savant et al., [2]; Ghanbari-Malidareh [22]; Balastra et al., [23]; and Deren et al., [24] reported that the high rates of silicon fertilization was substantially enhanced grain mass, straw yield leaf culm strengthens because, majority of the silica absorbed by the plant is precipitated and deposited inside or between plant cells and forming 'phytoliths', which offer mechanical strength and help protect the plant from physical, chemical, and biological stressors. Phytoliths are found in a wide range of monocot and dicot plants. The nutrient treatments positively influenced the harvest index but however, it was not significant in the interaction of nitrogen and silica treatments in the experiment (Fig. 3) [25].




Fig. 1. Effect of Nitrogen and silica on grain yield in paddy of CO51 (g pot⁻¹)

T_1 - N_0S_0
 T_2 - N_0S_1
 T_3 - N_0S_2
 T_4 - N_0S_3

T_5 - N_1S_0
 T_6 - N_1S_1
 T_7 - N_1S_2
 T_8 - N_1S_3

T_9 - N_2S_0
 T_{10} - N_2S_1
 T_{11} - N_2S_2
 T_{12} - N_2S_3

T_{13} - N_3S_0
 T_{14} - N_3S_1
 T_{15} - N_3S_2
 T_{16} - N_3S_3

Table 4. Effect of Nitrogen and silica on total dry matter production in paddy at 30, 60 DAT and harvest stage of CO51 (g pot⁻¹)

Treatment	30 DAT					60 DAT					Harvest				
	S ₀	S ₁	S ₂	S ₃	Mean	S ₀	S ₁	S ₂	S ₃	Mean	S ₀	S ₁	S ₂	S ₃	Mean
N ₀	10.3	13.6	15.1	16.2	13.8	18.1	21.2	23.8	24.9	22	27.9	31.8	33.6	34.1	31.8
N ₁	17.8	18.7	24.6	31.2	23.0	25.1	26.3	32.9	39.4	30.9	35.9	36.4	42.1	51.2	41.4
N ₂	18.7	24.6	34.5	22.1	24.1	30.7	31.8	43.6	35.1	35.3	40.6	41.1	55.8	46.7	46.0
N ₃	19.5	20.9	21.6	9.8	17.9	27.4	28.2	29.8	17.4	25.7	37.2	38.4	39.6	24.6	34.9
Mean	16.5	19.4	23.9	19.8		25.3	26.8	32.5	29.2		35.4	36.9	42.7	39.1	
Factor	S	N	S*N			S	N	S*N			S	N	S*N		
SEd	0.5	0.7	1.1			0.7	1.0	1.5			0.9	1.4	2.0		
CD	1.0	1.6	2.2			1.6	2.2	3.1			2.1	3.0	4.3		

N₀ –without urea
N₁– 100 kg ha⁻¹ urea

N₂– 150 kg ha⁻¹ urea
N₃– 175 kg ha⁻¹ urea

S₀–0 kg ha⁻¹ of Calcium silicate
S₁– 150 ha⁻¹ kg of Calcium silicate

S₂–200 kg ha⁻¹ of Calcium silicate
S₃–250 ha⁻¹ kg of Calcium silicate




Fig. 2. Effect of Nitrogen and silica on straw yield in paddy of CO51 (g pot⁻¹)

T₁– N₀S₀
 T₂– N₀S₁
 T₃– N₀S₂
 T₄– N₀S₃

T₅– N₁S₀
 T₆– N₁S₁
 T₇– N₁S₂
 T₈– N₁S₃

T₉– N₂S₀
 T₁₀– N₂S₁
 T₁₁– N₂S₂
 T₁₂– N₂S₃

T₁₃– N₃S₀
 T₁₄– N₃S₁
 T₁₅– N₃S₂
 T₁₆– N₃S₃




Fig. 3. Effect of Nitrogen and silica on harvest index in paddy of CO51 (g pot⁻¹)

T₁– N₀S₀
 T₂– N₀S₁
 T₃– N₀S₂
 T₄– N₀S₃

T₅– N₁S₀
 T₆– N₁S₁
 T₇– N₁S₂
 T₈– N₁S₃

T₉– N₂S₀
 T₁₀– N₂S₁
 T₁₁– N₂S₂
 T₁₂– N₂S₃

T₁₃– N₃S₀
 T₁₄– N₃S₁
 T₁₅– N₃S₂
 T₁₆– N₃S₃

The correlation coefficient were established between grain yield and growth parameters. The effect of plant height (r =0.98**), number of tillers per plant (r =0.98**), total drymatter

production (r =0.99**) were significant and positive in influencing the grain yield of paddy in pot experiment (Table 5. and Fig.4.)

Table 5. Correlation between yield and growth parameters

	Grain yield per pot	Plant height	No of tillers per plant	DMP per plant
Grain yield	1			
Plant height	0.991087	1		
No of tillers per plant	0.989314	0.979455	1	
DMP per plant	0.99525	0.981821	0.982373	1




Fig. 4. Regression between yield and growth parameters

4. CONCLUSION

The results revealed that the application of fertilizer like Nitrogen and Silica in combinations significantly improved the growth parameters and yield traits. Increased silicon levels resulted in a considerable increase in straw yield. This could be owing to silicon's function in improving photosynthetic activity, water use efficiency, and nutrient use efficiency. As a result, there is better vegetative growth. The increased plant height and number of tillers per hill were primarily responsible for the enhanced straw yield. Higher plant height, number of tillers per plant and total dry matter production were recorded in the treatment N₂S₂ (150 kg ha⁻¹ of urea and 200 kg ha⁻¹ of calcium silicate). which also boosted yield and yield attributes in this study. Silica fertilization positively correlated with plant height, number of tillers and total dry matter production which in turn significantly increased the yield. The production of paddy with application of

combined fertilizer is proven to be beneficial to the paddy farmers.

ACKNOWLEDGEMENT

The authors are sincerely thankful to the concerned departments of TNAU for help and support during the research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ministry of Agriculture and Farmers Welfare releases Third Advance Estimates of Principal Crops for; 2020-21. Available: <https://pib.gov.in/PressReleaselframePage.aspx?PRID=1721692>.

2. Savant NK, Snyder GH, Datnoff LE. Silicon management and sustainable rice production. *Advances in Agronomy*. 1996; 58:151-199.
3. Ma JF, Yamaji N. Functions and Transport of Silicon in Plants. *Cellular and Molecular Life Science*. 2008;65:3049-3057.
4. Singh V, Singh V, Singh S, Khanna R. Effect of zinc and silicon on growth and yield of aromatic rice (*Oryza sativa*) in North-Western plains of India. *J. Rice Res. Dev.* 2020;3(1):82-86.
5. Das A, Samant PK, Sahu G, Santra GH. Evaluation of silicon nutrition on yield, growth attributes and available nutrient status of paddy in coastal zone of Odisha. *Int. J. Pl. & Soil Sci.* 2021;33(23):277-288.
6. Lavinsky AO, Detmann KC, Reis JV, Ávila RT, Sanglard ML, Pereira LF, DaMatta FM. Silicon improves rice grain yield and photosynthesis specifically when supplied during the reproductive growth stage. *Journal of Plant Physiology*. 2016; 206:125-132.
7. Tripathi DK, Vishwakarma K, Singh VP, Prakash V, Sharma S, Muneer S, Corpas FJ. Silicon crosstalk with reactive oxygen species, phytohormones and other signaling molecules. *Journal of Hazardous Materials*. 2021;408:124820.
8. Ma JF, Takahashi E. *Soil, Fertilizer and Plant Silicon Research in Japan*. Elsevier. Amsterdam; 2002.
9. Malav JK, Patel KC, Sajid M, Ramani VP. Effect of silicon levels on growth, yield attributes and yield of rice in typical ustochrepts soils. *Eco. Env. & Cons.* 2005;205-208.
10. Singh KK, Singh K, Singh R, Singh Y, Singh CS. Response of nitrogen and silicon levels on growth, yield and nutrient uptake of rice (*Oryza sativa* L.). *Oryza*. 2006;43(3):220.
11. Gomez KA, Gomez AA. *Statistical Procedures for Agricultural Research*. John Wiley & Sons; 1984.
12. Arf O, Rodrigues RAF, Sá MED, Crusciol CA. CResposta de cultivares de arroz de sequeiro ao preparo do solo e à irrigação por aspersão. *Pesquisa Agropecuária Brasileira*. 2001;3:871-879.
13. Chu M, Liu M, Ding Y, Wang S, Liu Z, Tang S, Li G. Effect of nitrogen and silicon on rice submerged at tillering stage. *Agronomy Journal*. 2018;110(1): 183-192.
14. Mazid A. Impact of modern agricultural technologies on durum wheat production in Syria. Aleppo, Syria: International Center for Agricultural Research in the Dry Areas (ICARDA); 2003.
15. Pati S, Pal B, Badole S, Hazra GC, Mandal B. Effect of Silicon Fertilization on Growth, Yield and Nutrient Uptake of Rice. *Communications in Soil Science and Plant Analysis*. 2001;47:284-290.
16. Dorairaj D, Ismail MR, Sinniah UR, Tan KB. Silicon mediated improvement in agronomic traits, physiological parameters and fiber content in *Oryza sativa*. *Acta Physiologiae Plantarum*. 2020;42(3):1-11.
17. Cuong TX, Ullah H, Datta A, Hanh. Effects of silicon-based fertilizer on growth, yield and nutrient uptake of rice in tropical zone of Vietnam. *Rice Science*. 2001;24(5):283-290.
18. Yogendra ND, Kumara BH, Chandrashekar N, Prakash NB, Anantha MS, Jeyadeva HM. Effect of silicon on real time nitrogen management in a rice ecosystem. *African Journal of Agricultural Research*. 2014;9(9):831-840.
19. Agarie S, Agata W, Kubota F, Kaufman, PB. Physiological roles of silicon in photosynthesis and dry matter production in rice plants: I. Effects of silicon and shading treatments. *Japanese Journal of Crop Science*, 1991;61(2):200-206.
20. Chen JX, Tu NM, Yi ZX, Zhu HL. Effect of silicon fertilizer on nitrogen utilization efficiency of super early rice. *Crop Research*. 2011;01.
21. Sudhakara N, Anjaiah T, Qureshi AA, Sagar, GCV, Chary DS. Effect of different sources and levels of Silicon application on growth, yield attributes and yield of rice (*Oryza sativa* L.). *The J. Res. Pjtsau*. 2020;48(55):1-99.
22. Ghanbari-Malidareh A. Silicon application and nitrogen on yield and yield components in rice (*Oryza sativa* L.) in two irrigation systems. *International Journal of Agricultural and Biosystems Engineering*. 2020;5(2):40-47.
23. Barbosa Filho MP. *Nutrição e adubação do arroz*. Piracicaba: Associação Brasileira para Pesquisa da Potassa e do Fosfato, 127.

24. Deren CW, Datnoff, LE, Snyder, GH, Martin, FG. Silicon concentration, disease response, and yield components of rice genotypes grown on flooded organic Histosols. Crop Science.1994;34:73.
25. Yoshida S, Naveser SA, Ramirez EA. Effects of silica and nitrogen supply on some leaf characters of rice plant. Plant and Soil.1969;31:48-56.

© 2022 Sindhu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/82318>