



# Identification and Molecular Characterization of Bacteria and Fungi Associated with Three Fresh Edible Mushrooms

Caroline Iruoma Anyakorah <sup>a\*</sup> and Adeniji Omolara Folakemi <sup>a</sup>

<sup>a</sup> Department of Biological Sciences Bells University of Technology Ota Ogun State Nigeria.

## Authors' contributions

This work was carried out in collaboration between both authors. Author CIA designed the study, wrote the protocol, and corrected the draft of the manuscript. Author AOF managed the analyses of the study and wrote the draft of the manuscript. Both authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/JALSI/2022/v25i6582

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92636>

Original Research Article

Received 07 August 2022  
Accepted 13 October 2022  
Published 26 October 2022

## ABSTRACT

**Aims:** Mushrooms are health-food used as food supplements and fortification but are susceptible to microbial spoilage due to high moisture content and nutrient status. Fresh mushrooms sometimes show signs of spoilage after harvest and makes storage difficult. The objective of the study was to identify and characterize bacteria and fungi associated with three fresh edible mushrooms, under ambient and cold temperatures.

**Methodology:** The study was conducted at the Bells University of Technology, Nigeria, between December 2020 and October 2021. *Pleurotus ostreatus* and *Calocybe indica* fruitbodies were procured from the Federal Institute of Industrial Research Oshodi while *Pleurotus tuber-regium* fruitbodies were obtained from sclerotia planted in loamy soil. The fruitbodies were kept at ambient (28°C) and cold (15°C) temperatures respectively. The bacterial and fungal counts on each of the mushrooms were taken at 0,3, 5, and 7 days after harvest. The isolated bacteria were identified by conventional methods; Analytical Profile Index (API) 20E kits (BioMerieux), while Fungi were identified by morphological features and PCR amplification using ITS 1f/ ITS 4r universal primers.

**Results:** The bacterial and fungal counts on the fruitbodies ranged from 5.7 log CFU/ml – 6.3 log CFU/ml and 5.0 log CFU/ml – 5.9 log CFU/ml respectively. Seven genera of bacteria isolated were gram-negative bacteria. At ambient temperature, *Pseudomonas aeruginosa*, *Enterobacter asburiae*, *Klebsiella oxytoca*, *Klebsiella ornithinolytica*, *Serratia marcescens*, *Chryseobacterium*

\*Corresponding author: E-mail: Anyacaro2003@yahoo.com;

*meningosepticum* and *Cedecea davisae* were isolated while *Enterobacter cloacae*, *Enterobacter sakazakii* and *Citrobacter braakii* were isolated at cold temperature. *Aspergillus*, *Penicillium*, and *Fusarium* were isolated at both temperatures while *Alternaria* was isolated at ambient temperature.

**Conclusion:** Isolated bacteria and fungi were mostly enteric pathogens and potential mycotoxin-producing fungi. This is an indication that strict hygiene and control measures should be put in place during the production and storage of these mushrooms in order to improve the quality and food safety of fresh mushrooms in Nigeria.

**Keywords:** Bacteria; fungi; identification; microbial counts; mushroom; storage temperature.

## 1. INTRODUCTION

Mushrooms are edible fungi with abundant nutritional and health benefits considered to be essential for humans [1,2]. They are rich in protein, vitamins, fiber, minerals, and bioactive substances and are regarded as health-food. In some parts of Nigeria, the locals use mushrooms to replace meat in their diet such that mushrooms are gaining considerable attention, and youths are being empowered to acquire skills in this regard.

*Calocybe indica*, *Pleurotus ostreatus*, and *Pleurotus tuber-regium* are tropical edible mushrooms grown commercially in Nigeria and provide high returns for small-scale farmers. Due to their nutrient content and high moisture content, mushrooms are predisposed to different microorganisms and could harbour pathogenic microorganisms depending on cultural and environmental conditions of production and storage. Microbial infestation can occur during production, harvest, postharvest handling, storage, transportation, and processing. *Agaricus bisporus* mushrooms of good quality immediately after harvest developed brown blotches at retail, even while kept at refrigeration temperatures [3]. Temperature is one of the most measured factors accountable for mushroom deterioration and storage temperature is an important factor that affects the postharvest life of mushrooms. Higher storage temperature can activate tyrosinase and increase microbial growth [1]. The study revealed that different bacteria and fungi were associated with the surface of fresh mushrooms at both ambient and refrigeration temperatures. Microbial growth is a function of a variety of factors, both intrinsic (nutrient content, pH, water activity, physical structure of food oxidation-reduction potential) and extrinsic (temperature, relative humidity and food microbiota) [4]. Mushrooms are known to possess the above attributes, so fresh mushrooms can provide an ideal growth substrate for microorganisms. Therefore, the

identification of the microorganisms associated with fresh mushrooms is essential with respect to food safety, quality control, enhanced productivity, and marketability. The study was to identify and characterize microorganisms associated with fresh *Calocybe indica*, *Pleurotus ostreatus*, and *Pleurotus tuber-regium* under ambient and refrigeration temperatures.

## 2. MATERIALS AND METHODS

### 2.1 Materials

*Calocybe indica* and *Pleurotus ostreatus* were procured from the Federal Institute of Industrial Research Oshodi, Lagos, Nigeria; while the sclerotia of *Pleurotus tuber-regium* purchased from a local market (Ojuore market in Ota, Ogun State), were planted in the soil, watered until fruitbodies were produced and harvested. The study was conducted between December 2020 and October 2021. The fruit bodies were kept at ambient (28°C) and cold temperature (15°C) respectively.

### 2.2 Microbiological Examination of Mushroom Samples

Bacterial and fungal counts were carried out, according to the method of [5] using potato dextrose agar and nutrient agar. The procedure was repeated on days 3, 5, and 7 respectively. Incubation of bacterial and fungal plates was at 37°C and 28°C respectively.

### 2.3 Identification of Bacterial Isolates

The isolates were identified by conventional methods, morphological, cultural, and Gram stain reactions. The Analytical Profile Index (API) 20E test kits by Bio Merieux SA was employed for biochemical identifications. The API kit was prepared according to the manufacturer's specifications.

## 2.4 DNA Extraction and PCR of Fungal Isolates

Fungi were identified using morphological and macroscopic features according to [6]. Extracted fungal DNA was amplified with ITS 1f/ ITS 4r universal primers for fungi according to [8].

## 3. RESULTS AND DISCUSSION

### 3.1 Total Viable Counts of Bacteria Isolate under Ambient and Cold Temperatures

The total bacterial counts from the fresh mushrooms (day 0) at ambient temperature, were 5.8 log cfu/ml. The number increased up to the 7<sup>th</sup> day of storage except at day 5 for *C. indica* (Fig. 1). Continuous storage fruitbodies at ambient temperature, increased the bacterial counts up to 6.4 log cfu/ml, 6.2 log cfu/ml and 6.3 log cfu/ml over a 7-day period in *C. indica*, *P. ostreatus* and *P. tuber-regium* respectively.

This observation is in agreement with [3] who reported that in a 5-day post-harvest storage of sliced unwashed mushrooms, population of *Listeria monocytogenes* increased from 4 log cfu/g to 6.8 log cfu/g. Also, the brown blotch discoloration of *Agaricus bisporus* under refrigeration temperature was reported to be caused by high bacterial populations [9]. It was reported that 60% of fresh mushrooms and dried mushroom products examined in Bangladesh

were contaminated with coliforms, *E coli*, *Salmonella* and the highest bacterial load recorded was  $8.7 \times 10^8$  cfu / gm [10]. Contrary to this observation, a low occurrence of pathogenic bacteria in fresh produce sampled in Norway was reported though the fresh mushrooms harboured non-toxinogenic *Staphylococcus aureus* [11]. Plate count of aerobic mesophilic microorganisms in food is one of the microbial indicators of food quality and large counts are regarded as harmful and reflect existence of favourable conditions for multiplication. The increasing number of aerobic counts showed that continued storage could be dangerous and may have serious consequences for food safety.

Under cold temperature, the increase in the bacterial counts were not so rapid until the 5<sup>th</sup> day (5.9 log cfu/ml - 6.1 log cfu/ml) and 7<sup>th</sup> day of storage (6.0 log cfu/ml - 6.3 log cfu/ml) (Fig. 2). The result agrees with the suggestion of Singh *et al.*, [1] that higher storage temperature can activate tyrosinase and increase microbial growth. This was affirmed by [5] that temperature had significant effect on contaminants of some mushrooms. *Calocybe indica*, the milk mushroom supported the highest growth of microflora which could be due to the succulent nature and high nutrient status of the mushroom. The Pearson Correlation coefficient indicated a strong positive linear correlation existed between the three mushroom samples and is significant at 0.01 level (Table 1).





Fig. 1. Bacterial counts on fruitbodies of three fresh mushrooms stored under ambient temperature (28°C)

**Table 1. Pearson Correlation Coefficient of the three mushroom samples**

		<i>Calocybe indica</i>	<i>P. ostreatus</i>	<i>P. tuber-regium</i>
<i>Calocybe indica</i>	Pearson Correlation	1	.819**	.845**
	Sig. (2-tailed)		.000	.000
	N	16	16	16
<i>P. ostreatus</i>	Pearson Correlation	.819**	1	.996**
	Sig. (2-tailed)	.000		.000
	N	16	16	16
<i>P. tuber regium</i>	Pearson Correlation	.845**	.996**	1
	Sig. (2-tailed)	.000	.000	
	N	16	16	16

\*\* Correlation is significant at the 0.01 level (2-tailed)

**Fig. 2. Bacterial counts on fruitbodies of three fresh mushrooms stored under cold temperature (15°C)**

Freshly harvested mushrooms were reported to harbour approximately 3 log cfu of moulds and 6 log cfu of native yeast per gram of fresh tissue and bacterial populations tend to increase from 7.3 to 8.4 log cfu g<sup>-1</sup> during a 1-day storage period at 4°C [12].

### 3.2 Identification of Bacterial Isolates using API 20E kits

Seven genera of bacteria belonging to the family Enterobacteriaceae were identified by the Analytical Profile Index 20E Kits; among them were *Enterobacter*, *Pseudomonas*, *Chryseobacterium*, *Klebsiella*, *Citrobacter*, *Serratia* and *Cedecea species*. Dominant among them were *Enterobacter* (30%) and *Klebsiella* species (20%). In another study, *Enterobacter* specie was among the eight bacteria isolated from diseased mycelia of *Pleurotus eryngii* [13]. In a similar study, *Enterobacter* and *Pseudomonas* were associated with the substrate used in the cultivation of *P. ostreatus* [14]. Most Enterobacteriaceae are mesophiles

and thrive better under ambient temperature than refrigeration temperature [15]. This was corroborated by [4] that Enterobacteriaceae are slow growers under chill temperatures and become more significant as temperature rises. The involvement of members of Enterobacteriaceae is indicative of environmental contamination as members of this family are enteric organisms spread through the unhygienic handling of foods. These organisms are of a public health concern as they are capable of causing serious ill health. Their presence in food could have serious implications for food safety and is indicative of poor hygiene. This calls for adequate hygiene and good manufacturing practice during production and storage. Mushrooms should be produced and processed observing good hygienic practices (GHP) and consumed soon after harvest. Such hygienic procedures include proper composting of wastes and pasteurization at the right temperature to get rid of most contaminating organisms and observance of good hygiene during harvesting of the mushrooms.

ISOLATE	ONPG	ADH	LDC	ODC	CIT	H <sub>2</sub> S	URE	TDA	IND	VP	GEL	GLU	MAN	INO	SOR	RHA	SAC	MEL	AMY	ARA	OX	NO <sub>2</sub>	N <sub>2</sub>	MOB	M <sub>2</sub> C	OF-F	PROBABLE ORGANISM	SPECIMEN SOURCE
B1	+	+	-	+	+	-	-	-	-	+	-	+	+	-	-	-	+	-	+	-	-	+	+	+	+	+	<i>Cedecea davisae</i>	<i>Pleurotus tuberregium</i> ,
B2	+	+	-	+	+	-	-	-	-	+	-	+	+	+	-	+	+	+	+	+	-	-	+	+	+	+	<i>Enterobacter sakazakii</i>	<i>Pleurotus ostreatus</i>
B3	+	-	+	+	+	-	-	-	-	+	+	+	+	+	-	+	+	+	+	-	-	+	+	+	+	+	<i>Serratia marcescens</i>	<i>Pleurotus ostreatus</i>
B4	+	-	-	+	+	-	-	-	-	-	-	+	+	-	+	-	+	+	+	+	-	-	+	+	+	+	<i>Enterobacter asburiae</i>	<i>Calocybe indica</i>
B5	-	+	-	-	+	-	-	-	-	-	+	+	-	-	-	-	-	-	-	-	+	-	+	+	+	+	<i>Pseudomonas aeruginosa</i>	<i>Pleurotus ostreatus</i>
B6	+	+	-	+	+	-	-	-	-	+	-	+	+	-	+	+	+	+	+	+	+	-	+	+	+	+	<i>Enterobacter cloacae</i>	<i>Pleurotus tuber-regium</i>
B7	+	-	-	-	-	-	-	-	+	-	+	-	-	-	-	-	-	-	-	-	+	+	-	-	+	+	<i>Chryseobacterium meningosepticum</i>	<i>Pleurotus ostreatus</i>
B8	+	-	+	+	+	+	-	+	-	+	-	+	+	+	+	+	+	+	+	+	+	+	-	-	-	+	<i>Klebsiella ornithinolytica</i>	<i>Calocybe indica</i>
B9	+	-	+	-	+	-	+	-	+	+	-	+	+	+	+	+	+	+	+	+	+	-	-	-	+	+	<i>Klebsiella oxytoca</i>	<i>Calocybe indica</i>
B10	+	+	-	+	+	+	-	-	-	-	-	+	-	+	+	+	+	+	+	+	+	-	-	+	+	+	<i>Citrobacter braakii</i>	<i>Calocybe indica</i>

**Image 1. Identification of Microorganism**

Key: ONPG- Ortho-Nitrophenyl-β-galactoside, ADH- Arginine DiHydrolase, LDC- Lysine Decarboxy, ODC- Ornithine DeCarboxylase, CIT- Citrate, H<sub>2</sub>S- Hydrogen Sulphide Production, URE- Urease, TDA- Tryptophane DeAminase, IND- Indole Production, VP-Voges Praskauer, GEL- Gelatinase, GLU- D-Glucose, MAN- D-Mannitol, INO- Inositol, SOR- D-Sorbitol, RHA- L-Rhannose, SAC- Saccharose (D-Sucrose), MEL- D-Melibiose

This is contrary to previous reports that members of *Pseudomonads* were the dominant species [12]; the occurrence of *Pseudomonas* in this work was only 10% and is similar to [16] who reported 8% on vegetables. Bacterial species isolated at ambient temperature were *Pseudomonas aeruginosa*, *Serratia marcescens* and *Chryseobacterium meningosepticum* from *P. ostreatus*; *Cedecea davisae* from *P. tuber-regium* while *Enterobacter asburiae*, *Klebsiella ornithinolytica* and *Klebsiella oxytoca* were isolated from *Calocybe indica*. At cold temperature, *Enterobacter sakazaki*, *Enterobacter cloacae* and *Citrobacter braakii* were isolated from *P. ostreatus*, *P. tuber-regium* and *C. indica* respectively.

### 3.3 Total Viable Counts of Fungal Isolates under Ambient and Cold Temperatures

The fungal counts under ambient temperature ranged from 5.0 log cfu/ml to 5.9 log cfu/ml (Fig. 3) while at cold temperature the range was 5 log

cfu/ml to 5.6 log cfu/ml (Fig. 4). Same trend was observed for fungal counts as the counts at ambient temperature exceeded that of cold storage. Bacterial counts were more than fungal counts and increased with storage. This could be due the rapid generation time of bacteria compared to fungi. *Calocybe indica*, the milk mushroom supported the highest growth of fungi which could be due to the succulent nature and high nutrient status of the mushroom.


### 3.4 Identification of Fungal Isolates by Colony Morphology

Four genera of fungi were isolated from the mushrooms, three fungal isolates were isolated under both ambient and cold temperatures and include *Aspergillus spp.*, *Penicillium spp.*, and *Fusarium spp.* While *Alternaria spp.* was found under ambient condition (Table 2). The dominant fungal species were *Aspergillus* (41%) and *Penicillium* (25%). This observation was corroborated by [13] as *Aspergillus* had the highest frequency of occurrence.

**Table 2. Colony Morphology of isolated fungi from *C. indica*, *P. tuber-regium* and *P. ostreatus***

Mushroom species	Probable organism	Colour	Conidiophore	Phalides	Vesicle	A/R	Isolates
<i>P. ostreatus</i>	<i>Alternaria sp.</i>	Dark brown	Septate	Uniseriate	Zigzag	A	F1
<i>P. ostreatus</i>	<i>Aspergillus sp.</i>	Yellow-green	Glucose	Uniseriate	Round	A	F2
<i>P. ostreatus</i>	<i>Penicillium sp.</i>	Greenish	Downy	Uniseriate	Round	R	F8
<i>P. ostreatus</i>	<i>Aspergillus sp.</i>	Blue-green	smooth	uniseriate	round	R	F12
<i>C. indica</i>	<i>Penicillium sp.</i>	Greenish	Smooth	Uniseriate	Round	A	F3
<i>C. indica</i>	<i>Fusarium sp.</i>	Whitish	Smooth	Uniseriate	Round	A	F4
<i>C. indica</i>	<i>Fusarium sp.</i>	Whitish	Smooth	Uniseriate	Round	R	F5
<i>C. indica</i>	<i>Aspergillus sp.</i>	Yellow-green	Round	Uniseriate	Round	A	F6
<i>C. indica</i>	<i>Aspergillus sp.</i>	Black	Smooth	Uniseriate	Round	R	F7
<i>C. indica</i>	<i>Aspergillus sp.</i>	Blue-green	Septate	Uniseriate	Round	A	F13
<i>P. tuber-regium</i>	<i>Aspergillus sp.</i>	Blue-green	Smooth	Uniseriate	Round	A	F9
<i>P. tuber-regium</i>	<i>Aspergillus sp.</i>	Yellow-green	Smooth	Uniseriate	Round	A	F10
<i>p. tuber-regium</i>	<i>Penicillium sp.</i>	Greenish	Smooth	Uniseriate	Round	R	F11

A/R means ambient /refrigeration Cold temperature



**Fig. 3. Total viable counts of fungal isolates at ambient temperature**




Fig. 4. Total counts of fungal isolates under cold temperature




Fig. 5. PCR Amplification of fungal isolates using ITS1F/ITS4R Universal Primers and 100 bp ladder

m-molecular ladder; -ve- negative control; 1-13- fungal isolates

*Aspergillus*, *Fusarium* and *Penicillium* were among the fungi isolated from the substrates used in cultivation of *P. ostreatus* [13].

### 3.4 Molecular Identification of Fungal Isolates under Ambient and Cold Temperatures

Molecular approaches for identification of fungi included internal transcribed spacer region (ITS)

and had been used for detection of fungi. The PCR fragment sizes of fungal isolates using ITS1f and ITS4r were observed at about 600 bp for all isolates (Fig. 5).

Similar fragment size was obtained by [7] for isolates of *Aspergillus* species.

Storage temperature seems to have an effect on microbial counts as more numbers were observed under ambient temperature.

The implications of these results are that mushrooms could harbour large numbers of pathogenic organisms during post-harvest and storage periods so strict hygiene should be observed during production and storage conditions.

#### 4. CONCLUSION

Many enteric bacteria and potential mycotoxin producing fungi were associated with the three edible mushrooms both at ambient and cold temperatures. The bacteria identified include *Enterobacter asburiae*, *Klebsiella oxytoca*, *Klebsiella ornithinolytica*, *Pseudomonas aeruginosa*, *Serratia marcescens*, *Chryseobacterium meningosepticum*, *Cedecea davisae*, and *Citrobacter braakii* under ambient while under cold storage, *Enterobacter cloacae*, *Enterobacter sakazakii* and *Citrobacter braakii* were identified. The fungal species include *Aspergillus*, *Penicillium*, and *Fusarium* under both temperatures. Only *Alternaria spp.* was isolated under ambient temperature. The practical recommendation is that strict hygiene should be maintained during mushroom cultivation and post-harvest operations.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Singh P, Langowski H, Wani HA, Sangerlaub S. Recent advances in extending the shelf life of fresh *Agaricus* mushrooms: A review. *J Sci of Food Agric*. 2010;90(9):1393-402  
DOI:10.1002/jsfa.3971
- Anyakorah CI, Ojubanire M, Essien RE, Igbo UE. Green synthesis of silver nanoparticles using *Pleurotus ostreatus*. *J Appl Life Sci Int*. 2021: 1-10.  
Available:https://doi.org/10.9734/jalsi%2F2021%2Fv24i1230273
- Chikthimmah N. Microbial ecology of mushroom casing soils and preharvest strategies to enhance safety and quality of fresh mushrooms. A Thesis in Food Science, The Pennsylvania State University. 2006.
- Craig W. Microbial food spoilage: A Major Concern for Food Business Operators; 2020.  
Available: https://www.-newfoofmagazine.com
- Gbolagunte JS. Bacteria associated with cultures of *Psathyrella atroumbonata* (Pegler) and *Schizophyllum commune* (Fr. Ex. Fr). *Afr J Biomed Res*. 2004;7:139-143.
- Pitt JL, Basílico JC, Abarca ML, Lopez C. Mycotoxins and toxigenic fungi. *Med Mycol*. 2000;38:41-46.
- Henry T, Iwen PC, Hinrichs SH. Identification of *Aspergillus* species using internal transcribed spacer regions 1 and 2. *J Clin Microbio*. 2000;38:1510-1515.
- Mahajan PV, Oliveira FAR, Macedo I. Effect of temperature and humidity on the transpiration rate of the whole mushrooms. *J Food Eng*. 2008;84:281-288.
- Kamal AM, Begum F, Khair A. Assessment of Microbiological Quality of Fresh-cut, Processed and Preserved Mushrooms Available in and Around Dhaka City. *Bangladesh J Microbiol*. 2011;27(2):42–45.  
Available:https://doi.org/10.3329/bjm.v27i2.9170
- Johannessen GS, Loncarevic S, Kruse H. Bacteriological analysis of fresh produce in Norway. *Inter J Food Microbiol* 2002;77(3):199-204.
- Chikthimmah N, Beelman R. Microbial Spoilage of Fresh Mushrooms. In book: *Microbiology of Fruits and Vegetables*. 2005:135-158.  
DOI:10.1201/9781420038934.ch6
- Yunjung L, Jae SR, Shanliang S, Won N, Eonmi K, Quy VL *et al.* Isolation of bacteria associated with the king oyster mushroom, *Pleurotus eryngii*. *Mycobiol* 2008;36(1): 13-18.  
DOI: 10.4489/MYCO.2008.36.1.013
- Obire O. 2013. Cultivation of mushroom (*Pleurotus ostreatus*) and the microorganisms associated with the substrate used *J Sci Tech* 2013;8(4): 49-59



14. Brock DT, Thomas K, Brock M, David MW. Basic microbiology with applications. Antibiotics as other chemotherapeutics agents (3<sup>rd</sup> Ed.). 1986:114-115.
15. Desta W, Diriba M. Bacteriological contaminants of fresh vegetable irrigated with Awela River in Ethiopia. Adv Biol. 2016(1):1-11.

© 2022 Anyakorah and Adeniji; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/92636>